



echosens

PUBLICATIONS AND COMMUNICATIONS

MAY 15TH, 2020

FibroScan®
by echosens

<http://www.echosens.com>
communication@echosens.com

GUIDELINES

1. Chinese Foundation for Hepatitis Prevention and Control; Chinese Society of Infectious Disease and Chinese Society of Hepatology; Chinese Medical Association; Liver Disease Committee of Chinese Research Hospital Association. Consensus on clinical application of transient elastography detecting liver fibrosis: a 2018 update. *Zhonghua Gan Zang Bing Za Zhi* 2019;27(3):182-191.
2. Ferraioli, G., V. W. Wong, et al. Liver Ultrasound Elastography: An Update to the World Federation for Ultrasound in Medicine and Biology Guidelines and Recommendations. *Ultrasound in Medicine and Biology* 2018.
3. Thursz, M. EASL Clinical Practice Guidelines: Management of alcohol-related liver disease. *Journal of Hepatology* 2018;69(1):154-181.
4. Chalasani, N., Z. Younossi, et al. The diagnosis and management of nonalcoholic fatty liver disease: Practice guidance from the American Association for the Study of Liver Diseases. *Hepatology* 2018;67(1):328-357.
5. Association Française pour l'étude du Foie, A. Recommandations AFEF pour l'élimination de l'infection par le virus de l'hépatite C en France. - 2018.
6. Horowitz, J. M., I. R. Kamel, et al. ACR Appropriateness Criteria(R) Chronic Liver Disease. *J Am Coll Radiol* 2017;14(11s):S391-s405.
7. Lim, J. K., S. L. Flamm, et al. American Gastroenterological Association Institute Guideline on the Role of Elastography in the Evaluation of Liver Fibrosis. *Gastroenterology* 2017;152(6):1536-1543.
8. Singh, S., A. J. Muir, et al. American Gastroenterological Association Institute Technical Review on the Role of Elastography in Chronic Liver Diseases. *Gastroenterology* 2017;152(6):1544-1577.
9. Blond, E., E. Disse, et al. EASL-EASD-EASO clinical practice guidelines for the management of non-alcoholic fatty liver disease in severely obese people: do they lead to over-referral? *Diabetologia* 2017;60(7):1218-1222.
10. (APASL) Asian-Pacific Association for the Study of the Liver (APASL) consensus guidelines on invasive and non-invasive assessment of hepatic fibrosis: a 2016 update. *Hepatology International* 2017;11(1):1-30.
11. Dietrich, C. F., J. Bamber, et al. EFSUMB Guidelines and Recommendations on the Clinical Use of Liver Ultrasound Elastography, Update 2017 (Short Version). *Ultraschall in der Medizin* 2017;38(4):377-394.
12. (SAGES) (2016). SAGES endorsement statement VCTE, South African Gastroenterology Society (SAGES).
13. Barr, R. G., G. Ferraioli, et al. Elastography Assessment of Liver Fibrosis: Society of Radiologists in Ultrasound Consensus Conference Statement. *Ultrasound Quarterly* 2016;32(2):94-107.
14. Marchesini, G. EASL-EASD-EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *Journal of Hepatology* 2016;64(6):1388-402.
15. Acoustic Radiation Force Impulse Imaging for Diagnosis and Monitoring of Liver Fibrosis in Patients with Hepatitis C: A Review of Diagnostic Accuracy, Clinical Effectiveness, Cost-Effectiveness, and Guidelines. Ottawa ON, 2016 Canadian Agency for Drugs and Technologies in Health.
16. de Franchis, R. and V. I. F. Baveno. Expanding consensus in portal hypertension: Report of the Baveno VI Consensus Workshop: Stratifying risk and individualizing care for portal hypertension. *Journal of Hepatology* 2015;63(3):743-52.
17. International Network for Hepatitis in Substance Users (INHSU). Recommendations for the management of hepatitis C virus infection among people who inject drugs. *International Journal of Drug Policy* 2015;26(10):1028-38.
18. American Association for the Study of Liver Diseases (AALSD). AASLD guidelines for treatment of chronic hepatitis B. *Hepatology* 2015;--.
19. World Federation for Ultrasound in Medicine and Biology (WFUMB). WFUMB Guidelines and Recommendations for Clinical Use of Ultrasound Elastography: Part 1: Basic Principles and Terminology. *Ultrasound in Medicine and Biology* 2015;41(5):1126-1147.
20. World Federation for Ultrasound in Medicine and Biology (WFUMB). WFUMB guidelines and recommendations for clinical use of ultrasound elastography: Part 3: liver. *Ultrasound in Medicine and Biology* 2015;41(5):1161-79.
21. Ferraioli, G., C. Filice, et al. WFUMB guidelines and recommendations for clinical use of ultrasound elastography: Part 3: liver. *Ultrasound in Medicine and Biology* 2015;41(5):1161-79.
22. (WHO) (2015). Guidelines for the prevention, care and treatment of persons with chronic hepatitis B infection, World Health Organization.

23. Non-alcoholic Fatty Liver Disease and Metabolic Syndrome-Position Paper of the Indian National Association for the Study of the Liver, Endocrine Society of India, Indian College of Cardiology and Indian Society of Gastroenterology. *Journal of Clinical and Experimental Hepatology* 2015;5(1):51-68.
24. Authority, H. H. [Diagnosis and treatment of chronic hepatitis B and D. Hungarian national consensus guideline]. *Orvosi Hetilap* 2015;156(Suppl 1):25-35.
25. England, N. Clinical Commissioning Policy Statement: Treatment of chronic Hepatitis C in patients with cirrhosis. NHS Technology Adoption Center 2015;Reference: NHS England B07/P/a.
26. American Association for the Study of Liver Diseases; Infectious Diseases Society of America. Recommendations for Testing, Managing, and Treating Hepatitis C. HCV Guidelines 2015;-.
27. EASL-ALEH Clinical Practice Guidelines: Non-invasive tests for evaluation of liver disease severity and prognosis. *Journal of Hepatology* 2015;63(1):237-64.
28. EASL Recommendations on Treatment of Hepatitis C 2015. *Journal of Hepatology* 2015;63(1):199-236.
29. An update on the management of chronic hepatitis C: 2015 Consensus guidelines from the Canadian Association for the Study of the Liver. *Canadian Journal of Gastroenterology & Hepatology* 2015;29(1):19-34.
30. Authority, H. H. [Diagnosis and treatment of chronic hepatitis B and D. National consensus guideline in Hungary from 12 September 2015]. *Orvosi Hetilap* 2015;156(Suppl 2):25-36.
31. Chinese Society of Hepatology; Chinese Society of Infectious Diseases; Chinese Medical Association. The guideline of prevention and treatment for chronic hepatitis C: a 2015 update. *Chinese journal of clinical hepatology* 2015;31(12):1961-1979.
32. Hungarian Ministry of Health. The chronic inflammation of the liver caused by the hepatitis C virus diagnosis and treatment procedures of financing. *Hungarian Gazette* 2015;(071).
33. (AALSD) (2015). AASLD Guidelines for Treatment of Chronic Hepatitis B, American Association for the Study of Liver Diseases.
34. Organization and provision of prescription of direct action antiviral drugs for the treatment of HCV - chronic hepatitis, cirrhosis and relapse after transplant, and management of patients who have no treatment indication. *Official Gazette of the Region of Sicily* 2015;1(8):61, 62.
35. Chinese Society of Hepatology; Chinese Society of Infectious Diseases; Chinese Medical Association. The guideline of prevention and treatment for chronic hepatitis B: a 2015 update. *Chinese journal of clinical hepatology* 2015;31(12):1941-1960.
36. Romanian Ministry of Health; National Health Insurance House. [Therapeutic protocol corresponding position No. 116 code (LB01B)]. *Official Gazette of Romania* 2015;275/162(Annex 14).
37. Thailand Practice Guideline for Management of Chronic Hepatitis B and C 2015. Bangkok, Thai Association for the Study of the Liver.
38. Ministry of Labour Health and Social Affairs of Georgia. Elimination of hepatitis C program. 2015.
39. Transient Elastography for Assessment of Liver Fibrosis and Steatosis: An Evidence-Based Analysis. *Ontario Health Technology Assessment Series* 2015;15(18):1-45.
40. Transient Elastography for Assessment of Liver Fibrosis and Steatosis: OHTAC Recommendation. *Ontario Health Technology Assessment Series* 2015;--:1-4.
41. Transient Elastography and Controlled Attenuation Parameter for Diagnosing Liver Fibrosis and Steatosis in Ontario: An Economic Analysis. *Ontario Health Technology Assessment Series* 2015;15(19):1-58.
42. Prevention, H. C. f. D. C. (2015). [Guidelines for Therapeutic Intervention in Patients with Virus Infection with Hepatitis C]. Greece, Hellenic Center for Disease Control & Prevention - Ministry of Health.
43. Authority, H. H. [Diagnosis, treatment, and follow-up of hepatitis C-virus related liver disease. Hungarian national consensus guideline]. *Orvosi Hetilap* 2014;155(Suppl 2):3-24.
44. Hunyady, B., J. Gervain, et al. [Diagnosis, treatment, and follow-up of hepatitis C-virus related liver disease. Hungarian national consensus guideline]. *Orvosi Hetilap* 2014;155(Suppl 2):3-24.
45. KASL clinical practice guidelines: management of hepatitis C. *Clinical and Molecular Hepatology* 2014;20(2):89-136.
46. Romanian national guidelines and practical recommendations on liver elastography. *Medical Ultrasonography Journal* 2014;16(2):123-38.
47. JSH Guidelines for the Management of Hepatitis B Virus Infection. *Hepatology Research* 2014;44(Suppl S1):1-58.
48. Australian liver association (ALA) expert consensus recommendations for the use of transient elastography in chronic viral hepatitis.

- Journal of Gastroenterology & Hepatology 2014;-.
49. Latin American Association for the Study of the Liver recommendations on treatment of hepatitis C. Annals of Hepatology 2014;13 Suppl 2:s4-66.
50. (WHO) (2014). Guidelines for the screening, care and treatment of persons with hepatitis C infection, World Health Organization.
51. Notice of the Minister of Health regarding the list of reimbursed drugs, foodstuffs for special nutrition purposes and medical devices. Official Journal of the Minister of Health - Poland 2014;(Annex B.1; Annex B.2):670-680.
52. (ANRS) (2014). [Management of HCV and HBV infected patients]. France, Saint-Denis [France] : HAS (Haute Autorité de Santé).
53. Hellenic Center for Disease Control & Prevention (2014). [Guidelines for Therapeutic Intervention in Patients with Virus Infection with Hepatitis C]. Greece, Hellenic Center for Disease Control & Prevention - Ministry of Health.
54. Brazilian Ministry of Health (2013). Management of patients chronically infected with Hepatitis C (genotype 1) and advanced fibrosis, WHO Regional Office for Europe.
55. (SBU) (2013). Transient elastography with suspected fibrosis and cirrhosis of the liver. Stockholm, 2013 by the Swedish Council on Health Technology Assessment.
56. (NICE) (2013). NICE clinical guideline 165: Hepatitis B (chronic): Diagnosis and management of chronic hepatitis B in children, young people and adults.
57. EFSUMB guidelines and recommendations on the clinical use of ultrasound elastography. Part 2: Clinical applications. Ultraschall in der Medizin 2013;34(3):238-53.
58. KASL clinical practice guidelines: management of nonalcoholic fatty liver disease. Clinical and Molecular Hepatology 2013;19(4):325-48.
59. JSUM ultrasound elastography practice guidelines: liver. Journal of Medical Ultrasonics 2013;40(4):325-357.
60. The chronic inflammation of the liver caused by the hepatitis C virus diagnosis and treatment procedures of financing. Hungarian Gazette 2013;(071):50720, 50721, 50726, 50738.
61. Ministry of Health of Vietnam. Guideline on HCV diagnosis and treatment. Decision no. 4817/QĐ-BYT dated 28/11/2013 by Minister of Ministry of Health 2013.
62. Indications by the Italian Association for the Study of the Liver (AISF) on the use of triple therapy (pegylated alfa interferon + ribavirin + first-generation protease inhibitor) for the treatment of HCV genotype 1 patients. Roma,
- AISF, ASSOCIAZIONE ITALIANA PER LO STUDIO DEL FEGATO.
63. (BASL) (2013). Determination of needs for patient-centered and cost-effective HCV management, Belgian Association for the Study of the Liver.
64. Thai FDA and Bureau Of Drug Control (2013). [National List of Essential Medicine: User guideline Peginterferon alfa type 2a and 2b with Ribavirin Indication Chronic hepatitis C genotype 2,3].
65. Amending the decision on determining main list Croatian Health Insurance. Official Gazette of the Republic of Croatia 2013;(47):114, 120.
66. Minister of Health of the Russian Federation (2013). [Guidelines for the diagnosis and treatment of adult patients with hepatitis C], Minister of Health of the Russian Federation.
67. Brazilian Ministry of Health (2013). [Management of patients chronically infected with Hepatitis C (genotype 1) and advanced fibrosis]. Clinical protocol on treatment of viral Hepatitis and co-infections. Brasília, Brazilian Ministry of Health. **Suplement 1**.
68. Spanish society of liver transplantation (SETH) consensus recommendations on hepatitis C virus and liver transplantation. Liver International 2012;32(5):712-31.
69. Hungarian consensus guideline for the diagnosis and treatment of B, C, and D viral hepatitis. Orvosi Hetilap 2012;153(10):375-94.
70. Portal hypertension: recommendations for evaluation and treatment: consensus document sponsored by the Spanish Association for the Study of the Liver (AEEH) and the Biomedical Research Network Center for Liver and Digestive Diseases(CIBERehd). Gastroenterología y hepatología 2012;35(6):421-50.
71. The diagnosis and management of non-alcoholic fatty liver disease: practice Guideline by the American Association for the Study of Liver Diseases, American College of Gastroenterology, and the American Gastroenterological Association. Hepatology 2012;55(6):2005-23.
72. An update on the management of hepatitis C: consensus guidelines from the Canadian Association for the Study of the Liver. Canadian Journal of Gastroenterology 2012;26(6):359-75.
73. Myers, R. P., A. Ramji, et al. An update on the management of hepatitis C: consensus guidelines from the Canadian Association for the Study of the Liver. Canadian Journal of Gastroenterology 2012;26(6):359-75.
74. Consensus document of the Spanish Association for the Study of the Liver on the treatment of hepatitis B infection (2012).

- Gastroenterología y Hepatología 2012;35(7):512-28.
75. EASL clinical practical guidelines: management of alcoholic liver disease. Journal of Hepatology 2012;57(2):399-420.
76. APASL consensus statements and management algorithms for hepatitis C virus infection. Hepatology International 2012;6(2):409-435.
77. Spanish Ministry of Health (2012). [Criteria and recommendations for the treatment of HCV mono-infected patients]. Madrid - Spain, Ministerio de Sanidad, Servicios Sociales e Igualdad (Spanish Ministry of Health).
78. Spanish Ministry of Health (2012). [Criteria and recommendations for the treatment of HCV-HIV co-infected patients, transplanted patients and pediatric patients]. Madrid - Spain, Ministerio de Sanidad, Servicios Sociales e Igualdad (Spanish Ministry of Health).
79. Italian Association for the Study of the Liver (AISF) (2012). [AISF guidelines to define the appropriateness of centers to manage Hepatitis C genotype 1 patients receiving triple therapy]. Roma, AISF, ASSOCIAZIONE ITALIANA PER LO STUDIO DEL FEGATO.
80. KASL Clinical Practice Guidelines: Management of chronic hepatitis B. Clinical and Molecular Hepatology 2012;18(2):109-162.
81. Revision and update on clinical practice guideline for liver cirrhosis. The Korean Journal of Hepatology 2012;18(1):1-21.
82. World Health Organization (2012). Management of hepatitis B and HIV coinfection. HIV/AIDS treatment and care. Clinical protocols for the WHO European Region. 2012 Revisions, WHO Regional Office for Europe
83. World Health Organization (2012). Management of hepatitis C and HIV co-infection. HIV/AIDS treatment and care. Clinical protocols for the WHO European Region, WHO Regional Office for Europe
84. NHS Technology Adoption Center; Kings College NHS Foundation Trust; North West London Hospitals NHS Trust; East Cheshire NHS Trust. The Use of Ultrasound Elastography for the Detection and Assessment of Liver Fibrosis. A Technology Implementation Study Within Secondary Care. NHS Technology Adoption Center 2012;-.
85. Indonesian Association for the Study of the Liver (2012). [Guidelines Indonesia National consensus (Hepatitis B)]. Indonesia, Indonesian Association for the Study of the Liver.
86. World Gastroenterology Organisation (2012). World Gastroenterology Organisation global guidelines: Nonalcoholic fatty liver disease and nonalcoholic steatohepatitis, World Gastroenterology Organisation.
87. EASL clinical practice guidelines: Management of chronic hepatitis B virus infection. Journal of Hepatology 2011;57(1):167-85.
88. EASL Clinical Practice Guidelines: management of hepatitis C virus infection. Journal of Hepatology 2011;55(2):245-64.
89. The guideline of prevention and treatment for chronic hepatitis B (2010 version). Zhonghua Gan Zang Bing Za Zhi 2011;19(1):13-24.
90. Catalan Society of Gastroenterology (2011). [Consensus document: Diagnosis usefulness of FibroScan®]. Barcelona - Spain, Societat Catalana de Digestología (Catalan Society of Gastroenterology).
91. Spanish Agency for Health Technology Assessment (AETS) (2009). Transient Elastography (FibroScan®) for the diagnosis of liver fibrosis: a systematic review and meta-analysis. Madrid (Spain), AETS (Agencia de Evaluación de Tecnologías Sanitarias) (Spanish Agency for Health Technology Assessment).
92. Liver fibrosis: consensus recommendations of the Asian Pacific Association for the Study of the Liver (APASL). Hepatology International 2009;3(2):323-333.
93. Asian-Pacific consensus statement on the management of chronic hepatitis B: a 2008 update. Hepatology International 2008;2(3):263-83.
94. Haute Autorité de Santé (HAS) (2008). [Recommendations - Diagnosis of uncomplicated cirrhosis]. France, Saint-Denis [France] : HAS (Haute Autorité de Santé).

PUBLICATIONS

In Press

1. Siddiqui, M. S., M. O. Idowu, et al. Diagnostic Performance of Vibration Controlled Transient Elastography in Liver Transplant Recipients. *Clin Gastroenterol Hepatol* 2020; *in press*.
2. Wade, S. D., E. M. Yoshida, et al. Transient Elastography for Monitoring for Hepatotoxicity in Rheumatoid Arthritis Patients on Long-term Methotrexate. *Journal of clinical rheumatology* 2018; *In Press*.

2020

3. Tafur, M., A. Cheung, et al. Risk stratification in primary sclerosing cholangitis: comparison of biliary stricture severity on MRCP versus liver stiffness by MR elastography and vibration-controlled transient elastography. *European Radiology* 2020.
4. Pereira, P. F., V. Von Diemen, et al. Are Noninvasive Methods Comparable to Liver Biopsy in Postoperative Patients After Roux-en-Y Gastric Bypass? *Obesity surgery* 2020.
5. Shiha, G., R. Soliman, et al. A same day "test and treat" model for chronic HCV and HBV infection: Results from two community-based pilot studies in Egypt. *J Viral Hepat.* 2020.
6. Scorletti, E., P. R. Afolabi, et al. Symbiotic Alters Fecal Microbiomes, but not Liver Fat or Fibrosis, in a Randomized Trial of Patients With Non-alcoholic Fatty Liver Disease. *Gastroenterology* 2020.
7. Pourmarzi, D., A. Smirnov, et al. Enablers and barriers for the provision of community-based HCV treatment: A case study of a real-world practice. *Journal of viral hepatitis* 2020.
8. Paternostro, R., J. Pfeiffenberger, et al. Non-invasive diagnosis of cirrhosis and long-term disease monitoring by transient elastography in patients with Wilson disease. *Liver International* 2020.
9. Shima, T., K. Sakai, et al. Diagnostic accuracy of combined biomarker measurements and vibration-controlled transient elastography (VCTE) for predicting fibrosis stage of non-alcoholic fatty liver disease. *Journal of Gastroenterology* 2020;55(1):100-112.
10. Soni, A. and B. K. Tripathi. Correlation of FibroScan with FIB-4 index and APRI for Assessment of Liver Fibrosis in patients with Nonalcoholic Fatty Liver Disease. *J Assoc Physicians India* 2020;68(1):97.
11. Stefanescu, H. and G. Marasco. A novel spleen-dedicated stiffness measurement by FibroScan(R) improves the screening of high-risk oesophageal varices. *Liver Int.* 2020;40(1):175-185.
12. Said, M., R. Eletreby, et al. Fibro-indices versus liver stiffness for prediction of significant fibrosis in hepatitis B virus-infected Egyptian patients; a single-center experience. *Expert Rev Gastroenterol Hepatol.* 2020;1-7.
13. Schleiger, A., M. Salzmann, et al. Severity of Fontan-Associated Liver Disease Correlates with Fontan Hemodynamics. *Pediatr Cardiol* 2020.
14. Sebastiani, G., S. Saeed, et al. Vitamin E is an effective treatment for nonalcoholic steatohepatitis in HIV mono-infected patients. *Aids* 2020;34(2):237-244.
15. Russo, F. P. and A. Zanetto. Hepatitis C virus eradication with direct-acting antiviral improves insulin resistance. *J Viral Hepat.* 2020;27(2):188-194.
16. Sebastiani, G., S. Coccilillo, et al. Application of guidelines for the management of nonalcoholic fatty liver disease in three prospective cohorts of HIV-monoinfected patients. *HIV Medicine* 2020;21(2):96-108.
17. Sterling, R. K., W. C. King, et al. Evaluating Noninvasive Markers to Identify Advanced Fibrosis by Liver Biopsy in HBV/HIV Co-infected Adults. *Hepatology* 2020;71(2):411-421.
18. Zheng, S., M. Tong, et al. Lipid accumulation product independently correlate with hepatic steatosis quantified by controlled attenuation parameter in women with polycystic ovary syndrome. *Endocr Connect* 2020;9(2):154-162.
19. Siddiqui, M. S., M. O. Idowu, et al. Diagnostic Performance of Vibration Controlled Transient Elastography in Liver Transplant Recipients. *Clin Gastroenterol Hepatol* 2020; *in press*.
20. Patel, K. and G. Sebastiani. Limitations of non-invasive tests for assessment of liver fibrosis. *JHEP Rep* 2020;2(2):100067.
21. Shao, L. N., S. T. Zhang, et al. Platelet indices significantly correlate with liver fibrosis in HCV-infected patients. *PLoS One*. 2020;15(1):e0227544.
22. Soliman, H., D. Ziada, et al. Predictors for Fibrosis Regression in Chronic HCV Patients after the Treatment with DAAS: Results of a Real-world Cohort Study. *Endocr Metab Immune Disord Drug Targets* 2020;20(1):104-111.
23. Mantovani, A., A. Taliento, et al. PNPLA3 I148M gene variant and Chronic Kidney Disease in Type 2 Diabetic Patients With NAFLD: Clinical and Experimental Findings. *Liver International* 2020.
24. Foncea, C. G., A. Popescu, et al. Comparative study between pSWE and 2D-SWE techniques integrated in the same ultrasound machine, with Transient

- Elastography as the reference method. *Medical Ultrasonography Journal* 2020;22(1):15-21.
25. Abeysekera, K. W. M., G. S. Fernandes, et al. Prevalence of steatosis and fibrosis in young adults in the UK: a population-based study. *Lancet Gastroenterol Hepatol* 2020;5(3):295-305.
26. Bredin, C., S. Naimimohasses, et al. Development and relative validation of a short food frequency questionnaire for assessing dietary intakes of non-alcoholic fatty liver disease patients. *Eur J Nutr* 2020;59(2):571-580.
27. Mikolasevic, I., V. Lukenda Zanko, et al. Prospective evaluation of non-alcoholic fatty liver disease by elastographic methods of liver steatosis and fibrosis; controlled attenuation parameter and liver stiffness measurements. *Journal of diabetes and its complications* 2020;34(3):107512.
28. Horvath, G., T. Keleti, et al. Neurocognitive functions in patients with hepatitis C infection. *Ideggyogy Sz* 2020;73(1-2):43-49.
29. Liu, W. Y., J. H. Wang, et al. Transient elastography for assessment of fibrosis and steatosis of liver grafts from brain-death donors. *Clinics and Research in Hepatology and Gastroenterology* 2020.
30. Jain, V. and U. Poddar. Utility and accuracy of transient elastography in determining liver fibrosis: a case-control study. *Eur J Pediatr.* 2020.
31. Lian, X. X., Y. P. Sun, et al. [Correlation between intestinal mucosal permeability and prognosis in patients with liver cirrhosis]. *Zhonghua Gan Zang Bing Za Zhi* 2020;28(1):58-63.
32. Han, E., Y. H. Lee, et al. Ipragliflozin Additively Ameliorates Non-Alcoholic Fatty Liver Disease in Patients with Type 2 Diabetes Controlled with Metformin and Pioglitazone: A 24-Week Randomized Controlled Trial. *2020;9(1)*.
33. Kao, W. Y., I. W. Chang, et al. Fibroscan-Based Score to Predict Significant Liver Fibrosis in Morbidly Obese Patients with Nonalcoholic Fatty Liver Disease. *Obesity surgery* 2020.
34. Chuah, K. H., L. L. Lai, et al. Liver stiffness measurement in non-alcoholic fatty liver disease: Two is better than one. *Journal of Gastroenterology & Hepatology* 2020.
35. Casper, M., S. Zimmermann, et al. Risk of chemotherapy-associated liver injury (CALI) in PNPLA3 p.148M allele carriers: Preliminary results of a transient elastography-based study. *Digestive & Liver Disease* 2020;52(1):102-106.
36. Hyun Kim, K., B. Kyung Kim, et al. Sarcopenia assessed using bioimpedance analysis is associated independently with significant liver fibrosis in patients with chronic liver diseases. *European Journal of Gastroenterology & Hepatology* 2020;32(1):58-65.
37. Leong, W. L., L. L. Lai, et al. Comparing point shear wave elastography (ElastPQ) and transient elastography for diagnosis of fibrosis stage in non-alcoholic fatty liver disease. *Journal of Gastroenterology & Hepatology* 2020;35(1):135-141.
38. Inadomi, C., H. Takahashi, et al. Accuracy of the enhanced liver fibrosis (ELF) test and combination of ELF and noninvasive tests for the diagnosis of advanced liver fibrosis in patients with nonalcoholic fatty liver disease. *Hepatology Research* 2020.
39. Dong, T. S., W. Katzka, et al. A Microbial Signature Identifies Advanced Fibrosis in Patients with Chronic Liver Disease Mainly Due to NAFLD. *Sci Rep* 2020;10(1):2771.
40. Eskes, E. C. B., B. Sjouke, et al. Biochemical and imaging parameters in acid sphingomyelinase deficiency: Potential utility as biomarkers. *Mol Genet Metab* 2020.
41. Mueller, S., M. Luderer, et al. Open-label Study with Nalmefene as Needed Use in Alcohol-Dependent Patients with Evidence of Elevated Liver Stiffness and/or Hepatic Steatosis. *Alcohol Alcohol* 2020;55(1):63-70.
42. Costa, P., B. Rudolph, et al. Liver Stiffness Measurement by Vibration Controlled Transient Elastography Does Not Correlate to Hepatic Iron Overload in Children With Sickle Cell Disease. *J Pediatr Hematol Oncol* 2020.
43. Newsome, P. N., M. Sasso, et al. FibroScan-AST (FAST) score for the non-invasive identification of patients with non-alcoholic steatohepatitis with significant activity and fibrosis: a prospective derivation and global validation study. *Lancet Gastroenterol Hepatol* 2020.
44. Bischoff, J., C. Boesecke, et al. Has Increased Rollout of Direct Acting Antiviral Therapy Decreased the Burden of Late Presentation and Advanced Liver Disease in Patients Starting Hepatitis C Virus Therapy in Germany? *Journal of Clinical Gastroenterology* 2020;54(2):192-199.
45. Bluthner, E., J. Bednarsch, et al. Advanced liver function assessment in patients with intestinal failure on long-term parenteral nutrition. *Clinical Nutrition* 2020;39(2):540-547.
46. Cardoso, A. C., C. Cravo, et al. The performance of M and XL probes of FibroScan for the diagnosis of steatosis and fibrosis on a Brazilian nonalcoholic fatty liver disease cohort. *European Journal of Gastroenterology & Hepatology* 2020;32(2):231-238.
47. Da, B. L., P. Surana, et al. The Delta-4 fibrosis score (D4FS): A novel fibrosis score in chronic hepatitis D. *Antiviral Res* 2020;174:104691.
48. da Silva Franco, K. M. V. and W. B. Vieira. Doppler ultrasonography: A non-invasive method used to diagnose and follow up patients with chronic hepatitis C. *Journal of Gastroenterology & Hepatology* 2020;35(2):314-319.

49. Kohla, M. A. S., A. E. Fayoumi, et al. Early fibrosis regression by shear wave elastography after successful direct-acting anti-HCV therapy. *Clin Exp Med* 2020;20(1):143-148.
50. Lipinski, P., P. Szymanska-Rozek, et al. Controlled attenuation parameter and liver stiffness measurements using transient elastography by FibroScan in Gaucher disease. *Mol Genet Metab* 2020;129(2):125-131.
51. Lombardi, R. and L. Airaghi. Liver fibrosis by FibroScan((R)) independently of established cardiovascular risk parameters associates with macrovascular and microvascular complications in patients with type 2 diabetes. *Liver International* 2020;40(2):347-354.
52. Lombardi, R., S. Petta, et al. FibroScan Identifies Patients With Nonalcoholic Fatty Liver Disease and Cardiovascular Damage. *Clin Gastroenterol Hepatol* 2020;18(2):517-519.
53. Majumdar, A., S. Campos, et al. Defining the Minimum Acceptable Diagnostic Accuracy of Noninvasive Fibrosis Testing in Cirrhosis: A Decision Analytic Modeling Study. *Hepatology* 2020;71(2):627-642.
54. Navarro, J. G., S. J. Yang, et al. What are the most important predictive factors for clinically relevant posthepatectomy liver failure after right hepatectomy for hepatocellular carcinoma? *Ann Surg Treat Res*. 2020;98(2):62-71.
55. Gatos, I., P. Drazinos, et al. Comparison of Sound Touch Elastography, Shear Wave Elastography and Vibration-Controlled Transient Elastography in Chronic Liver Disease Assessment using Liver Biopsy as the "Reference Standard". *Ultrasound in Medicine and Biology* 2020;46(4):959-971.
56. Fayed, H. M. and H. S. Mahmoud. The Utility of Retinol-Binding Protein 4 in Predicting Liver Fibrosis in Chronic Hepatitis C Patients in Response to Direct-Acting Antivirals. *Clin Exp Gastroenterol* 2020;13:53-63.
60. Di Mauro, S. and A. Scamporrino. Serum coding and non-coding RNAs as biomarkers of NAFLD and fibrosis severity. *Liver Int*. 2019;39(9):1742-1754.
61. Hamesch, K., M. Mandorfer, et al. Liver Fibrosis and Metabolic Alterations in Adults With alpha-1-antitrypsin Deficiency Caused by the Pi*ZZ Mutation. *Gastroenterology* 2019;157(3):705-719.e18.
62. Kim, D. S., M. Y. Jeon, et al. Influence of hepatic steatosis on the outcomes of patients with chronic hepatitis B treated with entecavir and tenofovir. *Clinical and Molecular Hepatology* 2019;25(3):283-293.
63. Machado, C. M., N. C. Leite, et al. PNPLA3 gene polymorphism in Brazilian patients with type 2 diabetes: A prognostic marker beyond liver disease? *Nutr Metab Cardiovasc Dis* 2019;29(9):965-971.
64. Mahmud, N., S. D. Doshi, et al. Transient elastography reliably estimates liver fibrosis in autoimmune hepatitis. *Clin Exp Hepatol* 2019;5(3):244-249.
65. Mantovani, A., E. Sani, et al. Association between non-alcoholic fatty liver disease and bone turnover biomarkers in post-menopausal women with type 2 diabetes. *Diabetes Metab* 2019;45(4):347-355.
66. Shen, Y., S. D. Wu, et al. The prognostic role of liver stiffness in patients with chronic liver disease: a systematic review and dose-response meta-analysis. *Hepatology International* 2019;13(5):560-572.
67. Siddiqui, M. S., S. Carbone, et al. Prevalence and Severity of Nonalcoholic Fatty Liver Disease Among Caregivers of Patients With Nonalcoholic Fatty Liver Disease Cirrhosis. *Clin Gastroenterol Hepatol* 2019;17(10):2132-2133.
68. Sun, J., Y. Li, et al. Association between abdominal obesity and liver steatosis and fibrosis among patients with chronic hepatitis B measured by Fibroscan. *Experimental and therapeutic medicine* 2019;18(3):1891-1898.
69. Xu, Y., Y. Liu, et al. Comparison of FibroTouch and FibroScan for staging fibrosis in chronic liver disease: Single-center prospective study. *Digestive & Liver Disease* 2019;51(9):1323-1329.
70. Ferraioli, G. and L. B. Soares Monteiro. Ultrasound-based techniques for the diagnosis of liver steatosis. *World Journal of Gastroenterology* 2019;25(40):6053-6062.
71. Oeda, S., H. Takahashi, et al. Accuracy of liver stiffness measurement and controlled attenuation parameter using FibroScan((R)) M/XL probes to diagnose liver fibrosis and steatosis in patients with nonalcoholic fatty liver disease: a multicenter prospective study. *Journal of Gastroenterology* 2019.
72. Sharif, A., Z. Abbas, et al. Effect of Non-alcoholic Fatty Liver Disease on Transaminase Levels and

2019

57. Houck, K. K., A. P. Ifeachor, et al. Pharmacist-driven multidisciplinary pretreatment workup process for hepatitis C care: A novel model for same-day pretreatment workup. *J Am Pharm Assoc* (2003) 2019;59(5):710-716.
58. Lee, J. I., H. W. Lee, et al. Value of controlled attenuation parameter in fibrosis prediction in nonalcoholic steatohepatitis. *World Journal of Gastroenterology* 2019;25(33):4959-4969.
59. Stefanescu, H., C. Rusu, et al. Liver Stiffness Assessed by Ultrasound Shear Wave Elastography from General Electric Accurately Predicts Clinically Significant Portal Hypertension in Patients with Advanced Chronic Liver Disease. *Ultraschall in der Medizin* 2019.

- Transient Elastography in Patients with Chronic Hepatitis B. *Cureus* 2019;11(10):e5995.
73. Lin, Y. J., C. H. Lin, et al. Noninvasive and Convenient Screening of Metabolic Syndrome Using the Controlled Attenuation Parameter Technology: An Evaluation Based on Self-Paid Health Examination Participants. *J Clin Med* 2019;8(11).
74. Jafarov, F., E. Kaya, et al. The diagnostic utility of fibrosis-4 or nonalcoholic fatty liver disease fibrosis score combined with liver stiffness measurement by fibroscan in assessment of advanced liver fibrosis: a biopsy-proven nonalcoholic fatty liver disease study. *European Journal of Gastroenterology & Hepatology* 2019.
75. Kirkegaard-Klitbo, D. M., F. Bendtsen, et al. Poor Concordance Between Liver Stiffness and Noninvasive Fibrosis Scores in HIV Infection Without Viral Hepatitis. *Clin Gastroenterol Hepatol* 2019.
76. Wang, J. H., H. Y. Ou, et al. Usefulness of Controlled Attenuation Parameter in Detecting and Monitoring Hepatic Steatosis with MRI-PDFF as Reference. *Digestive Diseases & Sciences* 2019.
77. Gil-Garcia, A. I., A. Madejon, et al. Prevalence of hepatocarcinoma-related hepatitis B virus mutants in patients in grey zone of treatment. *World Journal of Gastroenterology* 2019;25(38):5883-5896.
78. Roccarina, D., L. logna Prat, et al. Establishing Reliability Criteria for Liver ElastPQ Shear Wave Elastography (ElastPQ-SWE): Comparison Between 10, 5 and 3 Measurements. *Ultraschall in der Medizin* 2019.
79. Alsenbesy, M., M. Rau, et al. A 2-step fast-track elastometry service for advanced workup of nonalcoholic fatty liver disease (NAFLD) patients - single-center real-world experience of outpatient clinical practice. *Zeitschrift für Gastroenterologie* 2019;57(10):1209-1217.
80. Bellan, M. and C. Colletta. Severity of Nonalcoholic Fatty Liver Disease in Type 2 Diabetes Mellitus: Relationship between Nongenetic Factors and PNPLA3/HSD17B13 Polymorphisms. *Diabetes Metab J* 2019;43(5):700-710.
81. Ben Yakov, G. and D. Sharma. Vibration Controlled Transient Elastography (Fibroscan(R)) in sickle cell liver disease - could we strike while the liver is hard? *Br J Haematol.* 2019;187(1):117-123.
82. Davyduke, T. and P. Tandon. Impact of Implementing a "FIB-4 First" Strategy on a Pathway for Patients With NAFLD Referred From Primary Care. *Hepatol Commun* 2019;3(10):1322-1333.
83. Ferraioli, G., L. Maiocchi, et al. Detection of Liver Steatosis With a Novel Ultrasound-Based Technique: A Pilot Study Using MRI-Derived Proton Density Fat Fraction as the Gold Standard. *Clinical and Translational Gastroenterology* 2019;10(10):e00081.
84. Honda, Y., K. Imajo, et al. Autotaxin is a valuable biomarker for the prediction of liver fibrosis in patients with non-alcoholic fatty liver disease. *Hepatology Research* 2019;49(10):1136-1146.
85. Izumi, T., T. Sho, et al. Assessing the risk of hepatocellular carcinoma by combining liver stiffness and the controlled attenuation parameter. *Hepatol Res.* 2019;49(10):1207-1217.
86. Koutsou-Tassopoulou, A., I. Papapostoli-Sklavounou, et al. Hepatic steatosis in patients with acromegaly. 2019;2(4):e00090.
87. Nan, S., Y. Liang, et al. Hepatic Steatosis in Chronic Hepatitis B as Measured by Controlled Attenuation Parameter: Frequency and Distribution. *J Coll Physicians Surg Pak* 2019;29(10):937-941.
88. Ortolan, A., M. Lorenzin, et al. Metabolic syndrome, non-alcoholic fatty liver disease and liver stiffness in psoriatic arthritis and psoriasis patients. *Clinical Rheumatology* 2019;38(10):2843-2850.
89. Staufer, K., E. Halilbasic, et al. Evaluation and comparison of six noninvasive tests for prediction of significant or advanced fibrosis in nonalcoholic fatty liver disease. *United European Gastroenterology Journal* 2019;7(8):1113-1123.
90. Winters, A. C., R. Mittal, et al. A review of the use of transient elastography in the assessment of fibrosis and steatosis in the post-liver transplant patient. *Clin Transplant* 2019;33(10):e13700.
91. Salavrakos, M., H. Piessevaux, et al. Fibroscan Reliably Rules Out Advanced Liver Fibrosis and Significant Portal Hypertension in Alcoholic Patients. *Journal of Clinical Gastroenterology* 2019;53(10):772-778.
92. Mantovani, A., T. Turino, et al. Screening for non-alcoholic fatty liver disease using liver stiffness measurement and its association with chronic kidney disease and cardiovascular complications in patients with type 2 diabetes. *Diabetes Metab* 2019.
93. Sun, D. Q., F. Z. Ye, et al. Higher liver stiffness scores are associated with early kidney dysfunction in patients with histologically proven non-cirrhotic NAFLD. *Diabetes Metab* 2019.
94. Lombardi, R., S. Fargion, et al. Combined use of genetic polymorphisms and elastographic techniques in NAFLD. Fact or fiction? *Curr Pharm Des* 2019.
95. Li, J., X. Q. Dong, et al. Unsatisfying antiviral therapeutic effect in patients with mother-to-child transmitted chronic hepatitis B virus infection: a prospective multi-center clinical study. *Chinese Medical Journal* 2019;132(22):2647-2656.
96. Da, B. L., P. Surana, et al. Vibration-controlled transient elastography for the detection of cirrhosis in chronic hepatitis D infection. *Journal of viral hepatitis* 2019.
97. Sung, J. C., C. Bosh, et al. Hepatitis C cure improved patient-reported outcomes in patients with and without liver fibrosis in a prospective study at a

- large urban medical center. *Journal of viral hepatitis* 2019.
98. Magdaleno-Tapia, J. and C. Valenzuela-Onate. Prevalence of non-alcoholic fatty liver and liver fibrosis in patients with moderate-severe psoriasis: A cross-sectional cohort study. *Australas J Dermatol* 2019.
99. Rosique-Oramas, D., M. Martinez-Castillo, et al. Production of insulin-like growth factor-binding proteins during the development of hepatic fibrosis due to chronic hepatitis C. *Revista de Gastroenterología de Mexico* 2019.
100. Joshita, S. and Y. Yamashita. Clinical utility of FibroScan as a non-invasive diagnostic test for primary biliary cholangitis. *Journal of Gastroenterology & Hepatology* 2019.
101. Mirizzi, A. and I. Franco. Effects of Some Food Components on Non-Alcoholic Fatty Liver Disease Severity: Results from a Cross-Sectional Study. *Nutrients* 2019;11(11).
102. Tellez, L., E. Rodriguez de Santiago, et al. Prevalence, features and predictive factors of liver nodules in Fontan surgery patients: The VALDIG Fonliver prospective cohort. *Journal of Hepatology* 2019.
103. Yoo, J. J., Y. J. Yoo, et al. Correlation of the grade of hepatic steatosis between controlled attenuation parameter and ultrasound in patients with fatty liver: a multi-center retrospective cohort study. *Korean J Intern Med* 2019.
104. Zhang, X., G. L. Wong, et al. Application of transient elastography in nonalcoholic fatty liver disease. *Clinical and Molecular Hepatology* 2019.
105. Surey, J., D. Menezes, et al. From peer-based to peer-led: redefining the role of peers across the hepatitis C care pathway: HepCare Europe. *The Journal of Antimicrobial Chemotherapy* 2019;74(Supplement_5):v17-v23.
106. Ajmera, V. H., E. Cachay, et al. MRI Assessment of Treatment Response in HIV-associated NAFLD: A Randomized Trial of a Stearyl-Coenzyme-A-Desaturase-1 Inhibitor (ARRIVE Trial). *Hepatology* 2019;70(5):1531-1545.
107. Anstee, Q. M., E. J. Lawitz, et al. Noninvasive Tests Accurately Identify Advanced Fibrosis due to NASH: Baseline Data From the STELLAR Trials. *Hepatology* 2019;70(5):1521-1530.
108. Chan, W. K., S. Treeprasertsuk, et al. Optimizing Use of Nonalcoholic Fatty Liver Disease Fibrosis Score, Fibrosis-4 Score, and Liver Stiffness Measurement to Identify Patients With Advanced Fibrosis. *Clin Gastroenterol Hepatol* 2019;17(12):2570-2580.e37.
109. Chanpong, A., N. Angkathunyakul, et al. Late allograft fibrosis in pediatric liver transplant recipients: Assessed by histology and transient elastography. *Pediatr Transplant*. 2019;23(7):e13541.
110. Crescenzi, L., A. Pecoraro, et al. Liver stiffness assessment by transient elastography suggests high prevalence of liver involvement in common variable immunodeficiency. *Digestive & Liver Disease* 2019;51(11):1599-1603.
111. Crossan, C., A. Majumdar, et al. Referral pathways for patients with NAFLD based on non-invasive fibrosis tests: Diagnostic accuracy and cost analysis. *Liver International* 2019;39(11):2052-2060.
112. Lewindon, P. J., M. V. Puertolas-Lopez, et al. Accuracy of Transient Elastography Data Combined With APRI in Detection and Staging of Liver Disease in Pediatric Patients With Cystic Fibrosis. *Clin Gastroenterol Hepatol* 2019;17(12):2561-2569.e5.
113. Mansour, A., M. R. Mohajeri-Tehrani, et al. Risk factors for non-alcoholic fatty liver disease-associated hepatic fibrosis in type 2 diabetes patients. *Acta Diabetologica* 2019;56(11):1199-1207.
114. Sohn, H. and S. Park. Predicting variceal bleeding in patients with biliary atresia. *Scandinavian Journal of Gastroenterology* 2019;54(11):1385-1390.
115. Uhrig, M., J. Mueller, et al. Susceptibility based multiparametric quantification of liver disease: Non-invasive evaluation of steatosis and iron overload. *Magnetic resonance imaging* 2019;63:114-122.
116. Wong, V. W., M. Irles, et al. Unified interpretation of liver stiffness measurement by M and XL probes in non-alcoholic fatty liver disease. *Gut* 2019;68(11):2057-2064.
117. Yilmaz, Y. and F. Eren. Hepatic fibrosis - and not steatosis - is the main determinant of arterial stiffness in non-alcoholic fatty liver disease. *Atherosclerosis* 2019;290:222-223.
118. Hanafy, A. S., W. M. Seleem, et al. Efficacy of a non-invasive model in predicting the cardiovascular morbidity and histological severity in non-alcoholic fatty liver disease. *Diabetes Metab Syndr* 2019;13(3):2272-2278.
119. Xu, M., Y. Li, et al. Interaction of TM6SF2 E167K and PNPLA3 I148M variants in NAFLD in northeast China. *Ann Hepatol* 2019;18(3):456-460.
120. Shimakawa, Y., G. Ndow, et al. Hepatitis B core-related antigen (HBcrAg): an alternative to HBV DNA to assess treatment eligibility in Africa. *Clinical Infectious Diseases* 2019.
121. Buzzetti, E. and A. Hall. Collagen proportionate area is an independent predictor of long-term outcome in patients with non-alcoholic fatty liver disease. *Alimentary Pharmacology & Therapeutics* 2019;49(9):1214-1222.
122. Chromy, D., M. Mandorfer, et al. Prevalence and Predictors of Hepatic Steatosis in Patients with HIV/HCV Coinfection and the Impact of HCV Eradication. *AIDS Patient Care STDS* 2019;33(5):197-206.
123. Colecchia, A., F. Ravaioli, et al. Liver Stiffness Measurement Allows Early Diagnosis of Veno-Occlusive Disease/Sinusoidal Obstruction Syndrome in Adult Patients Who Undergo

- Hematopoietic Stem Cell Transplantation: Results from a Monocentric Prospective Study. *Biology of Blood and Marrow Transplantation* 2019;25(5):995-1003.
124. Darweesh, S. K., R. A. AbdElAziz, et al. Serum cytokeratin-18 and its relation to liver fibrosis and steatosis diagnosed by FibroScan and controlled attenuation parameter in nonalcoholic fatty liver disease and hepatitis C virus patients. *European Journal of Gastroenterology & Hepatology* 2019;31(5):633-641.
125. Eddowes, P. J., M. Sasso, et al. Accuracy of FibroScan Controlled Attenuation Parameter and Liver Stiffness Measurement in Assessing Steatosis and Fibrosis in Patients With Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2019;156(6):1717-1730.
126. Leite, N. C., B. B. Viegas, et al. Efficacy of diacerein in reducing liver steatosis and fibrosis in patients with type 2 diabetes and non-alcoholic fatty liver disease: A randomized, placebo-controlled trial. *Diabetes Obes Metab* 2019;21(5):1266-1270.
127. Mjelle, A. B., A. Mulabecirovic, et al. Normal Liver Stiffness Values in Children: A Comparison of Three Different Elastography Methods. *Journal of Pediatric Gastroenterology and Nutrition* 2019;68(5):706-712.
128. Ravaioli, F. and A. Colecchia. Role of imaging techniques in liver veno-occlusive disease diagnosis: recent advances and literature review. *Expert Review of Gastroenterology & Hepatology* 2019;13(5):463-484.
129. Schmidt, T., D. Schwinge, et al. Th17 cell frequency is associated with low bone mass in primary sclerosing cholangitis. *Journal of Hepatology* 2019;70(5):941-953.
130. Sun, C., X. Xiao, et al. Histologically proven AMA positive primary biliary cholangitis but normal serum alkaline phosphatase: Is alkaline phosphatase truly a surrogate marker? *J Autoimmun* 2019;99:33-38.
131. Rout, G., B. Nayak, et al. Therapy with Oral Directly Acting Agents in Hepatitis C Infection Is Associated with Reduction in Fibrosis and Increase in Hepatic Steatosis on Transient Elastography. *Journal of Clinical and Experimental Hepatology* 2019;9(2):207-214.
132. Zuberi, F. F., B. F. Zuberi, et al. Non-specific impairment of Lung Function on Spirometry in Patients with Chronic Hepatitis-C. *Pak J Med Sci* 2019;35(2):360-364.
133. Safarian, M., S. Mohammadpour, et al. Effect of diet-induced weight loss on cytokeratin-18 levels in overweight and obese patients with liver fibrosis. *Diabetes Metab Syndr* 2019;13(2):989-994.
134. Fitriakusumah, Y. and C. R. A. Lesmana. The role of Small Intestinal Bacterial Overgrowth (SIBO) in Non-alcoholic Fatty Liver Disease (NAFLD) patients evaluated using Controlled Attenuation Parameter (CAP) Transient Elastography (TE): a tertiary referral center experience. 2019;19(1):43.
135. Doyle, M. A., C. Galanakis, et al. Hepatitis C Direct Acting Antivirals and Ribavirin Modify Lipid but not Glucose Parameters. *Cells* 2019;8(3).
136. Bader, R. M., M. M. Jonas, et al. Controlled attenuation parameter: A measure of hepatic steatosis in patients with cystic fibrosis. *Journal of Cystic Fibrosis* 2019;18(2):280-285.
137. Hsu, C., C. Caussy, et al. Magnetic Resonance vs Transient Elastography Analysis of Patients With Nonalcoholic Fatty Liver Disease: A Systematic Review and Pooled Analysis of Individual Participants. *Clin Gastroenterol Hepatol* 2019;17(4):630-637.e8.
138. Kracht, P. A. M., J. E. Arends, et al. REtrieval And cure of Chronic Hepatitis C (REACH): Results of micro-elimination in the Utrecht province. *Liver International* 2019;39(3):455-462.
139. Malin, J. J., C. Boesecke, et al. Liver stiffness regression after successful Hepatitis C treatment is independent of HIV coinfection. *HIV Medicine* 2019;20(3):230-236.
140. Park, H. E., H. Lee, et al. Clinical significance of hepatic steatosis according to coronary plaque morphology: assessment using controlled attenuation parameter. *Journal of Gastroenterology* 2019;54(3):271-280.
141. Silva, M., P. Costa Moreira, et al. Effect of Meal Ingestion on Liver Stiffness and Controlled Attenuation Parameter. *GE Port J Gastroenterol* 2019;26(2):99-104.
142. Soloveva, A., Z. Kobalava, et al. Relationship of Liver Stiffness With Congestion in Patients Presenting With Acute Decompensated Heart Failure. *J Card Fail* 2019;25(3):176-187.
143. Wang, X. K., P. Wang, et al. A study on spleen transient elastography in predicting the degree of esophageal varices and bleeding. *Medicine (Baltimore)* 2019;98(9):e14615.
144. Zhang, Y., Y. Zheng, et al. Comparison of Acoustic Structure Quantification, Transient Elastography (FibroScan) and Histology in Patients with Chronic Hepatitis B and without Moderate to Severe Hepatic Steatosis. *Ultrasound in Medicine and Biology* 2019;45(3):684-692.
145. Hirono, H., K. Watanabe, et al. Practical Use of Transient Elastography in Screening for Nonalcoholic Steatohepatitis in a Japanese Population. *Journal of Clinical and Translational Hepatology* 2019;7(2):127-131.
146. Li, Y., S. Liu, et al. Association between NAFLD and Risk of Colorectal Adenoma in Chinese Han Population. *Journal of Clinical and Translational Hepatology* 2019;7(2):99-105.
147. Bin, Z., W. Jian-Lin, et al. [Clinical value of transient elastography (FibroScan) in detection of

- clonorchiasis]. Zhongguo Xue Xi Chong Bing Fang Zhi Za Zhi 2019;31(3):319-322.
148. Kwon, Y. D. and K. O. Ko. Usefulness of Transient Elastography for Non-Invasive Diagnosis of Liver Fibrosis in Pediatric Non-Alcoholic Steatohepatitis. Journal of Korean medical science 2019;34(23):e165.
149. Mihaila, R. G. Liver stiffness in chronic hepatitis C virus infection. Romanian Journal of Internal Medicine 2019;57(2):85-98.
150. Abdelaziz, A. O., H. Abdelhalim, et al. Liver stiffness measurement changes following hepatocellular carcinoma treatment with percutaneous microwave ablation or transarterial chemoembolization: a cohort study. European Journal of Gastroenterology & Hepatology 2019;31(6):685-691.
151. Aberra, H., H. Desalegn, et al. The WHO guidelines for chronic hepatitis B fail to detect half of the patients in need of treatment in Ethiopia. Journal of Hepatology 2019;70(6):1065-1071.
152. Darand, M., Z. Darabi, et al. Nigella sativa and inflammatory biomarkers in patients with non-alcoholic fatty liver disease: Results from a randomized, double-blind, placebo-controlled, clinical trial. Complement Ther Med 2019;44:204-209.
153. Hong, C. R., S. M. Han, et al. Noninvasive assessment of liver fibrosis in pediatric intestinal failure patients using liver stiffness measurement by Vibration-Controlled Transient Elastography. Journal of pediatric surgery 2019;54(6):1174-1178.
154. Lee, S. J. and S. U. Kim. Noninvasive monitoring of hepatic steatosis: controlled attenuation parameter and magnetic resonance imaging-proton density fat fraction in patients with nonalcoholic fatty liver disease. Expert Review of Gastroenterology & Hepatology 2019;13(6):523-530.
155. Petta, S., V. Wai-Sun Wong, et al. Impact of Obesity and Alanine Aminotransferase Levels on the Diagnostic Accuracy for Advanced Liver Fibrosis of Noninvasive Tools in Patients With Nonalcoholic Fatty Liver Disease. American Journal of Gastroenterology 2019;114(6):916-928.
156. Sanyal, A. J. Past, present and future perspectives in nonalcoholic fatty liver disease. Nature reviews. Gastroenterology & hepatology 2019;16(6):377-386.
157. Stenberg Ribeiro, M., H. Hagstrom, et al. Transient liver elastography in normal pregnancy - a longitudinal cohort study. Scandinavian Journal of Gastroenterology 2019;54(6):761-765.
158. Sun, J., Y. Li, et al. Dynamic Changes of the Aspartate Aminotransferase-to-Platelet Ratio and Transient Elastography in Predicting a Histologic Response in Patients With Chronic Hepatitis B After Entecavir Treatment. Journal of ultrasound in medicine 2019;38(6):1441-1448.
159. Taniguchi, T., T. Ohtani, et al. Liver Stiffness Reflecting Right-Sided Filling Pressure Can Predict Adverse Outcomes in Patients With Heart Failure. JACC Cardiovasc Imaging 2019;12(6):955-964.
160. Vuppulanchi, R., R. Weber, et al. Is Fasting Necessary for Individuals With Nonalcoholic Fatty Liver Disease to Undergo Vibration-Controlled Transient Elastography? American Journal of Gastroenterology 2019;114(6):995-997.
161. Xia, S., X. Ren, et al. A Noninvasive Method-Shear-Wave Elastography Compared With Transient Elastography in Evaluation of Liver Fibrosis in Patients With Chronic Hepatitis B. Ultrasound Quarterly 2019;35(2):147-152.
162. Knop, V., S. C. Neuberger, et al. Intestinal failure-associated liver disease in patients with short bowel syndrome: Evaluation by transient elastography. Nutrition 2019;63-64:134-140.
163. Srivastava, A., S. Jong, et al. Cost-comparison analysis of FIB-4, ELF and fibroscan in community pathways for non-alcoholic fatty liver disease. BMC Gastroenterology 2019;19(1):122.
164. Attallah, A. M., M. S. Albannan, et al. A panel of a mitogenic (PDGF), biochemical (albumin) and demographic (age) parameters for the non-invasive assessment of hepatic fibrosis. Br J Biomed Sci 2019;76(3):105-110.
165. Ballestri, S., F. Nascimbeni, et al. A critical appraisal of the use of ultrasound in hepatic steatosis. Expert Review of Gastroenterology & Hepatology 2019;13(7):667-681.
166. El-Garem, H., M. AbdAllah, et al. DAAs therapy associated with improved hepatic fibrosis in HCV-GT4 patients co-infected with HIV. Expert Rev Gastroenterol Hepatol. 2019;13(7):693-698.
167. Huijbers, A., L. M. Koggel, et al. Systematic Review: Noninvasive Assessments of Intestinal Failure-Associated Liver Disease in the Adult Population. JPEN J Parenter Enteral Nutr 2019;43(5):615-626.
168. Kardashian, A., Y. Ma, et al. High Kynurenone:Tryptophan Ratio Is Associated With Liver Fibrosis in HIV-Monoinfected and HIV/Hepatitis C Virus-Coinfected Women. Open Forum Infectious Diseases 2019;6(7):ofz281.
169. Mak, L. Y., C. H. Lee, et al. Association of adipokines with hepatic steatosis and fibrosis in chronic hepatitis B patients on long-term nucleoside analogue. Liver International 2019;39(7):1217-1225.
170. Shen, F., Y. Q. Mi, et al. Moderate to severe hepatic steatosis leads to overestimation of liver stiffness measurement in chronic hepatitis B patients without significant fibrosis. Alimentary Pharmacology & Therapeutics 2019;50(1):93-102.
171. White, L., A. Azzam, et al. Facilitating treatment of HCV in primary care in regional Australia: closing the access gap. Frontline Gastroenterol 2019;10(3):210-216.

172. Rout, G., S. Kedia, et al. Controlled Attenuation Parameter for Assessment of Hepatic Steatosis in Indian Patients. *Journal of Clinical and Experimental Hepatology* 2019;9(1):13-21.
173. Lopez-Trujillo, M. A., J. M. Olivares-Gazca, et al. Nonalcoholic Fatty Liver Disease and Thrombocytopenia III: Its Association With Insulin Resistance. *Clin Appl Thromb Hemost* 2019;25:1076029619888694.
174. Bastian, W. P., I. Hasan, et al. Gut Microbiota Profiles in Nonalcoholic Fatty Liver Disease and Its Possible Impact on Disease Progression Evaluated with Transient Elastography: Lesson Learnt from 60 Cases. *Case Reports in Gastroenterology* 2019;13(1):125-133.
175. Garcia-Pajares, F., J. Tejedor-Tejada, et al. Efficacy of Direct-acting Antivirals to Improve Clinical Condition, Fibrosis, and Liver Function in Liver Transplant Recipients Infected by Hepatitis C. *Transplantation Proceedings* 2019;51(1):74-76.
176. Gravito-Soares, M., E. Gravito-Soares, et al. Lysosomal Acid Lipase: Can it be a New Non-Invasive Serum Biomarker of Cryptogenic Liver Fibrosis and Cirrhosis? *Ann Hepatol* 2019;18(1):78-88.
177. Sandra, S., C. R. A. Lesmana, et al. Hyperuricemia as an independent risk factor for non-alcoholic fatty liver disease (NAFLD) progression evaluated using controlled attenuation parameter-transient elastography: Lesson learnt from tertiary referral center. *Diabetes Metab Syndr* 2019;13(1):424-428.
178. Kraemer, M. and M. Krawczyk. Increased Circulating VAP-1 Levels Are Associated with Liver Fibrosis in Chronic Hepatitis C Infection. *J Clin Med.* 2019;8(1).
179. Bazerbachi, F., S. Haffar, et al. Range of Normal Liver Stiffness and Factors Associated With Increased Stiffness Measurements in Apparently Healthy Individuals. *Clin Gastroenterol Hepatol* 2019;17(1):54-64.e1.
180. Chow, J. C. and G. L. Wong. Repeating measurements by transient elastography in non-alcoholic fatty liver disease patients with high liver stiffness. *Journal of Gastroenterology & Hepatology* 2019;34(1):241-248.
181. Zhang, Y., C. Wang, et al. Decreased liver stiffness by transient elastography indicates lower incidence of hepatocellular carcinoma in patients with chronic hepatitis B. *Medicine (Baltimore)* 2019;98(3):e13929.
182. Lambert, J. S., R. Murtagh, et al. 'HepCheck Dublin': an intensified hepatitis C screening programme in a homeless population demonstrates the need for alternative models of care. *BMC Infect Dis* 2019;19(1):128.
183. Bandyopadhyay, D., K. Ashish, et al. Role of liver stiffness in prediction of adverse outcomes in heart failure. *J Cardiol* 2019;73(2):185-186.
184. Feng, G., N. He, et al. A simpler diagnostic formula for screening nonalcoholic fatty liver disease. *Clinical Biochemistry* 2019;64:18-23.
185. Ferraioli, G., A. De Silvestri, et al. Evaluation of Inter-System Variability in Liver Stiffness Measurements. *Ultraschall in der Medizin* 2019;40(1):64-75.
186. Guillaud, O., J. Dumortier, et al. Assessment of liver fibrosis by transient elastography (Fibroscan(R)) in patients with A1AT deficiency. *Clinics and Research in Hepatology and Gastroenterology* 2019;43(1):77-81.
187. Jhaveri, K. S., H. Hosseini-Nik, et al. The development and validation of magnetic resonance elastography for fibrosis staging in primary sclerosing cholangitis. *European Radiology* 2019;29(2):1039-1047.
188. Lertnawapan, R. and S. Chonprasertsuk. Association between cumulative methotrexate dose, non-invasive scoring system and hepatic fibrosis detected by Fibroscan in rheumatoid arthritis patients receiving methotrexate. *Int J Rheum Dis* 2019;22(2):214-221.
189. Li, B., L. Zhang, et al. A noninvasive indicator for the diagnosis of early hepatitis B virus-related liver fibrosis. *European Journal of Gastroenterology & Hepatology* 2019;31(2):218-223.
190. Matthews, K., A. MacGilchrist, et al. A nurse-led FibroScan(R) outreach clinic encourages socially deprived heavy drinkers to engage with liver services. *J Clin Nurs* 2019;28(3-4):650-662.
191. Moreno, C., S. Mueller, et al. Non-invasive diagnosis and biomarkers in alcohol-related liver disease. *Journal of Hepatology* 2019;70(2):273-283.
192. So-Armah, K., M. Freiberg, et al. Liver fibrosis and accelerated immune dysfunction (immunosenescence) among HIV-infected Russians with heavy alcohol consumption - an observational cross-sectional study. *BMC Gastroenterology* 2019;20(1):1.
193. Wang, L., M. Zhu, et al. Liver Stiffness Measurement Can Reflect the Active Liver Necroinflammation in Population with Chronic Liver Disease: A Real-world Evidence Study. *Journal of Clinical and Translational Hepatology* 2019;7(4):313-321.
194. Zeng, Q. and Z. Song. Controlled attenuation parameter by vibration-controlled transient elastography for steatosis assessment in members of the public undergoing regular health checkups with reference to magnetic resonance imaging-based proton density fat fraction. 2019.
195. Hamurcu Varol, P., E. Kaya, et al. Role of intensive dietary and lifestyle interventions in the treatment of lean nonalcoholic fatty liver disease patients. *European Journal of Gastroenterology & Hepatology* 2019.
196. Unger, L. W., B. Forstner, et al. Hepatic Steatosis in Lean Patients: Risk Factors and Impact on Mortality. *Digestive Diseases & Sciences* 2019.

197. Nguyen, T. H., R. Wardell, et al. When the liver gets stiff, the tough get moving. *Journal of Gastroenterology & Hepatology* 2019.
198. Nguyen, T. H., R. Wardell, et al. When the liver gets stiff, the tough get moving. *Journal of Gastroenterology & Hepatology* 2019.
199. Cossiga, V., V. Lembo, et al. Berberis aristata, Elaeis guineensis and Coffea canephora Extracts Modulate the Insulin Receptor Expression and Improve Hepatic Steatosis in NAFLD Patients: A Pilot Clinical Trial. *2019;11(12)*.
200. Caussy, C., J. Brissot, et al. Prospective, Same-day, Direct Comparison of CAP With the M vs the XL Probe in Patients With Nonalcoholic Fatty Liver Disease, Using Magnetic Resonance Imaging Proton Density Fat Fraction as the Standard. *Clin Gastroenterol Hepatol* 2019.
201. Piccinni, R., S. G. Rodrigues, et al. Controlled attenuation parameter reflects steatosis in compensated advanced chronic liver disease. *Liver International* 2019.
202. Elkabany, Z. A., R. T. Hamza, et al. Serum visfatin level as a noninvasive marker for nonalcoholic fatty liver disease in children and adolescents with obesity: relation to transient elastography with controlled attenuation parameter. *European Journal of Gastroenterology & Hepatology* 2019.
203. Osaka, T. and Y. Hashimoto. Reduction of Fat to Muscle Mass Ratio Is Associated with Improvement of Liver Stiffness in Diabetic Patients with Non-Alcoholic Fatty Liver Disease. *J Clin Med.* 2019;8(12).
204. Udompap, P., K. Sukonrut, et al. Prospective comparison of transient elastography, point shear wave elastography, APRI and FIB-4 for staging liver fibrosis in chronic viral hepatitis. *Journal of viral hepatitis* 2019.
205. Batman, B., H. Altun, et al. The Effect of Laparoscopic Sleeve Gastrectomy on Nonalcoholic Fatty Liver Disease. *Surg Laparosc Endosc Percutan Tech* 2019;29(6):509-512.
206. Cazzagon, N., S. Lemoinne, et al. The Complementary Value of Magnetic Resonance Imaging and Vibration-Controlled Transient Elastography for Risk Stratification in Primary Sclerosing Cholangitis. *American Journal of Gastroenterology* 2019;114(12):1878-1885.
207. Dhiman, R. K., G. S. Grover, et al. Decentralized care with generic direct-acting antivirals in the management of chronic hepatitis C in a public health care setting. *Journal of Hepatology* 2019;71(6):1076-1085.
208. Jaafar, R. F., A. M. Hajj Ali, et al. Fibroscan and low-density lipoprotein as determinants of severe liver fibrosis in diabetic patients with nonalcoholic fatty liver disease. *European Journal of Gastroenterology & Hepatology* 2019;31(12):1540-1544.
209. Semmler, G., J. Stift, et al. Performance of Controlled Attenuation Parameter in Patients with Advanced Chronic Liver Disease and Portal Hypertension. *Digestive Diseases & Sciences* 2019;64(12):3642-3651.
210. Vallet-Pichard, A., L. Parlati, et al. [Epidemiology of non-alcoholic steatohepatitis. Extent/burden of the problem and its impact on public health]. *Presse Med* 2019;48(12):1459-1467.
211. Zhang, W., R. Huang, et al. Fat Accumulation, Liver Fibrosis, and Metabolic Abnormalities in Chinese Patients With Moderate/Severe Versus Mild Hepatic Steatosis. *Hepatol Commun* 2019;3(12):1585-1597.
212. Harry, S., L. Lee Lai, et al. Volumetric Liver Fat Fraction Determines Grade of Steatosis More Accurately Than Controlled Attenuation Parameter in Patients With Nonalcoholic Fatty Liver Disease. *Clin Gastroenterol Hepatol* 2019.
213. Tu, C. H., J. Li, et al. [Diagnostic value of endoscopic ultrasonography, fibroscan, acoustic radiation pulse imaging, serological index, and their combination for early stage liver cirrhosis]. *Zhonghua Gan Zang Bing Za Zhi* 2019;27(8):615-620.
214. Broussier, T., A. Lannes, et al. Simple blood fibrosis tests reduce unnecessary referrals for specialized evaluations of liver fibrosis in NAFLD and ALD patients. *Clinics and Research in Hepatology and Gastroenterology* 2019.
215. Serag, W. M. and M. M. Mohamed. Determination of liver fibrosis stages in Egyptian chronic hepatitis B patients by a noninvasive tool. *Turk J Med Sci* 2019;49(4):1145-1150.
216. Xu, X. Y., W. S. Wang, et al. Performance of common imaging techniques vs serum biomarkers in assessing fibrosis in patients with chronic hepatitis B: A systematic review and meta-analysis. *World J Clin Cases* 2019;7(15):2022-2037.
217. Arieira, C., S. Monteiro, et al. Transient elastography: should XL probe be used in all overweight patients? *Scandinavian Journal of Gastroenterology* 2019;54(8):1022-1026.
218. Boursier, J., M. Guillaume, et al. New sequential combinations of non-invasive fibrosis tests provide an accurate diagnosis of advanced fibrosis in NAFLD. *Journal of Hepatology* 2019;71(2):389-396.
219. Calza, L., V. Colangeli, et al. Improvement in liver steatosis after the switch from a ritonavir-boosted protease inhibitor to raltegravir in HIV-infected patients with non-alcoholic fatty liver disease. *Infectious diseases* 2019;51(8):593-601.
220. Darweesh, S. K., H. Omar, et al. The clinical usefulness of elastography in the evaluation of nonalcoholic fatty liver disease patients: A biopsy-controlled study. *European Journal of Gastroenterology & Hepatology* 2019;31(8):1010-1016.
221. Elshaarawy, O., J. Mueller, et al. Spleen stiffness to liver stiffness ratio significantly differs between ALD and HCV and predicts disease-specific complications. *JHEP Rep* 2019;1(2):99-106.

- 222.Lai, L. L., W. N. I. Wan Yusoff, et al. Screening for non-alcoholic fatty liver disease in patients with type 2 diabetes mellitus using transient elastography. *Journal of Gastroenterology & Hepatology* 2019;34(8):1396-1403.
- 223.Manka, P., L. Bechmann, et al. Low Free Triiodothyronine Is Associated with Advanced Fibrosis in Patients at High Risk for Nonalcoholic Steatohepatitis. *Digestive Diseases & Sciences* 2019;64(8):2351-2358.
- 224.Naimi, R. M., M. Hvistendahl, et al. Effects of glepaglutide, a novel long-acting glucagon-like peptide-2 analogue, on markers of liver status in patients with short bowel syndrome: findings from a randomised phase 2 trial. *EBioMedicine* 2019;46:444-451.
- 225.Nogami, A., M. Yoneda, et al. Assessment of 10-year changes in liver stiffness using vibration-controlled transient elastography in non-alcoholic fatty liver disease. *Hepatology Research* 2019;49(8):872-880.
- 226.Praktiknjo, M., N. Djayadi, et al. Fibroblast growth factor 21 is independently associated with severe hepatic steatosis in non-obese HIV-infected patients. *Liver International* 2019;39(8):1514-1520.
- 227.Sozio, M. S. Fibroscan for the Rest of Us-How to Incorporate Fibroscan Into Management of Patients With Liver Disease. *Clin Gastroenterol Hepatol* 2019;17(9):1714-1717.
- 228.Srivastava, A., R. Gailer, et al. Prospective evaluation of a primary care referral pathway for patients with non-alcoholic fatty liver disease. *Journal of Hepatology* 2019;71(2):371-378.
- 229.Tomeno, W., K. Imajo, et al. Distribution of liver stiffness in non-alcoholic fatty liver disease with higher fibrosis-4 index than low cut-off index. *Journal of Gastroenterology & Hepatology* 2019;34(8):1411-1416.
- 230.Zama, D., G. Bossu, et al. Longitudinal evaluation of liver stiffness in three pediatric patients with veno-occlusive disease. *Pediatr Transplant*. 2019;23(5):e13456.
- 231.Dazley, J., E. Szabela, et al. The Largest American Study Comparing Transient Elastography and the Aminotransferase-to-platelet Ratio Index Score, Two Noninvasive Tests for Liver Fibrosis Staging in HIV/HCV Coinfected Patients. *J Glob Infect Dis* 2019;11(2):80-82.
- 232.Siddiqui, M. S., R. Vuppalanchi, et al. Vibration-controlled Transient Elastography to Assess Fibrosis and Steatosis in Patients With Nonalcoholic Fatty Liver Disease. *Clinical Gastroenterology & Hepatology* 2019;17(1):156-163.
- 233.Zeng, J., X. Zhang, et al. Feasibility study and reference values of FibroScan 502 with M probe in healthy preschool children aged 5 years. *BMC Pediatr* 2019;19(1):129.
- 234.Baumeler, S., W. Jochum, et al. Controlled attenuation parameter for the assessment of liver steatosis in comparison with liver histology: a single-centre real life experience. *Swiss Medical Weekly* 2019;149:w20077.
- 235.Shin, J., M. J. Kim, et al. Quick assessment with controlled attenuation parameter for hepatic steatosis in children based on MRI-PDFF as the gold standard. *BMC Pediatr* 2019;19(1):112.
- 236.Pu, K., Y. Wang, et al. Diagnostic accuracy of controlled attenuation parameter (CAP) as a non-invasive test for steatosis in suspected non-alcoholic fatty liver disease: a systematic review and meta-analysis. *BMC Gastroenterology* 2019;19(1):51.
- 237.Iogna Prat, L., D. Roccarina, et al. Etiology and Severity of Liver Disease in HIV-Positive Patients With Suspected NAFLD: Lessons From a Cohort With Available Liver Biopsies. *Journal of Acquired Immune Deficiency Syndromes* 2019;80(4):474-480.
- 238.Lemoine, M., L. Assoumou, et al. Diagnostic Accuracy of Noninvasive Markers of Steatosis, NASH, and Liver Fibrosis in HIV-Monoinfected Individuals at Risk of Nonalcoholic Fatty Liver Disease (NAFLD): Results From the ECHAM Study. *Journal of Acquired Immune Deficiency Syndromes* 2019;80(4):e86-e94.
- 239.Agbim, U. and S. K. Asrani. Non-invasive assessment of liver fibrosis and prognosis: an update on serum and elastography markers. *Expert Review of Gastroenterology & Hepatology* 2019;13(4):361-374.
- 240.Berger, A., S. Shili, et al. Liver Stiffness Measurement With FibroScan: Use the Right Probe in the Right Conditions! *Clinical and Translational Gastroenterology* 2019;10(4):e00023.
- 241.Brown, M. A., W. Gellatley, et al. Medical complications of homelessness: a neglected side of men's health. *Internal Medicine Journal* 2019;49(4):455-460.
- 242.Dillon, A., K. Egan, et al. Liver stiffness and thrombin generation in compensated cirrhosis. *Res Pract Thromb Haemost* 2019;3(2):291-297.
- 243.Hanafy, A. S., S. Soliman, et al. Rescue therapy for chronic hepatitis C virus infection after repeated treatment failures: Impact on disease progression and risk of hepatocellular carcinoma. *Hepatology Research* 2019;49(4):377-384.
- 244.Handzlik, G., M. Holecki, et al. Evaluation of metformin therapy using controlled attenuation parameter and transient elastography in patients with non-alcoholic fatty liver disease. *Pharmacol Rep* 2019;71(2):183-188.
- 245.Sansom, S. E., J. Martin, et al. Steatosis Rates by Liver Biopsy and Transient Elastography With Controlled Attenuation Parameter in Clinical Experience of Hepatitis C Virus (HCV) and Human Immunodeficiency Virus/HCV Coinfection in a Large

- US Hepatitis Clinic. Open Forum Infectious Diseases 2019;6(4):ofz099.
- 246.Tan, E. C. and M. S. Tai. Association between non-alcoholic fatty liver disease evaluated by transient elastography with extracranial carotid atherosclerosis in a multiethnic Asian community. *JGH Open* 2019;3(2):117-125.
- 247.Zhou, J., J. Liu, et al. Haptoglobin 2-2 Genotype is Associated with More Advanced Disease in Subjects with Non-Alcoholic Steatohepatitis: A Retrospective Study. *Advances in therapy* 2019;36(4):880-895.
- 248.Appleby, R. N., I. Moghul, et al. Non-alcoholic fatty liver disease is associated with dysregulated bile acid synthesis and diarrhea: A prospective observational study. *PLoS ONE* 2019;14(1):e0211348.
- 249.Barreira, D. P., R. T. Marinho, et al. Hepatitis C Pretreatment Profile and Gender Differences: Cognition and Disease Severity Effects. *Front Psychol* 2019;10:2317.
- 250.Bellan, M. and C. Rigamonti. Liver Stiffness, Not Fat Liver Content, Predicts the Length of QTc Interval in Patients with Chronic Liver Disease. *Gastroenterol Res Pract* 2019;2019:6731498.
- 251.Cantero, I., M. Elorz, et al. Ultrasound/Elastography techniques, lipidomic and blood markers compared to Magnetic Resonance Imaging in non-alcoholic fatty liver disease adults. *Int J Med Sci* 2019;16(1):75-83.
- 252.Carvalho-Furtado, A. C. L., D. M. Carvalho-Louro, et al. Transient Elastography and Controlled Attenuation Parameter (CAP) in the Assessment of Liver Steatosis in Severe Adult Growth Hormone Deficiency. *Front Endocrinol (Lausanne)* 2019;10:364.
- 253.Dold, L., M. J. Nielsen, et al. Circulating levels of PRO-C3 reflect liver fibrosis and liver function in HIV positive patients receiving modern cART. *PLoS One*. 2019;14(7):e0219526.
- 254.Federico, A. and M. Dallio. Evaluation of the Effect Derived from Silybin with Vitamin D and Vitamin E Administration on Clinical, Metabolic, Endothelial Dysfunction, Oxidative Stress Parameters, and Serological Worsening Markers in Nonalcoholic Fatty Liver Disease Patients. 2019;2019:8742075.
- 255.Gorka-Dynysiewicz, J., M. Pazgan-Simon, et al. Pentraxin 3 Detects Clinically Significant Fibrosis in Patients with Chronic Viral Hepatitis C. *BioMed Research International* 2019;2019:2639248.
- 256.Hansen, J. F., K. M. Christiansen, et al. Combining liver stiffness with hyaluronic acid provides superior prognostic performance in chronic hepatitis C. *PLoS ONE* 2019;14(2):e0212036.
- 257.Hussain, A., M. A. Gul, et al. Validation of Novel Fibrosis Index (NFI) for assessment of liver fibrosis: comparison with transient elastography (FibroScan). *BMJ Open Gastroenterol* 2019;6(1):e000316.
- 258.Kamali, L., A. Adibi, et al. Diagnostic Performance of Ultrasonography in Detecting Fatty Liver Disease in Comparison with Fibroscan in People Suspected of Fatty Liver. *Adv Biomed Res* 2019;8:69.
- 259.Koizumi, Y., M. Hirooka, et al. New diagnostic technique to evaluate hepatic steatosis using the attenuation coefficient on ultrasound B mode. *PLoS ONE* 2019;14(8):e0221548.
- 260.Kuttner, C. S., R. Mancina, et al. Four-Week Omega-3 Supplementation in Carriers of the Prosteatotic PNPLA3 p.I148M Genetic Variant: An Open-Label Study. *Lifestyle Genom* 2019;12(1-6):10-17.
- 261.Lee, Y. C., T. H. Hu, et al. The change in liver stiffness, controlled attenuation parameter and fibrosis-4 index for chronic hepatitis C patients with direct-acting antivirals. *PLoS ONE* 2019;14(4):e0214323.
- 262.Li, Y., F. Xiong, et al. Increased Serum Angiotensin II Is a Risk Factor of Nonalcoholic Fatty Liver Disease: A Prospective Pilot Study. *Gastroenterol Res Pract* 2019;2019:5647161.
- 263.Londono, M. C., M. Riveiro-Barciela, et al. Effectiveness, safety/tolerability of OBV/PTV/r +/- DSV in patients with HCV genotype 1 or 4 with/without HIV-1 co-infection, chronic kidney disease (CKD) stage IIIb-V and dialysis in Spanish clinical practice - Vie-KinD study. *PLoS One*. 2019;14(9):e0221567.
- 264.Ou, H., Y. Fu, et al. Association between Smoking and Liver Fibrosis among Patients with Nonalcoholic Fatty Liver Disease. *Canadian Journal of Gastroenterology & Hepatology* 2019;2019:6028952.
- 265.Semmler, G., B. Scheiner, et al. The impact of hepatic steatosis on portal hypertension. *PLoS ONE* 2019;14(11):e0224506.
- 266.Shili-Masmoudi, S., P. Sogni, et al. Increased liver stiffness is associated with mortality in HIV/HCV coinfecting subjects: The French nationwide ANRS CO13 HEPAVIH cohort study. *PLoS ONE* 2019;14(1):e0211286.
- 267.Soleimani, D., G. Ranjbar, et al. Dietary patterns in relation to hepatic fibrosis among patients with nonalcoholic fatty liver disease. *Diabetes Metab Syndr Obes* 2019;12:315-324.
- 268.Somda, S., A. Lebrun, et al. Adaptation of controlled attenuation parameter (CAP) measurement depth in morbidly obese patients addressed for bariatric surgery. *PLoS ONE* 2019;14(5):e0217093.
- 269.Stepanov, Y. M., N. Y. Zavhorodnia, et al. Association of nonalcoholic fatty liver disease with small intestine bacterial overgrowth in obese children. *Wiad Lek* 2019;72(3):350-356.
- 270.Szakacs, Z., B. Eross, et al. Baveno Criteria Safely Identify Patients With Compensated Advanced Chronic Liver Disease Who Can Avoid Variceal Screening Endoscopy: A Diagnostic Test Accuracy Meta-Analysis. *Front Physiol* 2019;10:1028.

- 271.Tantanavipas, S., O. Vallabhakara, et al. Abdominal Obesity as a Predictive Factor of Nonalcoholic Fatty Liver Disease Assessed by Ultrasonography and Transient Elastography in Polycystic Ovary Syndrome and Healthy Women. *BioMed Research International* 2019;2019:9047324.
- 272.Ting, Y. W., S. W. Wong, et al. Metabolic Syndrome Is Associated With Advanced Liver Fibrosis Among Pediatric Patients With Non-alcoholic Fatty Liver Disease. *Front Pediatr* 2019;7:491.
- 273.Yamashima, M., H. Miyaaki, et al. The Long-term Efficacy of Sodium Glucose Co-transporter 2 Inhibitor in Patients with Non-alcoholic Fatty Liver Disease. *Intern Med* 2019;58(14):1987-1992.
- 274.Yoo, H., Y. Kim, et al. Usefulness of noninvasive methods including assessment of liver stiffness by 2-dimensional shear wave elastography for predicting esophageal varices. *Digestive & Liver Disease* 2019;51(12):1706-1712.
- 275.Zeng, D. B., L. Di, et al. The Effect of Splenectomy on the Reversal of Cirrhosis: a Prospective Study. *Gastroenterol Res Pract* 2019;2019:5459427.

2018

- 276.Xia, B., F. Wang, et al. Feasibility and Efficacy of Transient Elastography using the XL probe to diagnose liver fibrosis and cirrhosis: A meta-analysis. *Medicine (Baltimore)* 2018;97(39):e11816.
- 277.Zhang, M., L. Yuan, et al. Human epididymis protein 4 concentration is not associated with liver fibrosis and cirrhosis in a case control study. *Clinica Chimica Acta* 2018;484:213-217.
- 278.Yoon, J. S., Y. R. Lee, et al. Comparison of acoustic radiation force impulse elastography and transient elastography for prediction of hepatocellular carcinoma recurrence after radiofrequency ablation. *European Journal of Gastroenterology & Hepatology* 2018;30(10):1230-1236.
- 279.Zhang, X. Y., J. Qian, et al. [Clinical value of noninvasive method in diagnosing hepatic fibrosis about chronic HBV carriers]. *Zhonghua Gan Zang Bing Za Zhi* 2018;26(5):332-336.
- 280.Wong, G. L. Non-invasive assessments for liver fibrosis: The crystal ball we long for. *Journal of gastroenterology and hepatology* 2018;33(5):1009-1015.
- 281.Zheng, J. P., H. X. Miao, et al. Risk factors for osteoporosis in liver cirrhosis patients measured by transient elastography. *Medicine (Baltimore)* 2018;97(20):e10645.
- 282.Wu, S., Y. Kong, et al. On-treatment changes of liver stiffness at week 26 could predict 2-year clinical outcomes in HBV-related compensated cirrhosis. *Liver International* 2018;38(6):1045-1054.
- 283.Wu, S., Y. Tseng, et al. Evaluation of transient elastography in assessing liver fibrosis in patients

- with advanced schistosomiasis japonica. *Parasitol Int* 2018;67(3):302-308.
- 284.Zhu, Q., W. Wang, et al. Transient Elastography Identifies the Risk of Esophageal Varices and Bleeding in Patients With Hepatitis B Virus-Related Liver Cirrhosis. *Ultrasound Quarterly* 2018;34(3):141-147.
- 285.Younossi, Z., Q. M. Anstee, et al. Global burden of NAFLD and NASH: trends, predictions, risk factors and prevention. *Nature reviews. Gastroenterology & hepatology* 2018;15(1):11-20.
- 286.Wu, H. M., L. Sheng, et al. Performance of transient elastography in assessing liver fibrosis in patients with autoimmune hepatitis-primary biliary cholangitis overlap syndrome. *World Journal of Gastroenterology* 2018;24(6):737-743.
- 287.Yan, L. B., Q. B. Zhang, et al. Serum S100 calcium binding protein A4 improves the diagnostic accuracy of transient elastography for assessing liver fibrosis in hepatitis B. *Clinics and Research in Hepatology and Gastroenterology* 2018;42(1):64-71.
- 288.Yen, Y. H., F. Y. Kuo, et al. Predicting Hepatic Steatosis in Living Liver Donors Via Controlled Attenuation Parameter. *Transplantation Proceedings* 2018;50(10):3533-3538.
- 289.Wong, V. W., G. L. Wong, et al. Beneficial Effects of Lifestyle intervention in non-obese patients with non-alcoholic fatty liver disease. *Journal of Hepatology* 2018;69(6):1349-1356.
- 290.Wu, J. F., C. S. Lee, et al. Transient elastography is useful in diagnosing biliary atresia and predicting prognosis after hepatoperoenterostomy. *Hepatology* 2018;68(2):616-624.
- 291.Wu, S. D., L. L. Liu, et al. Longitudinal monitoring of liver fibrosis status by transient elastography in chronic hepatitis B patients during long-term entecavir treatment. *Clin Exp Med* 2018;18(3):433-443.
- 292.Wu, J. F., S. H. Song, et al. Clinical Predictors of Liver Fibrosis in Patients With Chronic Hepatitis B Virus Infection From Children to Adults. *The Journal of Infectious Diseases* 2018;217(9):1408-1416.
- 293.Wong, G. L. H., R. Kwok, et al. A new screening strategy for varices by liver and spleen stiffness measurement (LSSM) in cirrhotic patients: A randomized trial. *Liver International* 2018;38(4):636-644.
- 294.Wilder, J., S. S. Choi, et al. Vibration-Controlled Transient Elastography for Diagnosing Cirrhosis and Staging Hepatic Fibrosis. *The Journal of the American Medical Association* 2018;320(19):2031-2032.
- 295.Wang, S., W. Zhang, et al. [Correlation between FibroScan, APRI Detecting the degree of esophageal and gastric varices of liver cirrhosis]. *Zhonghua Gan Zang Bing Za Zhi* 2018;26(5):342-346.

296. Wilson, T. G., Y. d'Udekem, et al. Hepatic and renal end-organ damage in the Fontan circulation: A report from the Australian and New Zealand Fontan Registry. *Int J Cardiol* 2018;273:100-107.
297. Vuppalanchi, R., M. S. Siddiqui, et al. Performance characteristics of vibration-controlled transient elastography for evaluation of nonalcoholic fatty liver disease. *Hepatology* 2018;67(1):134-144.
298. Webb, M., A. Zimran, et al. Are transient and shear wave elastography useful tools in Gaucher disease? *Blood Cells, Molecules & Diseases* 2018;68:143-147.
299. Wade, S. D., E. M. Yoshida, et al. Transient Elastography for Monitoring for Hepatotoxicity in Rheumatoid Arthritis Patients on Long-term Methotrexate. *Journal of clinical rheumatology* 2018;In Press.
300. Wasfy, E., G. Elkassas, et al. Predicting esophageal varices in cirrhotic hepatitis C virus patients using noninvasive measurement of insulin resistance variables. *Endocr Metab Immune Disord Drug Targets* 2018;18(6):573-580.
301. Wicher, D., I. Jankowska, et al. Transient Elastography for Detection of Liver Fibrosis in Children With Autosomal Recessive Polycystic Kidney Disease. *Front Pediatr* 2018;6:422.
302. Vinikoor, M. J., E. Sinkala, et al. Chronic hepatitis B virus monoinfection at a university hospital in Zambia. *World Journal of Hepatology* 2018;10(9):622-628.
303. Vinciguerra, T., A. Brunati, et al. Transient elastography for non-invasive evaluation of post-transplant liver graft fibrosis in children. *Pediatric transplantation* 2018;22(2).
304. Trzepizur, W., J. Boursier, et al. Increased liver stiffness in patients with severe sleep apnoea and metabolic comorbidities. *Eur Respir J* 2018;51(6).
305. van Santen, D. K., M. F. Schim van der Loeff, et al. High proportions of liver fibrosis and cirrhosis in an ageing population of people who use drugs in Amsterdam, The Netherlands. *European Journal of Gastroenterology & Hepatology* 2018;30(10):1168-1176.
306. Vuppalanchi, R., K. Mathur, et al. Liver Stiffness Measurements in Patients with Non-cirrhotic Portal Hypertension - The Devil is In the Details. *Hepatology* 2018;68(6):2438-2440.
307. Tsochatzis, E. A. and P. N. Newsome. Non-alcoholic fatty liver disease and the interface between primary and secondary care. *Lancet Gastroenterol Hepatol* 2018;3(7):509-517.
308. Tsai, E. and T. P. Lee. Diagnosis and Evaluation of Nonalcoholic Fatty Liver Disease/Nonalcoholic Steatohepatitis, Including Noninvasive Biomarkers and Transient Elastography. *Clinics in liver disease* 2018;22(1):73-92.
309. Tseng, Y., F. Li, et al. Spleen and liver stiffness for noninvasive assessment of portal hypertension in cirrhotic patients with large esophageal varices. *Journal of Clinical Ultrasound* 2018;46(7):442-449.
310. Tomic, D., W. W. Kemp, et al. Nonalcoholic fatty liver disease: current concepts, epidemiology and management strategies. *European Journal of Gastroenterology & Hepatology* 2018;30(10):1103-1115.
311. Song, J., K. Kim, et al. Imaging Assessment of Hepatic Changes after Fontan Surgery. *Int Heart J* 2018;59(5):1008-1014.
312. Sun, J., Y. Li, et al. Dynamic Changes of the Aspartate Aminotransferase-to-Platelet Ratio and Transient Elastography in Predicting a Histologic Response in Patients With Chronic Hepatitis B After Entecavir Treatment. *Journal of ultrasound in medicine* 2018;38(6):1441-1448.
313. Qin, S., S. Wang, et al. Liver stiffness assessed by transient elastography as a potential indicator of chronic kidney disease in patients with nonalcoholic fatty liver disease. *J Clin Lab Anal* 2018;33(2):e22657.
314. Shimizu, M., K. Suzuki, et al. Evaluation of the effects of dapagliflozin, a sodium-glucose co-transporter-2 inhibitor, on hepatic steatosis and fibrosis using transient elastography in patients with type 2 diabetes and non-alcoholic fatty liver disease. *Diabetes Obes Metab* 2018;21(2):285-292.
315. Pose, E. and P. Gines. Transient elastography for alcoholic liver disease: a step forward. *Lancet Gastroenterol Hepatol* 2018;3(9):589-591.
316. Taneja, S., A. Duseja, et al. Successful treatment of chronic hepatitis C infection with directly acting antivirals in renal transplant recipients. *Nephrology (Carlton)* 2018;23(9):876-882.
317. Ruzman, L., I. Mikolasevic, et al. Advances in diagnosis of chronic liver diseases in pediatric patients. *World journal of pediatrics* 2018;14(6):541-547.
318. Pietsch, V., K. Deterding, et al. Long-term changes in liver elasticity in hepatitis C virus-infected patients with sustained virologic response after treatment with direct-acting antivirals. *United European Gastroenterology Journal* 2018;6(8):1188-1198.
319. Sutton, H., E. Fitzpatrick, et al. Transient Elastography Measurements of Spleen Stiffness as a Predictor of Clinically Significant Varices in Children. *Journal of Pediatric Gastroenterology and Nutrition* 2018;67(4):446-451.
320. Tanner, S., A. Julai, et al. Non-alcoholic steatohepatitis associated with Netherton syndrome. *BMJ Case Rep* 2018;11(1).
321. Petta, S., G. Sebastiani, et al. Noninvasive Prediction of Esophageal Varices by Stiffness and Platelet in Nonalcoholic Fatty Liver Disease Cirrhosis. *Journal of Hepatology* 2018;69(4):878-885.
322. Shimizu, K., Y. Soroidsa, et al. Eradication of hepatitis C virus is associated with the attenuation of

- steatosis as evaluated using a controlled attenuation parameter. *Sci Rep* 2018;8(1):7845.
323. Shah, N., R. Alhusayen, et al. Methotrexate in the Treatment of Moderate to Severe Atopic Dermatitis: A Retrospective Study. *J Cutan Med Surg* 2018;22(5):484-487.
324. Tapper, E. B. and R. Loomba. Noninvasive imaging biomarker assessment of liver fibrosis by elastography in NAFLD. *Nature reviews Gastroenterology & hepatology* 2018;15(5):274-282.
325. Thiele, M., V. Rausch, et al. Controlled attenuation parameter and alcoholic hepatic steatosis: Diagnostic accuracy and role of alcohol detoxification. *Journal of Hepatology* 2018;68(5):1025-1032.
326. Petta, S., V. Di Marco, et al. Prevalence and severity of nonalcoholic fatty liver disease by transient elastography: Genetic and metabolic risk factors in a general population. *Liver International* 2018;38(11):2060-2068.
327. Shimakawa, Y., R. Njie, et al. Development of a simple score based on HBeAg and ALT for selecting patients for HBV treatment in Africa. *Journal of Hepatology* 2018;69(4):776-784.
328. Saroli Palumbo, C., S. Restellini, et al. Screening for Nonalcoholic Fatty Liver Disease in Inflammatory Bowel Diseases: A Cohort Study Using Transient Elastography. *Inflamm Bowel Dis* 2018;25(1):124-133.
329. Song, J., Z. Ma, et al. Comparison of three cut-offs to diagnose clinically significant portal hypertension by liver stiffness in chronic viral liver diseases: a meta-analysis. *European Radiology* 2018;28(12):5221-5230.
330. Piekarska, A., E. K. Berkan, et al. Surveillance for portal hypertension in the course of liver cirrhosis. *Clin Exp Hepatol* 2018;4(2):49-54.
331. Shousha, H. I., R. A. Abdelaziz, et al. Effect of treatment with direct acting antivirals on body mass index and hepatic steatosis in chronic hepatitis C. *Journal of Medical Virology* 2018;90(6):1099-1105.
332. Song, B. G., D. H. Sinn, et al. Additional role of liver stiffness measurement in stratifying residual hepatocellular carcinoma risk predicted by serum biomarkers in chronic hepatitis B patients under antiviral therapy. *European Journal of Gastroenterology & Hepatology* 2018;30(12):1447-1452.
333. Shiha, G., A. M. Metwally, et al. An educate, test, and treat programme towards elimination of hepatitis C infection in Egypt: a community-based demonstration project. *Lancet Gastroenterol Hepatol* 2018;3(11):778-789.
334. Sood, V., R. Khanna, et al. Study of Family Clustering and PNPLA3 Gene Polymorphism in Pediatric Non Alcoholic Fatty Liver Disease. *Indian Pediatrics* 2018;55(7):561-567.
335. Radmard, A. R., F. Khorasanizadeh, et al. Prevalence and Clinical Outcomes of Common Bile Duct Dilation in Patients Who Use Opium. *Am J Med Sci* 2018;356(1):39-46.
336. Taniguchi, T., T. Ohtani, et al. Liver Stiffness Reflecting Right-Sided Filling Pressure Can Predict Adverse Outcomes in Patients With Heart Failure. *JACC Cardiovasc Imaging* 2018;12(6):955-964.
337. Thu Thuy, P. T., C. Bunchorntavakul, et al. Sofosbuvir-ledipasvir with or without ribavirin for chronic hepatitis C genotype-1 and 6: real-world experience in Vietnam. *Antiviral Therapy* 2018;23(5):415-423.
338. Singh, S., A. Facciorusso, et al. Magnitude and Kinetics of Decrease in Liver Stiffness After Antiviral Therapy in Patients With Chronic Hepatitis C: A Systematic Review and Meta-analysis. *Clinical Gastroenterology & Hepatology* 2018;16(1):27-38.e4.
339. Schmidt, T., C. Schmidt, et al. Disease Duration and Stage Influence Bone Microstructure in Patients With Primary Biliary Cholangitis. *Journal of bone and mineral research* 2018;33(6):1011-1019.
340. Sarkar, M., N. Terrault, et al. The Association of Hispanic Ethnicity with Nonalcoholic Fatty Liver Disease in Polycystic Ovary Syndrome. *Curr Opin Gynecol Obstet* 2018;1(1):24-33.
341. Piecha, F., D. Paech, et al. Rapid change of liver stiffness after variceal ligation and TIPS implantation. *American journal of physiology. Gastrointestinal and liver physiology* 2018;314(2):G179-G187.
342. Roccarina, D., M. Rosselli, et al. Elastography methods for the non-invasive assessment of portal hypertension. *Expert Review of Gastroenterology & Hepatology* 2018;12(2):155-164.
343. Runge, J. H., L. P. Smits, et al. MR Spectroscopy-derived Proton Density Fat Fraction Is Superior to Controlled Attenuation Parameter for Detecting and Grading Hepatic Steatosis. *Radiology* 2018;286(2):547-556.
344. Pokora Rodak, A., S. Kiciak, et al. Neutrophil-lymphocyte ratio and mean platelet volume as predictive factors for liver fibrosis and steatosis in patients with chronic hepatitis B. *Ann Agric Environ Med* 2018;25(4):690-692.
345. Ramirez-Velez, R. and M. Izquierdo. Liver Fat Content and Body Fat Distribution in Youths with Excess Adiposity. 2018;7(12).
346. Poujois, A. and F. Woimant. Wilson's disease: A 2017 update. *Clinics and Research in Hepatology and Gastroenterology* 2018;42(6):512-520.
347. Schwarzkopf, K., J. Bojunga, et al. Use of Antiplatelet Agents Is Inversely Associated With Liver Fibrosis in Patients With Cardiovascular Disease. *Hepatol Commun* 2018;2(12):1601-1609.

348. Procopet, B., P. Fischer, et al. Good performance of liver stiffness measurement in the prediction of postoperative hepatic decompensation in patients with cirrhosis complicated with hepatocellular carcinoma. *Medical Ultrasonography Journal* 2018;20(3):272-277.
349. Sporea, I., R. Mare, et al. Comparative study between four ultrasound Shear Waves Elastographic methods for liver fibrosis assessment. *Medical Ultrasonography Journal* 2018;20(3):265-271.
350. Shiha, G. E., S. El-Etreby, et al. Chronic Hepatitis C Patients with Obesity: Do we Need two Operators for Accurate Evaluation of Liver Stiffness? *Ann Hepatol* 2018;17(5):795-801.
351. Seitz, H. K., R. Bataller, et al. Alcoholic liver disease. *Nat Rev Dis Primers* 2018;4(1):16.
352. Scheiner, B., L. Steininger, et al. Controlled attenuation parameter (CAP) does not predict hepatic decompensation in patients with advanced chronic liver disease. *Liver International* 2018;39(1):127-135.
353. Salavrakos, M., H. Piessevaux, et al. Fibroscan Reliably Rules Out Advanced Liver Fibrosis and Significant Portal Hypertension in Alcoholic Patients. *Journal of Clinical Gastroenterology* 2018;53(10):772-778.
354. Rinaldi, L., A. Ascione, et al. Influence of antiviral therapy on the liver stiffness in chronic HBV hepatitis. *Infection* 2018;46(2):231-238.
355. Seto, W. K., R. W. H. Hui, et al. Association Between Hepatic Steatosis, Measured by Controlled Attenuation Parameter, and Fibrosis Burden in Chronic Hepatitis B. *Clinical Gastroenterology & Hepatology* 2018;16(4):575-583 e2.
356. Shida, T., K. Akiyama, et al. Skeletal muscle mass to visceral fat area ratio is an important determinant affecting hepatic conditions of non-alcoholic fatty liver disease. *Journal of Gastroenterology & Hepatology* 2018;53(4):535-547.
357. Sousos, N., E. Sinakos, et al. Deferasirox improves liver fibrosis in beta-thalassaemia major patients. A five-year longitudinal study from a single thalassaemia centre. *British Journal of Haematology* 2018;181(1):140-142.
358. Sutton, H., A. Dhawan, et al. Non-invasive Markers of Portal Hypertension: Appraisal of Adult Experience and Potential Utilisation in Children. *Journal of Pediatric Gastroenterology and Nutrition* 2018;66(4):559-569.
359. Thiele, M., B. S. Madsen, et al. Accuracy of the Enhanced Liver Fibrosis Test vs FibroTest, Elastography, and Indirect Markers in Detection of Advanced Fibrosis in Patients With Alcoholic Liver Disease. *Gastroenterology* 2018;154(5):1369-1379.
360. Qi, X., M. An, et al. Transient Elastography for Significant Liver Fibrosis and Cirrhosis in Chronic Hepatitis B: A Meta-Analysis. *Canadian Journal of Gastroenterology & Hepatology* 2018;2018:3406789.
361. Saraiva, G. N., N. F. do Rosario, et al. Restoring Inflammatory Mediator Balance after Sofosbuvir-Induced Viral Clearance in Patients with Chronic Hepatitis C. *Mediators Inflamm* 2018;2018:8578051.
362. Sirinawasatien, A., T. Techasirioangkun, et al. Prevalence and Determinants of Significant Liver Fibrosis by Vibration-Controlled Transient Elastography in Thai Chronic Hepatitis B Patients. *International journal of hepatology* 2018;2018:4310102.
363. Nguyen-Khac, E., M. Thiele, et al. Non-invasive diagnosis of liver fibrosis in patients with alcohol-related liver disease by transient elastography: an individual patient data meta-analysis. *Lancet Gastroenterol Hepatol* 2018;3(9):614-625.
364. Park, S. J., A. R. Kim, et al. The Efficacy and Safety of Direct-acting Antiviral Treatment for Chronic Hepatitis C Patients: A Single Center Study. *Korean Journal of Gastroenterology* 2018;72(4):197-204.
365. Park, H. E., H. Lee, et al. Clinical significance of hepatic steatosis according to coronary plaque morphology: assessment using controlled attenuation parameter. *Journal of Gastroenterology & Hepatology* 2018;54(3):271-280.
366. Nakagomi, R., R. Tateishi, et al. Liver stiffness measurements in chronic hepatitis C: treatment evaluation and risk assessment. *Journal of gastroenterology and hepatology* 2018;34(5):921-928.
367. Pan, J. J., F. Bao, et al. Morphometry Confirms Fibrosis Regression From Sustained Virologic Response to Direct-Acting Antivirals for Hepatitis C. *Hepatol Commun* 2018;2(11):1320-1330.
368. Perazzo, H., S. W. Cardoso, et al. Predictive factors associated with liver fibrosis and steatosis by transient elastography in patients with HIV mono-infection under long-term combined antiretroviral therapy. *Journal of the International AIDS Society* 2018;21(11):e25201.
369. Mondal, S. A., D. Dutta, et al. Neck Circumference to Height Ratio is a Reliable Predictor of Liver Stiffness and Nonalcoholic Fatty Liver Disease in Prediabetes. *Indian J Endocrinol Metab* 2018;22(3):347-354.
370. Park, H. E., H. Lee, et al. Usefulness of controlled attenuation parameter for detecting increased arterial stiffness in general population. *Digestive & Liver Disease* 2018;50(10):1062-1067.
371. Nickel, F., C. Tapking, et al. Bariatric Surgery as an Efficient Treatment for Non-Alcoholic Fatty Liver Disease in a Prospective Study with 1-Year Follow-up : BariScan Study. *Obesity surgery* 2018;28(5):1342-1350.
372. Omran, D., R. A. Zayed, et al. Evaluating Diagnostic Accuracy of Noninvasive Tests in Assessment of Significant Liver Fibrosis in Chronic Hepatitis C

- Egyptian Patients. *Viral Immunol* 2018;31(4):315-320.
373. Noureddin, M., M. M. Wong, et al. Fatty liver in hepatitis C patients post-sustained virological response with direct-acting antivirals. *World Journal of Gastroenterology* 2018;24(11):1269-1277.
374. Nogueras Lopez, F., A. Lopez Garrido, et al. Therapy With Direct-Acting Antiviral Agents for Hepatitis C in Liver Transplant Recipients. *Transplantation Proceedings* 2018;50(2):631-633.
375. Omar, H., M. Said, et al. Longitudinal assessment of hepatic fibrosis in responders to direct-acting antivirals for recurrent hepatitis C after liver transplantation using noninvasive methods. *Clinical Transplantation* 2018;32(8):e13334.
376. Nacif, L. S., D. C. Paranagua-Vezozzo, et al. HIGHER VALUES IN LIVER ELASTOGRAPHY AND MELD SCORE ARE MORTALITY PREDICTORS ON LIVER TRANSPLANT WAITING LIST. *Brazilian archives of digestive surgery* 2018;31(1):e1360.
377. Mohamed, Z., J. Rwegasha, et al. The hepatitis C cascade of care in people who inject drugs in Dar es Salaam, Tanzania. *Journal of viral hepatitis* 2018;25(12):1438-1445.
378. Olveira, A., L. Dominguez, et al. Persistently altered liver test results in hepatitis C patients after sustained virological response with direct-acting antivirals. *Journal of viral hepatitis* 2018;25(7):818-824.
379. Paul, J., R. V. Venugopal, et al. MEASUREMENT OF CONTROLLED ATTENUATION PARAMETER: A SURROGATE MARKER OF HEPATIC STEATOSIS IN PATIENTS OF NONALCOHOLIC FATTY LIVER DISEASE ON LIFESTYLE MODIFICATION - A PROSPECTIVE FOLLOW-UP STUDY. *Arquivos de Gastroenterologia* 2018;55(1):7-13.
380. Ooi, G. J., A. Ernest, et al. Evaluating feasibility and accuracy of non-invasive tests for nonalcoholic fatty liver disease in severe and morbid obesity. *International journal of obesity* 2018;42(11):1900-1911.
381. Nobili, V. and P. Socha. Pediatric Nonalcoholic Fatty Liver Disease: Current Thinking. *Journal of Pediatric Gastroenterology and Nutrition* 2018;66(2):188-192.
382. Ogasawara, N., M. Kobayashi, et al. Serial changes in liver stiffness and controlled attenuation parameter following direct-acting antiviral therapy against hepatitis C virus genotype 1b. *Journal of Medical Virology* 2018;90(2):313-319.
383. Ooi, G. J., S. Mgaieth, et al. Systematic review and meta-analysis: non-invasive detection of non-alcoholic fatty liver disease related fibrosis in the obese. *Obesity reviews* 2018;19(2):281-294.
384. Patel, P. J., X. Banh, et al. Underappreciation of non-alcoholic fatty liver disease by primary care clinicians: limited awareness of surrogate markers of fibrosis. *Internal Medicine Journal* 2018;48(2):144-151.
385. Pateu, E., F. Oberti, et al. The noninvasive diagnosis of esophageal varices and its application in clinical practice. *Clinics and Research in Hepatology and Gastroenterology* 2018;42(1):6-16.
386. Paul, J., R. V. Venugopal, et al. Effects of lifestyle modification on liver enzyme and Fibroscan in Indian patients with non-alcoholic fatty liver disease. *Gastroenterology Report* 2018;6(1):49-53.
387. Ozturk, A., J. R. Grajo, et al. Quantitative Hepatic Fat Quantification in Non-alcoholic Fatty Liver Disease Using Ultrasound-Based Techniques: A Review of Literature and Their Diagnostic Performance. *Ultrasound in Medicine and Biology* 2018;44(12):2461-2475.
388. Nascimbeni, F., E. Cassinero, et al. Prevalence and predictors of liver fibrosis evaluated by vibration controlled transient elastography in type 1 Gaucher disease. *Mol Genet Metab* 2018;125(1-2):64-72.
389. Ojha, R. P., B. R. MacDonald, et al. Different Cutpoints for Transient Elastography Lead to Different Associations With Cirrhosis. *Clinical Gastroenterology & Hepatology* 2018;16(8):1359-1360.
390. Patel, P. J., F. Hossain, et al. Controlled attenuation parameter in NAFLD identifies risk of suboptimal glycaemic and metabolic control. *Journal of diabetes and its complications* 2018;32(8):799-804.
391. Perez-Latorre, L., M. Sanchez-Conde, et al. Prognostic value of liver stiffness in HIV/HCV-Coinfected patients with decompensated cirrhosis. *BMC Infectious Diseases* 2018;18(1):170.
392. Mohr, R., C. Boesecke, et al. Return-to-health effect of modern combined antiretroviral therapy potentially predisposes HIV patients to hepatic steatosis. *Medicine (Baltimore)* 2018;97(17):e0462.
393. Mulabecirovic, A., A. B. Mjelle, et al. Repeatability of shear wave elastography in liver fibrosis phantoms-Evaluation of five different systems. *PLoS ONE* 2018;13(1):e0189671.
394. Mulabecirovic, A., A. B. Mjelle, et al. Liver elasticity in healthy individuals by two novel shear-wave elastography systems-Comparison by age, gender, BMI and number of measurements. *PLoS ONE* 2018;13(9):e0203486.
395. Neesgaard, B., M. Ruhwald, et al. Determination of anti-HCV and quantification of HCV-RNA and IP-10 from dried blood spots sent by regular mail. *PLoS ONE* 2018;13(7):e0201629.
396. Papagianni, M. and K. Tziomalos. Non-Alcoholic Fatty Liver Disease in Patients with HIV Infection. *AIDS reviews* 2018;20(3):171-173.
397. Moctezuma-Velazquez, C., F. Saffioti, et al. Non-invasive Prediction of High-risk Varices in Patients with Primary Biliary Cholangitis and Primary Sclerosing Cholangitis. *American Journal of Gastroenterology* 2018;114(3):446-452.

398. Mazur, R., M. Celmer, et al. Clinical applications of spleen ultrasound elastography - a review. *J Ultrason* 2018;18(72):37-41.
399. McDonald, N., P. J. Eddowes, et al. Multiparametric magnetic resonance imaging for quantitation of liver disease: a two-centre cross-sectional observational study. *Sci Rep* 2018;8(1):9189.
400. McCombe, G., D. Swan, et al. Integrated Hepatitis C Care for People Who Inject Drugs (Heplink): Protocol for a Feasibility Study in Primary Care. *JMIR Res Protoc* 2018;7(6):e149.
401. Meijer, B., C. K. van Everdingen, et al. Transient elastography to assess liver stiffness in patients with inflammatory bowel disease. *Digestive & Liver Disease* 2018;50(1):48-53.
402. Mazzola, G., L. Adamoli, et al. Suboptimal performance of APRI and FIB-4 in ruling out significant fibrosis and confirming cirrhosis in HIV/HCV co-infected and HCV mono-infected patients. *Infection* 2018;47(3):409-415.
403. Matthias, A. T., A. N. R. Fernandopulle, et al. Survey on knowledge of non-alcoholic fatty liver disease (NAFLD) among doctors in Sri Lanka: a multicenter study. *BMC Res Notes* 2018;11(1):556.
404. Mendes, L. C., P. A. Ferreira, et al. Controlled attenuation parameter for steatosis grading in chronic hepatitis C compared with digital morphometric analysis of liver biopsy: impact of individual elastography measurement quality. *European Journal of Gastroenterology & Hepatology* 2018;30(8):959-966.
405. Mendes, L. C., P. A. Ferreira, et al. Elastogram quality assessment score in vibration-controlled transient elastography: Diagnostic performance compared to digital morphometric analysis of liver biopsy in chronic hepatitis C. *Journal of viral hepatitis* 2018;25(4):335-343.
406. Miyake, T., S. Yoshida, et al. Ipragliflozin Ameliorates Liver Damage in Non-alcoholic Fatty Liver Disease. *Open medicine* 2018;13:402-409.
407. Fouad, H., A. Mohamed, et al. The Effectiveness of Ledipasvir/Sofosbuvir in Youth with Genotype 4 Hepatitis C Virus; A Single Egyptian Center Study. *Pediatr Infect Dis J* 2018.
408. Garg, G., V. K. Dixit, et al. Impact of Direct Acting Antiviral Drugs in Treatment Naïve HCV Cirrhosis on Fibrosis and Severity of Liver Disease: A Real Life Experience from a Tertiary Care Center of North India. *Journal of Clinical and Experimental Hepatology* 2018;8(3):241-249.
409. Hasan, E. M., R. A. Abd Al Aziz, et al. Genetic Variants in nicotinamide-N-methyltransferase (NNMT) gene are related to the stage of non-alcoholic fatty liver disease diagnosed by controlled attenuation parameter (CAP)-fibroscan. *Journal of gastrointestinal and liver diseases* : JGLD 2018;27(3):265-272.
410. Marasco, G., A. Colecchia, et al. Role of liver and spleen stiffness in predicting the recurrence of hepatocellular carcinoma after resection. *Journal of Hepatology* 2018;70(3):440-448.
411. Knop, V., W. P. Hofmann, et al. Estimation of liver fibrosis by non-commercial serum markers in comparison to transient elastography in patients with chronic hepatitis C virus infection receiving direct acting antiviral treatment. *Journal of viral hepatitis* 2018;26(2):224-230.
412. Mantovani, A., E. Sani, et al. Association between non-alcoholic fatty liver disease and bone turnover biomarkers in post-menopausal women with type 2 diabetes. *Diabetes & Metabolism* 2018;45(4):347-355.
413. Labenz, C., Y. Huber, et al. Predictors of advanced fibrosis in non-cirrhotic non-alcoholic fatty liver disease in Germany. *Alimentary Pharmacology & Therapeutics* 2018;48(10):1109-1116.
414. Giannini, E. G., M. Crespi, et al. Improvement in HCV patients with advanced, compensated liver disease after SVR to antivirals. *European Journal of Clinical Investigation* 2018;49(3):e13056.
415. Garcia-Broncano, P., L. M. Medrano, et al. Dysregulation of the Immune System in HIV/HCV-Coinfected Patients According to Liver Stiffness Status. *Cells* 2018;7(11).
416. Harris, R., T. R. Card, et al. The XL probe: A luxury or a necessity? Risk stratification in an obese community cohort using transient elastography. *United European Gastroenterology Journal* 2018;6(9):1372-1379.
417. Liu, L. Z., J. Sun, et al. Improvements in the management of chronic hepatitis B virus infection. *Expert Review of Gastroenterology & Hepatology* 2018;12(11):1153-1166.
418. Maponga, T. G., M. I. Andersson, et al. HBV and HIV viral load but not microbial translocation or immune activation are associated with liver fibrosis among patients in South Africa. *BMC Infectious Diseases* 2018;18(1):214.
419. Kamal, N., P. Surana, et al. Liver disease in patients with cystic fibrosis. *Current Opinion in Gastroenterology* 2018;34(3):146-151.
420. Kobayashi, N., H. Iijima, et al. Changes in liver stiffness and steatosis among patients with hepatitis C virus infection who received direct-acting antiviral therapy and achieved sustained virological response. *European Journal of Gastroenterology & Hepatology* 2018;30(5):546-551.
421. Kwak, M. S., G. E. Chung, et al. Clinical implications of controlled attenuation parameter in a health check-up cohort. *Liver International* 2018;38(5):915-923.
422. Li, Q., L. Chen, et al. Changes of FibroScan, APRI, and FIB-4 in chronic hepatitis B patients with significant liver histological changes receiving 3-

- year entecavir therapy. *Clin Exp Med* 2018;18(2):273-282.
- 423.Li, J., R. C. Mao, et al. A novel noninvasive index for the prediction of moderate to severe fibrosis in chronic hepatitis B patients. *Digestive & Liver Disease* 2018;50(5):482-489.
- 424.Li, C., R. Li, et al. Progress in non-invasive detection of liver fibrosis. *Cancer Biol Med* 2018;15(2):124-136.
- 425.Heo, J. Y., B. K. Kim, et al. Combination of Transient Elastography and an Enhanced Liver Fibrosis Test to Assess the Degree of Liver Fibrosis in Patients with Chronic Hepatitis B. *Gut Liver* 2018;12(2):190-200.
- 426.Gominon, A. L., E. Frison, et al. Assessment of Liver Disease Progression in Cystic Fibrosis Using Transient Elastography. *Journal of Pediatric Gastroenterology and Nutrition* 2018;66(3):455-460.
- 427.Liang, X., Q. Xie, et al. Interpretation of liver stiffness measurement-based approach for the monitoring of hepatitis B patients with antiviral therapy: A 2-year prospective study. *Journal of viral hepatitis* 2018;25(3):296-305.
- 428.Lv, S., S. Jiang, et al. Noninvasive Quantitative Detection Methods of Liver Fat Content in Nonalcoholic Fatty Liver Disease. *Journal of Clinical and Translational Hepatology* 2018;6(2):217-221.
- 429.Hsu, C., C. Caussy, et al. Magnetic Resonance vs Transient Elastography Analysis of Patients With Nonalcoholic Fatty Liver Disease: A Systematic Review and Pooled Analysis of Individual Participants. *Clinical Gastroenterology & Hepatology* 2018;17(4):630-637.
- 430.Matsui, N., K. Imajo, et al. Magnetic resonance elastography increases usefulness and safety of non-invasive screening for esophageal varices. *Journal of gastroenterology and hepatology* 2018;33(12):2022-2028.
- 431.Gamil, M., M. Alboraei, et al. Novel scores combining AFP with non-invasive markers for prediction of liver fibrosis in chronic hepatitis C patients. *Journal of Medical Virology* 2018;90(6):1080-1086.
- 432.Iacob, S., R. Cerban, et al. 100% sustained virological response and fibrosis improvement in real-life use of direct acting antivirals in genotype-1b recurrent hepatitis C following liver transplantation. *Journal of gastrointestinal and liver diseases : JGLD* 2018;27(2):139-144.
- 433.Lindner, F., R. Muhlberg, et al. Predictive value of liver and spleen stiffness in advanced alcoholic cirrhosis with refractory ascites. *Zeitschrift für Gastroenterologie* 2018;56(6):561-568.
- 434.Mandorfer, M., T. Bucsics, et al. Liver disease in adults with alpha1-antitrypsin deficiency. *United European Gastroenterology Journal* 2018;6(5):710-718.
- 435.Jhaveri, K. S., H. Hosseini-Nik, et al. The development and validation of magnetic resonance elastography for fibrosis staging in primary sclerosing cholangitis. *European Radiology* 2018;29(2):1039-1047.
- 436.Gao, Y., J. Zheng, et al. Liver Fibrosis with Two-dimensional US Shear-Wave Elastography in Participants with Chronic Hepatitis B: A Prospective Multicenter Study. *Radiology* 2018;289(2):407-415.
- 437.Keller, S., J. Sedlacik, et al. Prospective comparison of diffusion-weighted MRI and dynamic Gd-EOB-DTPA-enhanced MRI for detection and staging of hepatic fibrosis in primary sclerosing cholangitis. *European Radiology* 2018;29(2):818-828.
- 438.Hedenstierna, M., A. Nangarhari, et al. Cirrhosis, high age and high body mass index are risk factors for persisting advanced fibrosis after sustained virological response in chronic hepatitis C. *Journal of viral hepatitis* 2018;25(7):802-810.
- 439.Lee, J., M. Y. Kim, et al. The gamma-glutamyl transferase to platelet ratio and the FIB-4 score are noninvasive markers to determine the severity of liver fibrosis in chronic hepatitis B infection. *Br J Biomed Sci* 2018;75(3):128-132.
- 440.Lee, C. K., P. D. Mitchell, et al. Validation of Transient Elastography Cut Points to Assess Advanced Liver Fibrosis in Children and Young Adults: The Boston Children's Hospital Experience. *The Journal of pediatrics* 2018;198:84-89 e2.
- 441.Galaski, J., L. Schulz, et al. Discordance in steatosis classification between liver biopsy and transient elastography for high controlled attenuation parameter (CAP) values. *Zeitschrift für Gastroenterologie* 2018;56(1):36-42.
- 442.Garg, H., S. Aggarwal, et al. Utility of transient elastography (fibroscan) and impact of bariatric surgery on nonalcoholic fatty liver disease (NAFLD) in morbidly obese patients. *Surg Obes Relat Dis* 2018;14(1):81-91.
- 443.Herrmann, E., V. de Ledinghen, et al. Assessment of biopsy-proven liver fibrosis by two-dimensional shear wave elastography: An individual patient data-based meta-analysis. *Hepatology* 2018;67(1):260-272.
- 444.Hui, R. W. H., W. K. Seto, et al. Inverse relationship between hepatic steatosis and hepatitis B viremia: Results of a large case-control study. *Journal of viral hepatitis* 2018;25(1):97-104.
- 445.Kamarajah, S. K., W. K. Chan, et al. Repeated liver stiffness measurement compared with paired liver biopsy in patients with non-alcoholic fatty liver disease. *Hepatology International* 2018;12(1):44-55.
- 446.Lopez, J. J., N. Perez-Alvarez, et al. Optimal Use of Transient Elastography and Acoustic Radiation Force Impulse to Stage Liver Fibrosis in HIV/HCV-Coinfected Patients in Clinical Practice. *Journal of ultrasound in medicine* 2018;37(1):113-121.
- 447.Ibrahim, M., I. Mostafa, et al. New developments in managing variceal bleeding. *Gastroenterology* 2018;154(7):1964-1969.

- 448.Harman, D. J., S. D. Ryder, et al. Obesity and type 2 diabetes are important risk factors underlying previously undiagnosed cirrhosis in general practice: a cross-sectional study using transient elastography. *Alimentary Pharmacology & Therapeutics* 2018;47(4):504-515.
- 449.Kim, B. S., Y. S. Seo, et al. Reduced risk of hepatocellular carcinoma by achieving a subcirrhotic liver stiffness through antiviral agents in hepatitis B virus-related advanced fibrosis or cirrhosis. *Journal of gastroenterology and hepatology* 2018;33(2):503-510.
- 450.Leung, J. C., T. C. Loong, et al. Invasive and non-invasive assessment of portal hypertension. *Hepatology International* 2018;12(Suppl 1):44-55.
- 451.Lin, Z., J. Liang, et al. Diverse correlations between fibrosis-related factors and liver stiffness measurement by transient elastography in chronic hepatitis B. *European Journal of Gastroenterology & Hepatology* 2018;30(2):217-225.
- 452.Lai, L. L., W. Yusoff, et al. Screening for non-alcoholic fatty liver disease in patients with type 2 diabetes mellitus using transient elastography. *Journal of gastroenterology and hepatology* 2018;34(8):1396-1403.
- 453.logna Prat, L., D. Roccarina, et al. Aetiology and severity of liver disease in HIV positive patients with suspected NAFLD: lessons from a cohort with available liver biopsies. *Journal of Acquired Immune Deficiency Syndromes* 2018.
- 454.Mahfouz, R. A., M. Gouda, et al. Interatrial septal fat thickness and left atrial stiffness are mechanistic links between nonalcoholic fatty liver disease and incident atrial fibrillation. *Echocardiography* 2018;36(2):249-256.
- 455.Kiatbumrung, R., N. Chuaypen, et al. The Association of PNPLA3, COX-2 and DHCR7 Polymorphisms with Advanced Liver Fibrosis in Patients with HCV Mono- Infection and HCV/HIV Co-Infection. *Asian Pacific Journal of Cancer Prevention* 2018;19(8):2191-2197.
- 456.Jiang, W., S. Huang, et al. Diagnostic accuracy of point shear wave elastography and transient elastography for staging hepatic fibrosis in patients with non-alcoholic fatty liver disease: a meta-analysis. *BMJ Open* 2018;8(8):e021787.
- 457.Kim, D. W., H. M. Yoon, et al. Diagnostic Performance of Ultrasound Elastography for Evaluating Portal Hypertension in Children: A Systematic Review and Meta-analysis. *Journal of ultrasound in medicine* 2018;38(3):747-759.
- 458.Hwang, J. Y., H. M. Yoon, et al. Diagnostic Performance of Transient Elastography for Liver Fibrosis in Children: A Systematic Review and Meta-Analysis. *AJR Am J Roentgenol* 2018;211(5):W257-W266.
- 459.Leoni, S., F. Tovoli, et al. Current guidelines for the management of non-alcoholic fatty liver disease: A systematic review with comparative analysis. *World Journal of Gastroenterology* 2018;24(30):3361-3373.
- 460.Ishikawa, T., R. Sasaki, et al. Liver stiffness measured by transient elastography as predictor of prognoses following portosystemic shunt occlusion. *Journal of gastroenterology and hepatology* 2018;34(1):215-223.
- 461.Fujiwara, Y., H. Kuroda, et al. The B-Mode Image-Guided Ultrasound Attenuation Parameter Accurately Detects Hepatic Steatosis in Chronic Liver Disease. *Ultrasound in Medicine and Biology* 2018;44(11):2223-2232.
- 462.Margini, C., G. Murgia, et al. Prognostic Significance of Controlled Attenuation Parameter in Patients With Compensated Advanced Chronic Liver Disease. *Hepatol Commun* 2018;2(8):929-940.
- 463.He, C. Y., Y. Lyu, et al. [Diagnostic value of transient elastography for diagnosis of idiopathic non-cirrhotic portal hypertension]. *Zhonghua Gan Zang Bing Za Zhi* 2018;26(4):310-312.
- 464.Garcia-Saenz-de-Sicilia, M. and A. Duarte-Rojo. Hepatic Steatosis and its Effects on Fibrosis in Patients With Chronic Hepatitis B Virus Infection. *Clinical Gastroenterology & Hepatology* 2018;16(4):491-494.
- 465.Jansen, C., P. Moller, et al. Increase in liver stiffness after transjugular intrahepatic portosystemic shunt is associated with inflammation and predicts mortality. *Hepatology* 2018;67(4):1472-1484.
- 466.Jeon, M. Y., H. W. Lee, et al. Feasibility of dynamic risk prediction for hepatocellular carcinoma development in patients with chronic hepatitis B. *Liver International* 2018;38(4):676-686.
- 467.Karlas, T., D. Petroff, et al. Impact of controlled attenuation parameter on detecting fibrosis using liver stiffness measurement. *Alimentary Pharmacology & Therapeutics* 2018;47(7):989-1000.
- 468.Kamal, S., K. K. Aldossari, et al. Clinicopathological and immunological characteristics and outcome of concomitant coeliac disease and non-alcoholic fatty liver disease in adults: a large prospective longitudinal study. *BMJ Open Gastroenterol* 2018;5(1):e000150.
- 469.Kawagishi, N., G. Suda, et al. Liver steatosis and dyslipidemia after HCV eradication by direct acting antiviral agents are synergistic risks of atherosclerosis. *PLoS ONE* 2018;13(12):e0209615.
- 470.Kelly, M. L., S. M. Riordan, et al. Capacity of non-invasive hepatic fibrosis algorithms to replace transient elastography to exclude cirrhosis in people with hepatitis C virus infection: A multi-centre observational study. *PLoS ONE* 2018;13(2):e0192763.
- 471.Kosasih, S., W. Zhi Qin, et al. Relationship between Serum Cytokeratin-18, Control Attenuation Parameter, NAFLD Fibrosis Score, and Liver Steatosis in Nonalcoholic Fatty Liver Disease.

- International journal of hepatology 2018;2018:9252536.
472. Erman, A., A. Sathya, et al. Estimating chronic hepatitis C prognosis using transient elastography-based liver stiffness: A systematic review and meta-analysis. *Journal of viral hepatitis* 2018;25(5):502-513.
473. Fabbri, G., I. Mastorosa, et al. Liver stiffness reduction and serum fibrosis score improvement in HIV/hepatitis C virus-coinfected patients treated with direct-acting antivirals. *HIV Medicine* 2018;19(8):578-584.
474. Ester, C., R. Cerban, et al. The Role of Beta-7 Integrin and Carbonic Anhydrase IX in Predicting the Occurrence of de Novo Nonalcoholic Fatty Liver Disease in Liver Transplant Recipients. *Chirurgia (Bucur)* 2018;113(4):534-541.
475. El Sagheer, G., E. Soliman, et al. Study of changes in lipid profile and insulin resistance in Egyptian patients with chronic hepatitis C genotype 4 in the era of DAAs. *Libyan J Med* 2018;13(1):1435124.
476. Facciorusso, A., H. A. Garcia Perdomo, et al. Systematic review with meta-analysis: Change in liver stiffness during anti-viral therapy in patients with hepatitis B. *Digestive & Liver Disease* 2018;50(8):787-794.
477. Ferraioli, G., A. De Silvestri, et al. Adherence to quality criteria improves concordance between transient elastography and ElastPQ for liver stiffness assessment-A multicenter retrospective study. *Digestive & Liver Disease* 2018;50(10):1056-1061.
478. El-Gohary, M. and M. Moore. Local care and treatment of liver disease (LOCATE) - A cluster-randomized feasibility study to discover, assess and manage early liver disease in primary care. *PLoS One*. 2018;13(12):e0208798.
479. Fabrellas, N., R. Hernandez, et al. Prevalence of hepatic steatosis as assessed by controlled attenuation parameter (CAP) in subjects with metabolic risk factors in primary care. A population-based study. *PLoS ONE* 2018;13(9):e0200656.
480. Fath-Elbab, H. K., E. Ahmed, et al. Event-related evoked potential versus clinical tests in assessment of subclinical cognitive impairment in chronic hepatitis C virus. *Egypt J Neurol Psychiatr Neurosurg* 2018;54(1):35.
481. Di Marco, V., A. Alberti, et al. Optimising management of patients with hepatitis C virus in the age of direct-acting antivirals: results of a Delphi consensus. *European Review for Medical and Pharmacological Sciences* 2018;22(20):7024-7033.
482. Demir, M., O. Deyneli, et al. Screening for hepatic fibrosis and steatosis in Turkish patients with type 2 diabetes mellitus: A transient elastography study. *The Turkish journal of gastroenterology* 2018;30(3):266-270.
483. Dharmalingam, M. and P. G. Yamasandhi. Nonalcoholic Fatty Liver Disease and Type 2 Diabetes Mellitus. *Indian J Endocrinol Metab* 2018;22(3):421-428.
484. Chin, C. Y., C. T. Hsu, et al. Monitoring of liver stiffness by transient elastography during the treatment of Gaucher disease. *Pediatr Neonatol* 2018;60(2):221-223.
485. Chan, W. K., N. R. Nik Mustapha, et al. Can the same controlled attenuation parameter cut-offs be used for M and XL probes for diagnosing hepatic steatosis? *Journal of gastroenterology and hepatology* 2018;33(10):1787-1794.
486. Conti, F., C. Serra, et al. Assessment of Liver Fibrosis with Elastography Point Quantification vs Other Non-Invasive Methods. *Clinical Gastroenterology & Hepatology* 2018;17(3):510-517.
487. Dimzova, M., I. Kondova-Topuzovska, et al. Noninvasive Biomarkers in Assessment of Liver Fibrosis in Patients with HBeAg Negative Chronic Hepatitis B. *Open Access Maced J Med Sci* 2018;6(6):1052-1058.
488. Chow, J. C., G. L. Wong, et al. Repeating measurements by transient elastography in nonalcoholic fatty liver disease patients with high liver stiffness. *Journal of gastroenterology and hepatology* 2018.
489. Dillon, A., Z. Galvin, et al. Transient elastography can stratify patients with Child-Pugh A cirrhosis according to risk of early decompensation. *European Journal of Gastroenterology & Hepatology* 2018;30(12):1434-1440.
490. Cristina, S. J. L., C. M. Marta, et al. Characterization and evaluation of liver fibrosis grade in patients with chronic hepatitis B virus infection and normal transaminases. *Clinical and Molecular Hepatology* 2018;24(4):384-391.
491. Cheng, F., H. Cao, et al. Meta-analysis of the accuracy of transient elastography in measuring liver stiffness to diagnose esophageal varices in cirrhosis. *MEDICINE* 2018;97(28):e11368.
492. Degasperi, E., A. Aghemo, et al. 12 weeks ombitasvir/paritaprevir-ritonavir+ribavirin achieve high SVR rates in HCV-4 patients with advanced fibrosis. *Digestive & Liver Disease* 2018;50(7):703-706.
493. Chen, Y. P., X. M. Hu, et al. Stepwise application of fibrosis index based on four factors, red cell distribution width-platelet ratio, and aspartate aminotransferase-platelet ratio for compensated hepatitis B fibrosis detection. *Journal of gastroenterology and hepatology* 2018;33(1):256-263.
494. Choi, Y. Comment on: utility of transient elastography (fibroscan) and impact of bariatric surgery on nonalcoholic fatty liver disease (NAFLD) in morbidly obese patients. *Surgery for obesity and related diseases* 2018;14(1):91-92.

- 495.Chan, J., N. Gogela, et al. Direct-Acting Antiviral Therapy for Chronic HCV Infection Results in Liver Stiffness Regression Over 12 Months Post-treatment. *Digestive Diseases & Sciences* 2018;63(2):486-492.
- 496.Chi, M., J. Hofmann, et al. Prevalence and Clinical Correlates of Chronic Hepatitis E Infection in German Renal Transplant Recipients With Elevated Liver Enzymes. *Transplant Direct* 2018;4(2):e341.
- 497.Chromy, D., P. Schwabl, et al. Non-invasive liver fibrosis assessment and HCV treatment initiation within a systematic screening program in HIV/HCV coinfected patients. *Wiener Klinische Wochenschrift* 2018;130(3-4):105-114.
- 498.Desalegn, H., H. Aberra, et al. Treatment of chronic hepatitis B in sub-Saharan Africa: 1-year results of a pilot program in Ethiopia. *BMC Medicine* 2018;16(1):234.
- 499.Choudhuri, G., S. Chaudhari, et al. Etiological Patterns, Liver Fibrosis Stages and Prescribing Patterns of Hepato-Protective Agents in Indian Patients with Chronic Liver Disease. *J Assoc Physicians India* 2018;66(12):58-63.
- 500.Clark, V. C., G. Marek, et al. Clinical and Histologic Features of Adults with Alpha-1 Antitrypsin Deficiency in a Non-Cirrhotic Cohort. *Journal of Hepatology* 2018;69(6):1357-1364.
- 501.Colecchia, A., F. Ravaioli, et al. A combined model based on spleen stiffness measurement and Baveno VI criteria to rule out high-risk varices in advanced chronic liver disease. *Journal of Hepatology* 2018;69(2):308-317.
- 502.D'Ambrosio, R., A. Aghemo, et al. Persistence of hepatocellular carcinoma risk in hepatitis C patients with a response to IFN and cirrhosis regression. *Liver International* 2018;38(8):1459-1467.
- 503.Di Lascio, N., C. Avigo, et al. Steato-Score: Non-Invasive Quantitative Assessment of Liver Fat by Ultrasound Imaging. *Ultrasound in Medicine and Biology* 2018;44(8):1585-1596.
- 504.Dong, H., C. Xu, et al. The combination of 5 serum markers compared to FibroScan to predict significant liver fibrosis in patients with chronic hepatitis B virus. *Clinica Chimica Acta* 2018;483:145-150.
- 505.Cheng, P. N., Y. C. Chiu, et al. The characteristics of residents with unawareness of hepatitis C virus infection in community. *PLoS ONE* 2018;13(2):e0193251.
- 506.Cheng, H. S. and M. Rademaker. Monitoring methotrexate-induced liver fibrosis in patients with psoriasis: utility of transient elastography. *Psoriasis (Auckl)* 2018;8:21-29.
- 507.Cheng, P. N., H. C. Chiu, et al. Comparison of FIB-4 and transient elastography in evaluating liver fibrosis of chronic hepatitis C subjects in community. *PLoS ONE* 2018;13(11):e0206947.
- 508.Delicou, S., K. Maragkos, et al. Transient Elastography (TE) is a Useful Tool for Assessing the Response of Liver Iron Chelation in Sickle Cell Disease Patients. *Mediterr J Hematol Infect Dis* 2018;10(1):e2018049.
- 509.Caussy, C., J. Chen, et al. Association Between Obesity and Discordance in Fibrosis Stage Determination by Magnetic Resonance vs Transient Elastography in Patients With Nonalcoholic Liver Disease. *Clinical Gastroenterology & Hepatology* 2018;16(12):1974-1982.
- 510.Castelnovo, B., F. Mubiru, et al. Antiretroviral treatment Long-Term (ALT) cohort: a prospective cohort of 10 years of ART-experienced patients in Uganda. *BMJ Open* 2018;8(2):e015490.
- 511.Castera, L. Diagnosis of non-alcoholic fatty liver disease/non-alcoholic steatohepatitis: Non-invasive tests are enough. *Liver International* 2018;38 Suppl 1:67-70.
- 512.Caussy, C., M. H. Alquiraish, et al. Optimal threshold of controlled attenuation parameter with MRI-PDFF as the gold standard for the detection of hepatic steatosis. *Hepatology* 2018;67(4):1348-1359.
- 513.Cardoso, A. C., R. M. Perez, et al. Prevalence and predictive factors of moderate/severe liver steatosis in chronic hepatitis C (CHC) infected patients evaluated with controlled attenuation parameter (CAP). *Journal of viral hepatitis* 2018;25(11):1244-1250.
- 514.Cao, L. H., F. M. Lu, et al. Study on the relationship between insulin growth factor 1 and liver fibrosis in patients with chronic hepatitis C with type 2 diabetes mellitus. *J Cell Biochem* 2018;119(11):9513-9518.
- 515.Bucsics, T., B. Grasl, et al. Point Shear Wave Elastography for Non-invasive Assessment of Liver Fibrosis in Patients with Viral Hepatitis. *Ultrasound in Medicine and Biology* 2018;44(12):2578-2586.
- 516.Bazerbachi, F., S. Haffar, et al. Range of Normal Liver Stiffness and Factors Associated With Increased Stiffness Measurements in Apparently Healthy Individuals. *Clinical Gastroenterology & Hepatology* 2018;17(1):54-64.
- 517.Bloom, S., W. Kemp, et al. Liver stiffness measurement in the primary care setting detects high rates of advanced fibrosis and predicts liver-related events in hepatitis C. *Journal of Hepatology* 2018;69(3):575-583.
- 518.Buechter, M., J. Thimm, et al. Liver Maximum Capacity: A Novel Test to Accurately Diagnose Different Stages of Liver Fibrosis. *Digestion* 2018;100(1):45-54.
- 519.Cai, S., Z. Ou, et al. Risk factors associated with liver steatosis and fibrosis in chronic hepatitis B patient with component of metabolic syndrome. *United European Gastroenterology Journal* 2018;6(4):558-566.

- 520.Bordbar, M., N. Shakibazad, et al. Effect of ursodeoxycholic acid and vitamin E in the prevention of liver injury from methotrexate in pediatric leukemia. *The Turkish journal of gastroenterology* 2018;29(2):203-209.
- 521.Calanchini, M., A. Moolla, et al. Liver biochemical abnormalities in Turner syndrome: A comprehensive characterization of an adult population. *Clin Endocrinol (Oxf)* 2018;89(5):667-676.
- 522.Caballeria, L., G. Pera, et al. High Prevalence of Liver Fibrosis Among European Adults With Unknown Liver Disease: A Population-Based Study. *Clinical Gastroenterology & Hepatology* 2018;16(7):1138-1145 e5.
- 523.Bachofner, J., P. V. Valli, et al. Excellent outcome of direct antiviral treatment for chronic hepatitis C in Switzerland. *Swiss Medical Weekly* 2018;148:w14560.
- 524.Bellan, M., P. P. Sainaghi, et al. Gas6 as a predictor of esophageal varices in patients affected by hepatitis C virus related-chronic liver disease. *Biomark Med* 2018;12(1):27-34.
- 525.Buechter, M., P. Manka, et al. Spleen stiffness is positively correlated with HVPG and decreases significantly after TIPS implantation. *Digestive & Liver Disease* 2018;50(1):54-60.
- 526.Berzigotti, A., G. Ferraioli, et al. Novel ultrasound-based methods to assess liver disease: The game has just begun. *Digestive & Liver Disease* 2018;50(2):107-112.
- 527.Begovac, J., J. Krznaric, et al. Successful treatment of genotype 3 hepatitis C infection in a noncirrhotic HIV infected patient on chronic dialysis with the combination of sofosbuvir and velpatasvir: A case report. *Medicine (Baltimore)* 2018;97(51):e13671.
- 528.Bae, J., D. H. Sinn, et al. Validation of the Baveno VI and the expanded Baveno VI criteria to identify patients who could avoid screening endoscopy. *Liver International* 2018;38(8):1442-1448.
- 529.Bastard, C., V. Miette, et al. A Novel FibroScan Examination Dedicated to Spleen Stiffness Measurement. *Ultrasound in Medicine and Biology* 2018;44(8):1616-1626.
- 530.Barr, R. G. Shear wave liver elastography. *Abdominal Radiology* 2018;43(4):800-807.
- 531.Benmassaoud, A., P. Ghali, et al. Screening for nonalcoholic steatohepatitis by using cytokeratin 18 and transient elastography in HIV mono-infection. *PLoS ONE* 2018;13(1):e0191985.
- 532.Akuta, N., J. Toyota, et al. Potent viral suppression and improvements in alpha-fetoprotein and measures of fibrosis in Japanese patients receiving a daclatasvir/asunaprevir/beclabuvir fixed-dose combination for the treatment of HCV genotype-1 infection. *Journal of Gastroenterology & Hepatology* 2018;53(9):1089-1097.
- 533.Arribas Anta, J., M. Garcia Gonzalez, et al. Prediction of the presence of esophageal varices using spleen stiffness measurement by transient elastography in cirrhotic patients. *Acta Gastroenterol Belg* 2018;81(4):496-501.
- 534.Ammon, F. J., A. Kohlhaas, et al. Liver stiffness reversibly increases during pregnancy and independently predicts preeclampsia. *World Journal of Gastroenterology* 2018;24(38):4393-4402.
- 535.Abdelaziz, A. O., H. I. Shousha, et al. Evaluation of liver steatosis, measured by controlled attenuation parameter, in patients with hepatitis C-induced advanced liver fibrosis and hepatocellular carcinoma. *European Journal of Gastroenterology & Hepatology* 2018;30(11):1384-1388.
- 536.Adinolfi, L. E., R. Nevola, et al. Hepatitis C virus clearance by direct-acting antiviral treatments and impact on insulin resistance in chronic hepatitis C patients. *Journal of gastroenterology and hepatology* 2018;33(7):1379-1382.
- 537.Ajmera, V. and R. Loomba. Can Elastography Differentiate Isolated Fatty Liver from Nonalcoholic Steatohepatitis? *Seminars in Liver Disease* 2018;38(1):14-20.
- 538.Alsebaey, A., M. Elhelbawy, et al. Platelets-to-lymphocyte ratio is a good predictor of liver fibrosis and insulin resistance in hepatitis C virus-related liver disease. *European Journal of Gastroenterology & Hepatology* 2018;30(2):207-211.
- 539.Ahn, S. B., D. W. Jun, et al. Optimal cutoff value for assessing changes in intrahepatic fat amount by using the controlled attenuation parameter in a longitudinal setting. *Medicine (Baltimore)* 2018;97(50):e13636.
- 540.Abdelhaleem, H., H. Gamal Eldeen, et al. Evaluation of acoustic radiation force impulse (ARFI) elastography as non-invasive diagnostic tool in living donor liver transplantation. *Abdominal Radiology* 2018;44(2):464-472.
- 541.Abozeid, M., A. Alsebaey, et al. High Efficacy of Generic and Brand Direct Acting Antivirals in Treatment of Chronic Hepatitis C. *International Journal of Infectious Diseases* 2018;75:109-144.
- 542.Anadol, E., K. Lust, et al. Exposure to previous cART is associated with significant liver fibrosis and cirrhosis in human immunodeficiency virus-infected patients. *PLoS ONE* 2018;13(1):e0191118.

2017

- 543.Yen, Y. H., K. C. Chang, et al. Elevated body mass index is a risk factor associated with possible liver cirrhosis across different etiologies of chronic liver disease. *Journal of the Formosan Medical Association* 2017.
- 544.Wang, Z., H. Yang, et al. Application of Ultrasound Elastography for Chronic Allograft Dysfunction in

- Kidney Transplantation. Journal of ultrasound in medicine 2017;36(9):1759-1769.
- 545.Wong, V. W., S. Petta, et al. Validity criteria for the diagnosis of fatty liver by M probe-based controlled attenuation parameter. Journal of Hepatology 2017;67(3):577-584.
- 546.Zizza, A., M. Guido, et al. Atherosclerosis is associated with a higher risk of hepatic steatosis in HIV-infected patients. J Prev Med Hyg 2017;58(3):E219-E224.
- 547.Staufer, K., M. Dengler, et al. The non-invasive serum biomarker soluble Axl accurately detects advanced liver fibrosis and cirrhosis. Cell Death Dis 2017;8(10):e3135.
- 548.Yeung, M. W., G. L. Wong, et al. Advanced liver fibrosis but not steatosis is independently associated with albuminuria in Chinese patients with type 2 diabetes. Journal of Hepatology 2017;S0168-8278(17):32334-6.
- 549.Trivedi, H. D., S. C. Lin, et al. Noninvasive Assessment of Fibrosis Regression in Hepatitis C Virus Sustained Virologic Responders. Gastroenterology & Hepatology 2017;13(10):587-595.
- 550.Tag-Adeen, M., M. Alsenbesy, et al. Liver stiffness measurement and spleen diameter as predictors for the presence of esophageal varices in chronic hepatitis C patients. Medicine (Baltimore) 2017;96(46):e8621.
- 551.Taneja, S., A. Borkakoty, et al. Assessment of Liver Fibrosis by Transient Elastography Should Be Done After Hemodialysis in End Stage Renal Disease Patients with Liver Disease. Digestive Diseases & Sciences 2017;62(11):3186-3192.
- 552.van de Peppel, I. P., A. Bertolini, et al. Diagnosis, follow-up and treatment of cystic fibrosis-related liver disease. Curr Opin Pulm Med 2017;23(6):562-569.
- 553.Voican, C. S., A. Louvet, et al. Transient elastography alone and in combination with FibroTest((R)) for the diagnosis of hepatic fibrosis in alcoholic liver disease. Liver International 2017;37(11):1697-1705.
- 554.Yen, Y. H., J. F. Chen, et al. The correlation of controlled attenuation parameter results with ultrasound-identified steatosis in real-world clinical practice. Journal of the Formosan Medical Association 2017;116(11):852-861.
- 555.Vinikoor, M. J., E. Sinkala, et al. Impact of Antiretroviral Therapy on Liver Fibrosis Among Human Immunodeficiency Virus-Infected Adults With and Without HBV Coinfection in Zambia. Clinical Infectious Diseases 2017;64(10):1343-1349.
- 556.Walker, A. P. Ischaemic stroke and liver fibrosis. Atherosclerosis 2017;260:153-155.
- 557.Torres-Valadez, R., S. Roman, et al. Early Detection of Liver Damage in Mexican Patients with Chronic Liver Disease. J Transl Int Med 2017;5(1):49-57.
- 558.Xu, Q., L. Sheng, et al. Evaluation of transient elastography in assessing liver fibrosis in patients with autoimmune hepatitis. Journal of gastroenterology and hepatology 2017;32(3):639-644.
- 559.Yoon, C. Y., M. Lee, et al. Fatty liver associated with metabolic derangement in patients with chronic kidney disease: A controlled attenuation parameter study. Kidney Res Clin Pract 2017;36(1):48-57.
- 560.Zachou, K., S. Gabeta, et al. COMP serum levels: A new non-invasive biomarker of liver fibrosis in patients with chronic viral hepatitis. European Journal of Internal Medicine 2017;38:83-88.
- 561.Zeremski, M. and A. Martinez. Liver Disease and Fibrosis Assessment in Substance Use-Related Disorders. Clin Pharmacol Drug Dev 2017;6(2):164-168.
- 562.Sporea, I., R. Lupusoru, et al. Dynamics of liver stiffness values by means of transient elastography in patients with HCV liver cirrhosis undergoing interferon free treatment. Journal of Gastrointestinal & Liver Diseases 2017;26(2):145-150.
- 563.Sugihara, T., M. Koda, et al. Risk Assessment of Hepatocellular Carcinoma in General Population by Liver Stiffness in Combination with Controlled Attenuation Parameter using Transient Elastography: A Cross Sectional Study. Yonago Acta Med 2017;60(2):106-112.
- 564.Toguchi, M., M. Tsurusaki, et al. Magnetic resonance elastography in the assessment of hepatic fibrosis: a study comparing transient elastography and histological data in the same patients. Abdominal Radiology 2017;42(6):1659-1666.
- 565.van Katwyk, S., D. Coyle, et al. Transient elastography for the diagnosis of liver fibrosis: a systematic review of economic evaluations. Liver International 2017;37(6):851-861.
- 566.Veiga, Z. S. T., C. A. Villela-Nogueira, et al. Transient elastography evaluation of hepatic and spleen stiffness in patients with hepatosplenic schistosomiasis. European Journal of Gastroenterology & Hepatology 2017;29(6):730-735.
- 567.Yan, L. B., X. Zhu, et al. Mutations in basal core promoter is associated with significant fibrosis in both HBeAg positive and negative treatment-naïve chronic hepatitis B. Clinics and Research in Hepatology and Gastroenterology 2017;41(3):296-302.
- 568.Tanajewski, L., R. Harris, et al. Economic evaluation of a community-based diagnostic pathway to stratify adults for non-alcoholic fatty liver disease: a Markov model informed by a feasibility study. BMJ Open 2017;7(6):e015659.
- 569.Tenggara, I. R., C. R. Lesmana, et al. Treatment Response Monitoring of Chronic Hepatitis B

- Patients using Transient Elastography and Aspartate Aminotransferase-to-Platelet Ratio Index (APRI). *Acta Medica Indonesiana* 2017;49(3):220-226.
- 570.Verdelho Machado, M. Controlled Attenuation Parameter as a Noninvasive Method to Detect and Quantify Hepatic Steatosis in Chronic Liver Disease: What Is the Clinical Relevance? *GE Port J Gastroenterol* 2017;24(4):157-160.
- 571.Welzel, T. M., T. Asselah, et al. Ombitasvir, paritaprevir, and ritonavir plus dasabuvir for 8 weeks in previously untreated patients with hepatitis C virus genotype 1b infection without cirrhosis (GARNET): a single-arm, open-label, phase 3b trial. *The lancet. Gastroenterology & hepatology* 2017;2(7):494-500.
- 573.Yip, T. C. and V. W. Wong. How to identify patients with advanced liver disease in the community? *Hepatology* 2017;66(1):7-9.
- 574.Zelber-Sagi, S., D. Shoham, et al. Predictors for advanced fibrosis in morbidly obese non-alcoholic fatty liver patients. *World Journal of Hepatology* 2017;9(2):91-98.
- 575.Stasi, C. and S. Milani. Evolving strategies for liver fibrosis staging: Non-invasive assessment. *World Journal of Gastroenterology* 2017;23(2):191-196.
- 576.Yada, N., T. Sakurai, et al. Influence of Liver Inflammation on Liver Stiffness Measurement in Patients with Autoimmune Hepatitis Evaluation by Combinational Elastography. *Oncology* 2017;92 Suppl 1:10-15.
- 577.You, M. W., K. W. Kim, et al. A Meta-analysis for the Diagnostic Performance of Transient Elastography for Clinically Significant Portal Hypertension. *Ultrasound in Medicine and Biology* 2017;43(1):59-68.
- 578.Zou, X., M. Y. Zhu, et al. Serum WFA+ -M2BP levels for evaluation of early stages of liver fibrosis in patients with chronic hepatitis B virus infection. *Liver International* 2017;37(1):35-44.
- 579.Sun, W. L., C. G. Sun, et al. [A clinical study of the association between hepatic controlled attenuation parameter and metabolic syndrome]. *Zhonghua Gan Zang Bing Za Zhi* 2017;25(2):128-133.
- 580.Zeng, J., S. Cai, et al. Dynamic Changes in Liver Stiffness Measured by Transient Elastography Predict Clinical Outcomes Among Patients With Chronic Hepatitis B. *Journal of ultrasound in medicine* 2017;36(2):261-268.
- 581.Younossi, Z. M., R. Loomba, et al. Diagnostic Modalities for Non-alcoholic Fatty Liver Disease (NAFLD), Non-alcoholic Steatohepatitis (NASH) and Associated Fibrosis. *Hepatology* 2017.
- 582.Song, J., J. Huang, et al. Performance of spleen stiffness measurement in prediction of clinical significant portal hypertension: A meta-analysis. *Clinics and Research in Hepatology and Gastroenterology* 2017;42(3):216-226.
- 583.Wang, D., P. Zhang, et al. Predictors for advanced liver fibrosis in chronic hepatitis B virus infection with persistently normal or mildly elevated alanine aminotransferase. *Experimental and therapeutic medicine* 2017;14(6):5363-5370.
- 584.Talme, T., P. Nikamo, et al. Transient Elastography May Improve Detection of Liver Fibrosis in Psoriasis Patients Treated with Methotrexate. *Acta dermatovenerologica* 2017;97(8):952-954.
- 585.Sousa, M., S. Fernandes, et al. The Baveno VI criteria for predicting esophageal varices: validation in real life practice. *Revista Española de Enfermedades Digestivas* 2017;109.
- 586.Wong, V. W., H. L. Chan, et al. Liver fibrosis progression in patients with chronic hepatitis B: a prospective study with paired transient elastography. *Hong Kong medical journal* 2017;23 Suppl 5(4):23-26.
- 587.Wu, D., E. Chen, et al. Predicting the risk of postoperative liver failure and overall survival using liver and spleen stiffness measurements in patients with hepatocellular carcinoma. *MEDICINE* 2017;96(34):e7864.
- 588.Xu, L., W. Lu, et al. A comparison of hepatic steatosis index, controlled attenuation parameter and ultrasound as noninvasive diagnostic tools for steatosis in chronic hepatitis B. *Digestive & Liver Disease* 2017;49(8):910-917.
- 589.Zeng, J., J. Zheng, et al. Comparison of 2-D Shear Wave Elastography and Transient Elastography for Assessing Liver Fibrosis in Chronic Hepatitis B. *Ultrasound in Medicine and Biology* 2017;43(8):1563-1570.
- 590.Summers, J. A., M. Radhakrishnan, et al. Virtual Touch Quantification to Diagnose and Monitor Liver Fibrosis in Hepatitis B and Hepatitis C: A NICE Medical Technology Guidance. *Applied Health Economics Health Policy* 2017;15(2):139-154.
- 591.Yu, J. H. and J. I. Lee. Current role of transient elastography in the management of chronic hepatitis B patients. *Ultrasonography* 2017;36(2):86-94.
- 592.Zeng, J., C. Sun, et al. Association between non-invasively diagnosed hepatic steatosis and chronic kidney disease in Chinese adults on their health check-up. *Journal of Digestive Diseases* 2017;18(4):229-236.
- 593.Sulyok, M., T. Ferenci, et al. Hepatic fibrosis and factors associated with liver stiffness in HIV mono-infected individuals. *PeerJ* 2017;5:e2867.
- 594.Tag-Adeen, M., A. M. Sabra, et al. Impact of hepatitis C virus genotype-4 eradication following direct acting antivirals on liver stiffness measurement. *Hepatic Medicine: Evidence and Research* 2017;9:45-53.
- 595.Zhao, J., F. Zhai, et al. Evaluating the Significance of Viscoelasticity in Diagnosing Early-Stage Liver

- Fibrosis with Transient Elastography. *PLoS ONE* 2017;12(1):e0170073.
596. Mauro, E., G. Crespo, et al. Portal pressure and liver stiffness measurements in the prediction of fibrosis regression after SVR in recurrent hepatitis C. *Hepatology* 2017.
597. Price, J. C., J. L. Dodge, et al. Controlled attenuation parameter and magnetic resonance spectroscopy-measured liver steatosis are discordant in obese HIV-infected adults. *Aids* 2017;31(15):2119-2125.
598. Mahady, S. E., P. Macaskill, et al. Diagnostic Accuracy of Noninvasive Fibrosis Scores in a Population of Individuals With a Low Prevalence of Fibrosis. *Clinical Gastroenterology & Hepatology* 2017;15(9):1453-1460 e1.
599. Marshall, A. D., J. Grebely, et al. 'I didn't want to let it go too far.' The decisions and experiences of people who inject drugs who received a liver disease assessment as part of a liver health promotion campaign: The LiveRLife study. *International Journal on Drug Policy* 2017;47:153-160.
600. Mulazzani, L., V. Salvatore, et al. Point shear wave ultrasound elastography with Esaote compared to real-time 2D shear wave elastography with supersonic imagine for the quantification of liver stiffness. *Journal of ultrasound* 2017;20(3):213-225.
601. Naveau, S., C. S. Voican, et al. Controlled attenuation parameter for diagnosing steatosis in bariatric surgery candidates with suspected nonalcoholic fatty liver disease. *European Journal of Gastroenterology & Hepatology* 2017;29(9):1022-1030.
602. Nunez-Torres, R., J. Macias, et al. Fat mass and obesity-associated gene variations are related to fatty liver disease in HIV-infected patients. *HIV Medicine* 2017;18(8):546-554.
603. Omran, D. A., S. K. Darweesh, et al. Serum Zinc Deficiency and its Relation to Liver Fibrosis in Chronic HCV: a Real-Life Egyptian Study. *Biol Trace Elem Res* 2017;179(1):1-7.
604. Potts, J. R., C. M. Maybury, et al. Diagnosing liver fibrosis: a narrative review of current literature for dermatologists. *British Journal of Dermatology* 2017;177(3):637-644.
605. Seko, Y., Y. Sumida, et al. Effect of sodium glucose cotransporter 2 inhibitor on liver function tests in Japanese patients with non-alcoholic fatty liver disease and type 2 diabetes mellitus. *Hepatology Research* 2017;47(10):1072-1078.
606. Sharma, P., R. Agarwal, et al. Transient Elastography (Fibroscan) in Patients with Non-cirrhotic Portal Fibrosis. *Journal of Clinical and Experimental Hepatology* 2017;7(3):230-234.
607. Pembroke, T., M. Deschenes, et al. Hepatic steatosis progresses faster in HIV mono-infected than HIV/HCV co-infected patients and is associated with liver fibrosis. *Journal of Hepatology* 2017;67(4):801-808.
608. Ragazzo, T. G., D. Paranagua-Vezozzo, et al. Accuracy of transient elastography-FibroScan(R), acoustic radiation force impulse (ARFI) imaging, the enhanced liver fibrosis (ELF) test, APRI, and the FIB-4 index compared with liver biopsy in patients with chronic hepatitis C. *Clinics (Sao Paulo)* 2017;72(9):516-525.
609. Rajakannu, M., D. Cherqui, et al. Liver stiffness measurement by transient elastography predicts late posthepatectomy outcomes in patients undergoing resection for hepatocellular carcinoma. *Surgery* 2017;162(4):766-774.
610. Seko, Y., Y. Sumida, et al. Effect of 12-week dulaglutide therapy in Japanese patients with biopsy-proven non-alcoholic fatty liver disease and type 2 diabetes mellitus. *Hepatology Research* 2017;47(11):1206-1211.
611. Rouhi, A., G. Hazlewood, et al. Prevalence and risk factors for liver fibrosis detected by transient elastography or shear wave elastography in inflammatory arthritis: a systematic review. *Clinical and Experimental Rheumatology* 2017;35(6):1029-1036.
612. Pozzi, R., I. Parzanese, et al. Point shear-wave elastography in chronic pancreatitis: A promising tool for staging disease severity. *Pancreatology* 2017;17(6):905-910.
613. Qu, H. J., L. Wang, et al. [Value of hydrogen proton magnetic resonance spectroscopy in the determination of liver triglyceride in patients with fatty liver disease and its influencing factors]. *Zhonghua Gan Zang Bing Za Zhi* 2017;25(11):858-863.
614. Llop, E., M. Lopez, et al. Validation of noninvasive methods to predict the presence of gastroesophageal varices in a cohort of patients with compensated advanced chronic liver disease. *Journal of gastroenterology and hepatology* 2017;32(11):1867-1872.
615. Maira, D., E. Cassinero, et al. Progression of liver fibrosis can be controlled by adequate chelation in transfusion-dependent thalassemia (TDT). *Ann Hematol* 2017;96(11):1931-1936.
616. Mattos, A. Z. and A. A. Mattos. Transient Elastography vs. Aspartate Aminotransferase to Platelet Ratio Index in Hepatitis C: A Meta-Analysis. *Annals of Hepatology* 2017;16(3):349-357.
617. Ryan, J. D. and E. A. Tsouchatzis. MRE in NAFLD: Promising but Further Validation is Required. *Annals of Hepatology* 2017;16(3):331-332.
618. Mauss, S., S. Pol, et al. Late presentation of chronic viral hepatitis for medical care: a consensus definition. *BMC Medicine* 2017;15(1):92.
619. Mukund, A., S. S. Pargewar, et al. Changes in Liver Congestion in Patients with Budd-Chiari Syndrome following Endovascular Interventions: Assessment

- with Transient Elastography. *Journal of Vascular and Interventional Radiology* 2017;28(5):683-687.
- 620.Romero-Gomez, M. and H. Cortez-Pinto. Detecting liver fat from viscoelasticity: How good is CAP in clinical practice? The need for universal cut-offs. *Journal of Hepatology* 2017;66(5):886-887.
- 621.Paranagua-Vezozzo, D. C., A. Andrade, et al. Concordance of non-invasive mechanical and serum tests for liver fibrosis evaluation in chronic hepatitis C. *World Journal of Hepatology* 2017;9(8):436-442.
- 622.Silva, M. J., C. Bernardes, et al. Baveno VI Recommendation on Avoidance of Screening Endoscopy in Cirrhotic Patients: Are We There Yet? *GE Port J Gastroenterol* 2017;24(2):79-83.
- 623.Olivares-Gazca, J. C., A. K. Nunez-Cortes, et al. More on the thrombocytopenia of the non-alcoholic fatty liver disease. *Hematology* 2017;22(5):316-319.
- 624.Patel, P. J., K. L. Hayward, et al. Multimorbidity and polypharmacy in diabetic patients with NAFLD: Implications for disease severity and management. *Medicine (Baltimore)* 2017;96(26):e6761.
- 625.Loong, T. C., J. L. Wei, et al. Application of the combined FibroMeter vibration-controlled transient elastography algorithm in Chinese patients with non-alcoholic fatty liver disease. *Journal of gastroenterology and hepatology* 2017;32(7):1363-1369.
- 626.Okajima, A., Y. Sumida, et al. Liver stiffness measurement to platelet ratio index predicts the stage of liver fibrosis in non-alcoholic fatty liver disease. *Hepatology Research* 2017;47(8):721-730.
- 627.Pang, J., W. Xu, et al. Significant positive association of endotoxemia with histological severity in 237 patients with non-alcoholic fatty liver disease. *Alimentary Pharmacology & Therapeutics* 2017;46(2):175-182.
- 628.Pavlides, M., R. Banerjee, et al. Multiparametric magnetic resonance imaging for the assessment of non-alcoholic fatty liver disease severity. *Liver International* 2017;37(7):1065-1073.
- 629.Piscaglia, F., V. Salvatore, et al. Differences in liver stiffness values obtained with new ultrasound elastography machines and Fibroscan: A comparative study. *Digestive & Liver Disease* 2017;49(7):802-808.
- 630.Rongngern, P., L. Chularojanamontri, et al. Diagnostic performance of transient elastography for detection of methotrexate-induced liver injury using Roenigk classification in Asian patients with psoriasis: a retrospective study. *Arch Dermatol Res* 2017;309(5):403-408.
- 631.Sinakos, E., D. Kountouras, et al. Treatment of chronic hepatitis C with direct-acting antivirals in patients with beta-thalassaemia major and advanced liver disease. *British Journal of Haematology* 2017;178(1):130-136.
- 632.Pu, K., J. H. Shi, et al. Diagnostic accuracy of transient elastography (FibroScan) in detection of esophageal varices in patients with cirrhosis: A meta-analysis. *World Journal of Gastroenterology* 2017;23(2):345-356.
- 633.Ou, G., H. H. Ko, et al. Utility of Transient Elastography in Estimating Hepatic Iron Concentration in Comparison to Magnetic Resonance Imaging in Patients Who are Transfusion-Dependent: A Canadian Center Experience. *Hemoglobin* 2017;41(1):21-25.
- 634.Paternostro, R., T. Reiberger, et al. Plasma renin concentration represents an independent risk factor for mortality and is associated with liver dysfunction in patients with cirrhosis. *Journal of Gastroenterology & Hepatology* 2017;32(1):184-190.
- 635.Schwabl, P., M. Mandorfer, et al. Interferon-free regimens improve portal hypertension and histological necroinflammation in HIV/HCV patients with advanced liver disease. *Alimentary Pharmacology & Therapeutics* 2017;45(1):139-149.
- 636.Sherman, D., P. Lung, et al. Virtual touch quantification (VTq) elastography for non-invasive assessment of liver disease and its complications: what the clinician needs to know. *Frontline Gastroenterol* 2017;8(1):37-44.
- 637.Puigvehi, M., T. Broquetas, et al. Impact of Anthropometric Features on the Applicability and Accuracy of FibroScan(R) (M and XL) in Overweight/Obese Patients. *Journal of gastroenterology and hepatology* 2017;32(10):1746-1753.
- 638.Mare, R., I. Sporea, et al. The value of ElastPQ for the evaluation of liver stiffness in patients with B and C chronic hepatopathies. *Ultrasonics* 2017;77:144-151.
- 639.Ou, H., S. Cai, et al. A noninvasive diagnostic model to assess nonalcoholic hepatic steatosis in patients with chronic hepatitis B. *Therapeutic Advances in Gastroenterology* 2017;10(2):207-217.
- 640.Parikh, P., J. D. Ryan, et al. Fibrosis assessment in patients with chronic hepatitis B virus (HBV) infection. *Ann Transl Med* 2017;5(3):40.
- 641.Park, C. C., P. Nguyen, et al. Magnetic Resonance Elastography vs Transient Elastography in Detection of Fibrosis and Noninvasive Measurement of Steatosis in Patients With Biopsy-Proven Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2017;152(3):598-607 e2.
- 642.Mohamed, S. Y., E. F. Mostafa, et al. The relationship between expression of Toll-like receptor 4 in chronic hepatitis C patients and different stages of liver fibrosis. *Gastroenterol Hepatol Bed Bench* 2017;10(4):278-283.
- 643.Seo, Y. S., B. K. Jang, et al. Validation of risk prediction models for the development of HBV-related HCC: a retrospective multi-center 10-year

- follow-up cohort study. *Oncotarget* 2017;8(68):113213-113224.
644. Mebus, S., N. Nagdyman, et al. Non-invasive assessment of liver changes in Eisenmenger patients. *Int J Cardiol* 2017;249:140-144.
645. Liu, K., V. W. Wong, et al. Prognostic Value of Controlled Attenuation Parameter by Transient Elastography. *American Journal of Gastroenterology* 2017;112(12):1812-1823.
646. Marino, Z., J. M. Pascasio-Acevedo, et al. High efficacy of Sofosbuvir plus Simeprevir in a large cohort of Spanish cirrhotic patients infected with genotypes 1 and 4. *Liver International* 2017;37(12):1823-1832.
647. Ong, S. Y., L. C. Gurrin, et al. Reduction of body iron in HFE-related haemochromatosis and moderate iron overload (Mi-Iron): a multicentre, participant-blinded, randomised controlled trial. *Lancet Haematol* 2017;4(12):e607-e614.
648. Paul, S. B., P. Das, et al. Assessment of liver fibrosis in chronic hepatitis: comparison of shear wave elastography and transient elastography. *Abdominal Radiology* 2017;42(12):2864-2873.
649. Qi, M., Y. Chen, et al. Clinical significance of preoperative liver stiffness measurements in primary HBV-positive hepatocellular carcinoma. *Future oncology* 2017;13(30):2799-2810.
650. Ramzy, I., A. Elsharkawy, et al. Impact of old Schistosomiasis infection on the use of transient elastography (Fibroscan) for staging of fibrosis in chronic HCV patients. *Acta Trop* 2017;176:283-287.
651. Roulot, D., F. Roudot-Thoraval, et al. Concomitant screening for liver fibrosis and steatosis in French type 2 diabetic patients using Fibroscan. *Liver International* 2017;37(12):1897-1906.
652. Nosotti, L., A. Petrelli, et al. Distribution of IL28B Polymorphism in a Cohort of Italians and Immigrants with HCV Infection: Association with Viraemia, Stage of Fibrosis and Response to Treatment. *Journal of Immigrant and Minority Health* 2017;19(4):876-882.
653. Peiseler, M., A. Creutzfeldt, et al. Potential impact of screening for fatty liver disease by transient elastography with liver stiffness and controlled attenuation parameter measurements: a pilot study. *Zeitschrift für Gastroenterologie* 2017;55(8):754-760.
654. Pons, M., B. Santos, et al. Rapid liver and spleen stiffness improvement in compensated advanced chronic liver disease patients treated with oral antivirals. *Therapeutic Advances in Gastroenterology* 2017;10(8):619-629.
655. Puustinen, L., A. Hakkarainen, et al. 31Phosphorus magnetic resonance spectroscopy of the liver for evaluating inflammation and fibrosis in autoimmune hepatitis. *Scandinavian Journal of Gastroenterology* 2017;52(8):886-892.
656. Seki, K., T. Shima, et al. Assessment of transient elastography in Japanese patients with non-alcoholic fatty liver disease. *Hepatology Research* 2017;47(9):882-889.
657. McLeay, S. C., W. M. Harvey, et al. Physical comorbidities of post-traumatic stress disorder in Australian Vietnam War veterans. *The Medical Journal of Australia* 2017;206(6):251-257.
658. Petta, S., V. W. Wong, et al. Improved noninvasive prediction of liver fibrosis by liver stiffness measurement in patients with nonalcoholic fatty liver disease accounting for controlled attenuation parameter values. *Hepatology* 2017;65(4):1145-1155.
659. Poinard, T., V. Peta, et al. Screening studies of transient elastography and FibroTest in the general population. *Lancet Gastroenterol Hepatol* 2017;2(4):246.
660. Prasetya, I. B., I. Hasan, et al. Prevalence and Profile of Fibrosis in Diabetic Patients with Non-alcoholic Fatty Liver Disease and the Associated Factors. *Acta Medica Indonesiana* 2017;49(2):91-98.
661. Raizner, A., N. Shillingford, et al. Hepatic Inflammation May Influence Liver Stiffness Measurements by Transient Elastography in Children and Young Adults. *Journal of Pediatric Gastroenterology and Nutrition* 2017;64(4):512-517.
662. Marconi, C., M. Bellan, et al. Cardiac dysfunction as an early predictor of portal hypertension in chronic hepatitis C. *Annals of gastroenterology* 2017;30(6):675-681.
663. Marozas, M., R. Zykus, et al. Noninvasive Evaluation of Portal Hypertension Using a Supervised Learning Technique. *J Healthc Eng* 2017;2017:6183714.
664. Parisi, S. G., M. Basso, et al. Liver stiffness is not associated with short- and long-term plasma HIV RNA replication in immunocompetent patients with HIV infection and with HIV/HCV coinfection. *Annals of gastroenterology* 2017;30(5):534-541.
665. Patel, P. J., D. Smith, et al. Alcohol Consumption in Diabetic Patients with Nonalcoholic Fatty Liver Disease. *Canadian Journal of Gastroenterology & Hepatology* 2017;2017:7927685.
666. Laufer, N., D. Ojeda, et al. CD4(+) T cells and natural killer cells: Biomarkers for hepatic fibrosis in human immunodeficiency virus/hepatitis C virus-coinfected patients. *World Journal of Hepatology* 2017;9(25):1073-1080.
667. Huang, R., X. Yan, et al. Can Transient Elastography Predict Fibrosis Regression in Patients with Chronic Hepatitis B During Long-Term Antiviral Therapy? *American Journal of Gastroenterology* 2017;112(9):1477-1478.
668. Hwang, J. A., T. W. Kang, et al. Association between non-hypervascular hypointense nodules on gadoxetic acid-enhanced MRI and liver stiffness or hepatocellular carcinoma. *European Journal of Radiology* 2017;95:362-369.

- 669.Joshi, S. S., D. Wong, et al. Peripartum cytokine flares in a multiethnic cohort of chronic hepatitis B carriers does not correlate with hepatitis B virus suppression or increased risk of liver disease. *American journal of reproductive immunology* 2017;78(4).
- 670.Lee, Y. H., K. J. Kim, et al. Association of non-alcoholic steatohepatitis with subclinical myocardial dysfunction in non-cirrhotic patients. *Journal of Hepatology* 2017;68(4):764-772.
- 671.Koizumi, Y., M. Hirooka, et al. Comparison between real-time tissue elastography and vibration-controlled transient elastography for the assessment of liver fibrosis and disease progression in patients with primary biliary cholangitis. *Hepatology Research* 2017;47(12):1252-1259.
- 672.Lee, J. E., K. S. Shin, et al. Non-invasive Assessment of Liver Fibrosis with ElastPQ: Comparison with Transient Elastography and Serologic Fibrosis Marker Tests, and Correlation with Liver Pathology Results. *Ultrasound in Medicine and Biology* 2017;43(11):2515-2521.
- 673.Karlas, T., L. Weise, et al. Correlation of cell-free DNA plasma concentration with severity of non-alcoholic fatty liver disease. *Journal of translational medicine* 2017;15(1):106.
- 674.Jeon, M. Y., H. W. Lee, et al. Subcirrhotic liver stiffness by FibroScan correlates with lower risk of hepatocellular carcinoma in patients with HBV-related cirrhosis. *Hepatology International* 2017;11(3):268-276.
- 675.Karlas, T., D. Petroff, et al. Individual patient data meta-analysis of controlled attenuation parameter (CAP) technology for assessing steatosis. *Journal of Hepatology* 2017;66(5):1022-1030.
- 676.Kim, S. U., D. Song, et al. Liver fibrosis assessed with transient elastography is an independent risk factor for ischemic stroke. *Atherosclerosis* 2017;260:156-162.
- 677.Lackner, C., R. Bataller, et al. Fibrosis evaluation by transient elastography in alcoholic liver disease: Is the histological scoring system impacting cutoff values? *Hepatology* 2017;65(5):1758-1761.
- 678.Lee, Y. R., S. Y. Park, et al. Using transient elastography to predict hepatocellular carcinoma recurrence after radiofrequency ablation. *Journal of gastroenterology and hepatology* 2017;32(5):1079-1086.
- 679.Kim, G., M. Y. Kim, et al. Transient elastography versus hepatic venous pressure gradient for diagnosing portal hypertension: a systematic review and meta-analysis. *Clinical and Molecular Hepatology* 2017;23(1):34-41.
- 680.Kim, B. K. Transient elastography can be integrated into routine clinical practice for the evaluation of portal hypertension? *Clinical and Molecular Hepatology* 2017;23(1):27-30.
- 681.Li, X., Q. Jin, et al. Changes in liver stiffness and its associated factors during oral antiviral therapy in Chinese patients with chronic hepatitis B. *Experimental and therapeutic medicine* 2017;13(3):1169-1175.
- 682.Karlas, T., S. Beer, et al. Do we need controlled attenuation parameter adjustment for fibrosis estimation in nonalcoholic fatty liver disease patients? *Hepatology* 2017;65(6):2126-2128.
- 683.Li, X., Q. Jin, et al. Chronic hepatitis B patients with high liver fibrosis levels should receive antiviral treatment. *Experimental and therapeutic medicine* 2017;13(6):3624-3630.
- 684.Latt, N. L., B. T. Yanny, et al. Eight-week ledipasvir/sofosbuvir in non-cirrhotic, treatment-naive hepatitis C genotype-1 patients with hepatitis C virus-RNA < 6 million: Single center, real world effectiveness and safety. *World Journal of Gastroenterology* 2017;23(26):4759-4766.
- 685.Lee, E. S., J. B. Lee, et al. Shear Wave Liver Elastography with a Propagation Map: Diagnostic Performance and Inter-Observer Correlation for Hepatic Fibrosis in Chronic Hepatitis. *Ultrasound in Medicine and Biology* 2017;43(7):1355-1363.
- 686.Kumar, A., N. M. Khan, et al. Correlation of transient elastography with hepatic venous pressure gradient in patients with cirrhotic portal hypertension: A study of 326 patients from India. *World Journal of Gastroenterology* 2017;23(4):687-696.
- 687.Huh, J. H., K. J. Kim, et al. Obesity is more closely related with hepatic steatosis and fibrosis measured by transient elastography than metabolic health status. *Metabolism: Clinical and Experimental* 2017;66:23-31.
- 688.Lee, J. M., Y. S. Seo, et al. The LAW index as an accurate indicator of the initiation of antiviral treatment in patients with chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2017;32(1):208-214.
- 689.Lee, Y. H., J. H. Kim, et al. Lobeglitazone, a Novel Thiazolidinedione, Improves Non-Alcoholic Fatty Liver Disease in Type 2 Diabetes: Its Efficacy and Predictive Factors Related to Responsiveness. *Journal of Korean medical science* 2017;32(1):60-69.
- 690.Lemoine, M., M. Thursz, et al. Diagnostic accuracy of the gamma-glutamyl transpeptidase to platelet ratio (GPR) using transient elastography as a reference. *Gut* 2017;66(1):195-196.
- 691.Leung, J. C., T. C. Loong, et al. Histological severity and clinical outcomes of nonalcoholic fatty liver disease in nonobese patients. *Hepatology* 2017;65(1):54-64.
- 692.Jaquet, A., G. Wandeler, et al. Alcohol use, viral hepatitis and liver fibrosis among HIV-positive persons in West Africa: a cross-sectional study. *Journal of the International AIDS Society* 2017;19(1):1-10.

- 693.Karlas, T., D. Petroff, et al. Collaboration, Not Competition: The Role of Magnetic Resonance, Transient Elastography, and Liver Biopsy in the Diagnosis of Nonalcoholic Fatty Liver Disease. *Gastroenterology* 2017;152(3):479-481.
- 694.Liang, X. E., C. Zhong, et al. Optimization of hepatitis B cirrhosis detection by stepwise application of transient elastography and routine biomarkers. *Journal of Gastroenterology & Hepatology* 2017;32(2):459-465.
- 695.Lei, J. W., X. Y. Ji, et al. Prediction of posthepatectomy liver failure using transient elastography in patients with hepatitis B related hepatocellular carcinoma. *BMC Gastroenterology* 2017;17(1):171.
- 696.Liang, X. E. and Y. P. Chen. Clinical Application of Vibration Controlled Transient Elastography in Patients with Chronic Hepatitis B. *Journal of Clinical and Translational Hepatology* 2017;5(4):368-375.
- 697.Leblebicioğlu, H., J. E. Arends, et al. Availability of hepatitis C diagnostics and therapeutics in European and Eurasia countries. *Antiviral Research* 2017;150:9-14.
- 698.Katwesigye, E., E. Seremba, et al. Low sero-prevalence of hepatitis delta antibodies in HIV/hepatitis B co-infected patients attending an urban HIV clinic in Uganda. *Afr Health Sci* 2017;17(4):974-978.
- 699.Krawczyk, M., J. Ligocka, et al. Does transient elastography correlate with liver fibrosis in patients with PSC? Laennec score-based analysis of explanted livers. *Scandinavian Journal of Gastroenterology* 2017;52(12):1407-1412.
- 700.Horowitz, J. M., S. K. Venkatesh, et al. Evaluation of hepatic fibrosis: a review from the society of abdominal radiology disease focus panel. *Abdominal Radiology* 2017;42(8):2037-2053.
- 701.Jangouk, P., L. Turco, et al. Validating, deconstructing and refining Baveno criteria for ruling out high-risk varices in patients with compensated cirrhosis. *Liver International* 2017;37(8):1177-1183.
- 702.Jaquet, A., M. Nouaman, et al. Hepatitis B treatment eligibility in West Africa: Uncertainties and need for prospective cohort studies. *Liver International* 2017;37(8):1116-1121.
- 703.Lee, H. W., B. K. Kim, et al. Prevalence and Predictors of Significant Fibrosis Among Subjects with Transient Elastography-Defined Nonalcoholic Fatty Liver Disease. *Digestive Diseases & Sciences* 2017;62(8):2150-2158.
- 704.Kashiyama, N., K. Toda, et al. Evaluation of right ventricular function using liver stiffness in patients with left ventricular assist device. *European journal of cardio-thoracic surgery* 2017;51(4):715-721.
- 705.Kowdley, K. V., V. Sundaram, et al. Eight weeks of ledipasvir/sofosbuvir is effective for selected patients with genotype 1 hepatitis C virus infection. *Hepatology* 2017;65(4):1094-1103.
- 706.Jun, B. G., W. Y. Park, et al. A prospective comparative assessment of the accuracy of the FibroScan in evaluating liver steatosis. *PLoS ONE* 2017;12(8):e0182784.
- 707.Karajamaki, A. J., O. Kettunen, et al. Presence of atrial fibrillation is associated with liver stiffness in an elderly Finnish population. *PLoS ONE* 2017;12(3):e0173855.
- 708.Kim, G., K. J. Kim, et al. Significant liver fibrosis assessed using liver transient elastography is independently associated with low bone mineral density in patients with non-alcoholic fatty liver disease. *PLoS ONE* 2017;12(7):e0182202.
- 709.Kjaergaard, M., M. Thiele, et al. High risk of misinterpreting liver and spleen stiffness using 2D shear-wave and transient elastography after a moderate or high calorie meal. *PLoS ONE* 2017;12(4):e0173992.
- 710.Klotter, V., C. Gunchick, et al. Assessment of pathologic increase in liver stiffness enables earlier diagnosis of CFLD: Results from a prospective longitudinal cohort study. *PLoS ONE* 2017;12(6):e0178784.
- 711.Lee, H. W., K. J. Kim, et al. The relationship between visceral obesity and hepatic steatosis measured by controlled attenuation parameter. *PLoS ONE* 2017;12(10):e0187066.
- 712.Lee, M. S., J. M. Bae, et al. Prospective comparison among transient elastography, supersonic shear imaging, and ARFI imaging for predicting fibrosis in nonalcoholic fatty liver disease. *PLoS ONE* 2017;12(11):e0188321.
- 713.Liang, J., F. Liu, et al. A Noninvasive Score Model for Prediction of NASH in Patients with Chronic Hepatitis B and Nonalcoholic Fatty Liver Disease. *BioMed Research International* 2017;2017:8793278.
- 714.Elsharkawy, A., S. A. Alem, et al. Changes in liver stiffness measurements and fibrosis scores following sofosbuvir based treatment regimens without interferon. *Journal of gastroenterology and hepatology* 2017;32(9):1624-1630.
- 715.Ferro, Y., I. Care, et al. Protein and vitamin B6 intake are associated with liver steatosis assessed by transient elastography, especially in obese individuals. *Clinical and Molecular Hepatology* 2017;23(3):249-259.
- 716.Gheorghe, L., S. Jacob, et al. Real-Life Use of 3 Direct-Acting Antiviral Regimen in a Large Cohort of Patients with Genotype-1b HCV Compensated Cirrhosis. *Journal of Gastrointestinal & Liver Diseases* 2017;26(3):275-281.
- 717.Gutierrez-Grobe, Y., E. Juarez-Hernandez, et al. Less liver fibrosis in metabolically healthy compared with metabolically unhealthy obese patients with non-alcoholic fatty liver disease. *Diabetes & Metabolism* 2017;43(4):332-337.

- 718.Guo, L., L. Zheng, et al. Transient Elastography (FibroScan) Performs Better Than Non-Invasive Markers in Assessing Liver Fibrosis and Cirrhosis in Autoimmune Hepatitis Patients. *Medical science monitor : international medical journal of experimental and clinical research* 2017;23:5106-5112.
- 719.Hartl, J., H. Ehlken, et al. Usefulness of biochemical remission and transient elastography in monitoring disease course in autoimmune hepatitis. *Journal of Hepatology* 2017;68(4):754-763.
- 720.Dolmazashvili, E., A. Abutidze, et al. Regression of liver fibrosis over a 24-week period after completing direct-acting antiviral therapy in patients with chronic hepatitis C receiving care within the national hepatitis C elimination program in Georgia: results of hepatology clinic HEPA experience. *European Journal of Gastroenterology & Hepatology* 2017;29(11):1223-1230.
- 721.Ebert, T., N. Linder, et al. Association of fetuin B with markers of liver fibrosis in nonalcoholic fatty liver disease. *Endocrine* 2017;58(2):246-252.
- 722.Ferraioli, G., V. Calcaterra, et al. Noninvasive assessment of liver steatosis in children: the clinical value of controlled attenuation parameter. *BMC Gastroenterology* 2017;17(1):61.
- 723.Grgurevic, I., T. Bokun, et al. Non-invasive diagnosis of portal hypertension in cirrhosis using ultrasound based elastography. *Medical Ultrasonography Journal* 2017;19(3):310-317.
- 724.Expert Panel on Gastrointestinal, I., J. M. Horowitz, et al. ACR Appropriateness Criteria(R) Chronic Liver Disease. *Journal of the American College of Radiology* 2017;14(5S):S103-S117.
- 725 Engelmann, G., J. Quader, et al. Limitations and opportunities of non-invasive liver stiffness measurement in children. *World Journal of Hepatology* 2017;9(8):409-417.
- 726.He, X. N., X. F. Diao, et al. Improved shear wave motion detection using coded excitation for transient elastography. *Sci Rep* 2017;7:44483.
- 727.Dolmazashvili, E., M. Karchava, et al. Comparative Study of Fib-4 Index and Transient Elastography among Patients with Chronic Hepatitis C Virus Infection in Georgia. *Georgian Medical News* 2017;(264):81-86.
- 728.Friedrich-Rust, M., F. Schoelzel, et al. Safety of transient elastography in patients with implanted cardiac rhythm devices. *Digestive & Liver Disease* 2017;49(3):314-316.
- 729.Ferraioli, G., L. Maiocchi, et al. Ruling-in and Ruling-out Significant Fibrosis and Cirrhosis in Patients with Chronic Hepatitis C Using a Shear Wave Measurement Method. *Journal of Gastrointestinal & Liver Diseases* 2017;26(2):139-143.
- 730.Grant, J., O. Agbaji, et al. Hepatitis B virus sequencing and liver fibrosis evaluation in HIV/HBV co-infected Nigerians. *Tropical medicine & international health* 2017;22(6):744-754.
- 731.Hong, Y. M., K. T. Yoon, et al. Clinical usefulness of controlled attenuation parameter to screen hepatic steatosis for potential donor of living donor liver transplant. *European Journal of Gastroenterology & Hepatology* 2017;29(7):805-810.
- 732.Hallager, S., P. Brehm Christensen, et al. Mortality Rates in Patients With Chronic Hepatitis C and Cirrhosis Compared With the General Population: A Danish Cohort Study. *The Journal of Infectious Diseases* 2017;215(2):192-201.
- 733.Hodge, A., S. Lim, et al. Coffee Intake Is Associated with a Lower Liver Stiffness in Patients with Non-Alcoholic Fatty Liver Disease, Hepatitis C, and Hepatitis B. *Nutrients* 2017;9(1).
- 734.European Paediatric, H. C.-i. S. G. i. t. E. P. and H. I. V. C. C. i. E. Paediatric. Coinfection with HIV and hepatitis C virus in 229 children and young adults living in Europe. *Aids* 2017;31(1):127-135.
- 735.Elkabbany, Z. A., N. S. Elbarbary, et al. Transient elastography as a noninvasive assessment tool for hepatopathies of different etiology in pediatric type 1 diabetes mellitus. *Journal of diabetes and its complications* 2017;31(1):186-194.
- 736.Gianni, E., P. Forte, et al. Prospective Evaluation of Liver Stiffness Using Transient Elastography in Alcoholic Patients Following Abstinence. *Alcohol Alcohol* 2017;52(1):42-47.
- 737.Drasar, E., E. Fitzpatrick, et al. Interim assessment of liver damage in patients with sickle cell disease using new non-invasive techniques. *British Journal of Haematology* 2017;176(4):643-650.
- 738.Gollisch, K. S., A. Lindhorst, et al. EndoBarrier Gastrointestinal Liner in Type 2 Diabetic Patients Improves Liver Fibrosis as Assessed by Liver Elastography. *Experimental and clinical endocrinology & diabetes* 2017;125(2):116-121.
- 739.Guardiola Arevalo, A., R. Gomez Rodriguez, et al. Characteristics and course of chronic hepatitis B e antigen-negative infection. *Gastroenterología y hepatología* 2017;40(2):59-69.
- 740.Eddowes, P. J., N. McDonald, et al. Utility and cost evaluation of multiparametric magnetic resonance imaging for the assessment of non-alcoholic fatty liver disease. *Alimentary Pharmacology & Therapeutics* 2017.
- 741.Elsharkawy, A., M. Alboraie, et al. Establishing ultrasound based transient elastography cutoffs for different stages of hepatic fibrosis and cirrhosis in Egyptian chronic hepatitis C patients. *Arab journal of gastroenterology* 2017;18(4):210-215.
- 742.Grgurevic, I., T. Bozin, et al. Hepatitis C is now curable, but what happens with cirrhosis and portal hypertension afterwards? *Clin Exp Hepatol* 2017;3(4):181-186.
- 743.Honda, Y., T. Kessoku, et al. Efficacy of glutathione for the treatment of nonalcoholic fatty liver disease:

- an open-label, single-arm, multicenter, pilot study. *BMC Gastroenterology* 2017;17(1):96.
- 744.Ducancelle, A., V. Leroy, et al. A Single Test Combining Blood Markers and Elastography is More Accurate Than Other Fibrosis Tests in the Main Causes of Chronic Liver Diseases. *Journal of Clinical Gastroenterology* 2017;51(7):639-649.
- 745.Honsawek, S., W. Udomsinprasert, et al. Elevated serum heat shock protein 70 and liver stiffness reflect hepatic dysfunction and severity in postoperative biliary atresia. *Pediatric Surgery International* 2017;33(8):893-899.
- 746.Gani, R. A., I. Hasan, et al. Evaluation of Acoustic Radiation Force Impulse (ARFI) for Fibrosis Staging in Chronic Liver Diseases. *Acta Medica Indonesiana* 2017;49(2):128-135.
- 747.Gines, P., I. Graupera, et al. Screening studies of transient elastography and FibroTest in the general population - Authors' reply. *Lancet Gastroenterol Hepatol* 2017;2(4):246-247.
- 748.Han, M. A., R. Saouaf, et al. Magnetic resonance imaging and transient elastography in the management of Nonalcoholic Fatty Liver Disease (NAFLD). *Expert Rev Clin Pharmacol* 2017;10(4):379-390.
- 749.Harris, R., D. J. Harman, et al. Prevalence of clinically significant liver disease within the general population, as defined by non-invasive markers of liver fibrosis: a systematic review. *Lancet Gastroenterol Hepatol* 2017;2(4):288-297.
- 750.Endo, M., Y. Soroida, et al. Ultrasound evaluation of liver stiffness: accuracy of ultrasound imaging for the prediction of liver cirrhosis as evaluated using a liver stiffness measurement. *J Med Dent Sci* 2017;64(2-3):27-34.
- 751.Fernandes, F. F., H. Perazzo, et al. Latent Class Analysis of Noninvasive Methods and Liver Biopsy in Chronic Hepatitis C: An Approach without a Gold Standard. *BioMed Research International* 2017;2017:8252980.
- 752.Fidai, A., F. Dallaire, et al. Non-invasive Investigations for the Diagnosis of Fontan-Associated Liver Disease in Pediatric and Adult Fontan Patients. *Front Cardiovasc Med* 2017;4:15.
- 753.Friedrich-Rust, M., F. Schoelzel, et al. Severity of coronary artery disease is associated with non-alcoholic fatty liver disease: A single-blinded prospective mono-center study. *PLoS ONE* 2017;12(10):e0186720.
- 754.Gustot, T. [Treatments of non-alcoholic steatohepatitis]. *Revue Médicale de Bruxelles* 2017;38(4):369-373.
- 755.Hallager, S., S. Ladelund, et al. Liver-related morbidity and mortality in patients with chronic hepatitis C and cirrhosis with and without sustained virologic response. *Clin Epidemiol* 2017;9:501-516.
- 756.Cheah, M. C., A. J. McCullough, et al. Current Modalities of Fibrosis Assessment in Non-alcoholic Fatty Liver Disease. *Journal of Clinical and Translational Hepatology* 2017;5(3):261-271.
- 757.de Ledinghen, V., J. B. Hiriart, et al. Controlled Attenuation Parameter (CAP) with the XL Probe of the Fibroscan(R): A Comparative Study with the M Probe and Liver Biopsy. *Digestive Diseases & Sciences* 2017;62(9):2569-2577.
- 758.Cooper, C. L., H. Hatashita, et al. Direct-Acting Antiviral Therapy Outcomes in Canadian Chronic Hepatitis C Telemedicine Patients. *Annals of Hepatology* 2017;16(6):874-880.
- 759.Cucchetti, A., M. Cescon, et al. Adding Liver Stiffness Measurement to the Routine Evaluation of Hepatocellular Carcinoma Resectability Can Optimize Clinical Outcome. *Ultraschall in der Medizin* 2017;38(5):515-522.
- 760.Desalegn, H., H. Aberra, et al. Are non-invasive fibrosis markers for chronic hepatitis B reliable in sub-Saharan Africa? *Liver International* 2017;37(10):1461-1467.
- 761.Cao, Z., Z. Li, et al. Assessment of serum Golgi protein 73 as a biomarker for the diagnosis of significant fibrosis in patients with chronic HBV infection. *Journal of viral hepatitis* 2017;24 Suppl 1:57-65.
- 762.Chen, J., M. Yin, et al. Diagnostic Performance of MR Elastography and Vibration-controlled Transient Elastography in the Detection of Hepatic Fibrosis in Patients with Severe to Morbid Obesity. *Radiology* 2017;283(2):418-428.
- 763.Colecchia, A., G. Marasco, et al. Usefulness of liver stiffness measurement in predicting hepatic veno-occlusive disease development in patients who undergo HSCT. *Bone Marrow Transplantation* 2017;52(3):494-497.
- 764.Boursier, J., V. de Ledinghen, et al. A stepwise algorithm using an at-a-glance first-line test for the non-invasive diagnosis of advanced liver fibrosis and cirrhosis. *Journal of Hepatology* 2017;66(6):1158-1165.
- 765.Chon, Y. E., J. Y. Park, et al. Improvement of Liver Fibrosis after Long-Term Antiviral Therapy Assessed by Fibroscan in Chronic Hepatitis B Patients With Advanced Fibrosis. *American Journal of Gastroenterology* 2017;112(6):882-891.
- 766.Cepeda, J. A., D. L. Thomas, et al. Increased Mortality Among Persons With Chronic Hepatitis C With Moderate or Severe Liver Disease: A Cohort Study. *Clinical Infectious Diseases* 2017;65(2):235-243.
- 767.Cordero-Ruiz, P., I. Carmona-Soria, et al. Long-term follow-up of patients with chronic hepatitis C treated with alpha-interferon and ribavirin antiviral therapy: clinical and fibrosis impact of treatment response. *European Journal of Gastroenterology & Hepatology* 2017;29(7):792-799.

- 768.Bolognesi, M., M. Di Pascoli, et al. Clinical role of non-invasive assessment of portal hypertension. *World Journal of Gastroenterology* 2017;23(1):1-10.
- 769.Della-Guardia, B., A. S. Evangelista, et al. Diagnostic Accuracy of Transient Elastography for Detecting Liver Fibrosis After Liver Transplantation: A Specific Cut-Off Value Is Really Needed? *Digestive Diseases & Sciences* 2017;62(1):264-272.
- 770.Boyd, A., J. Bottero, et al. Liver fibrosis regression and progression during controlled hepatitis B virus infection among HIV-HBV patients treated with tenofovir disoproxil fumarate in France: a prospective cohort study. *Journal of the International AIDS Society* 2017;20(1):1-12.
- 771.Chen, G. F., J. Ping, et al. [Correlation of liver stiffness measured by FibroTouch and FibroScan with Ishak fibrosis score in patients with chronic hepatitis B]. *Zhonghua Gan Zang Bing Za Zhi* 2017;25(2):145-150.
- 772.Chan, W. K., N. R. Nik Mustapha, et al. Controlled attenuation parameter using the FibroScan(R) XL probe for quantification of hepatic steatosis for non-alcoholic fatty liver disease in an Asian population. *United European Gastroenterology Journal* 2017;5(1):76-85.
- 773.Chong, C. C., G. L. Wong, et al. Liver stiffness measurement predicts high-grade post-hepatectomy liver failure: A prospective cohort study. *Journal of gastroenterology and hepatology* 2017;32(2):506-514.
- 774.Chen, J., M. L. Wang, et al. High value of controlled attenuation parameter predicts a poor antiviral response in patients with chronic hepatitis B. *Hepatobiliary & pancreatic diseases international* 2017;16(4):370-374.
- 775.Cheng, B. C., Y. H. Yen, et al. Transient elastography as a screening tool for liver fibrosis in a large hemodialysis population. *Sci Rep* 2017;7:46458.
- 776.Cales, P., J. Boursier, et al. Liver fibrosis diagnosis by blood test and elastography in chronic hepatitis C: agreement or combination? *Alimentary Pharmacology & Therapeutics* 2017;45(7):991-1003.
- 777.Chen, B., R. A. Schreiber, et al. Response to: Comment on "Assessment of Liver Stiffness in Pediatric Fontan Patients Using Transient Elastography". *Canadian Journal of Gastroenterology & Hepatology* 2017;2017:4217878.
- 778.Chen Yi Mei, S. L. G., A. J. Thompson, et al. Sustained virological response halts fibrosis progression: A long-term follow-up study of people with chronic hepatitis C infection. *PLoS ONE* 2017;12(10):e0185609.
- 779.Couto, I., M. Victoria, et al. Prevalence and predictors for compensated Advanced Chronic Liver Disease (c-ACLD) in patients with chronic Hepatitis Delta Virus (HDV) infection. *PLoS ONE* 2017;12(3):e0174453.
- 780.Dalgard, O., O. Weiland, et al. Sofosbuvir based treatment of chronic hepatitis C genotype 3 infections-A Scandinavian real-life study. *PLoS ONE* 2017;12(7):e0179764.
- 781.Agarwal, S. K., S. Bagchi, et al. Hemodialysis Patients Treated for Hepatitis C Using a Sofosbuvir-based Regimen. *Kidney Int Rep* 2017;2(5):831-835.
- 782.Anand, A. C., S. K. Agarwal, et al. Sofosbuvir and Ribavirin for 24 Weeks Is An Effective Treatment Option for Recurrent Hepatitis C Infection After Living Donor Liver Transplantation. *Journal of Clinical and Experimental Hepatology* 2017;7(3):165-171.
- 783.Ahn, S. J., J. M. Lee, et al. Prospective Validation of Intra- and Interobserver Reproducibility of a New Point Shear Wave Elastographic Technique for Assessing Liver Stiffness in Patients with Chronic Liver Disease. *Korean journal of radiology* 2017;18(6):926-935.
- 784.Bachofner, J. A., P. V. Valli, et al. Direct antiviral agent treatment of chronic hepatitis C results in rapid regression of transient elastography and fibrosis markers fibrosis-4 score and aspartate aminotransferase-platelet ratio index. *Liver International* 2017;37(3):369-376.
- 785.Barone, M., E. Shahini, et al. Letter: can transient elastography and body mass index predict fibrosis regression in HBV patients on long-term nucleoside therapy? *Alimentary Pharmacology & Therapeutics* 2017;45(5):765-766.
- 786.Abera, H., H. Desalegn, et al. Early experiences from one of the first treatment programs for chronic hepatitis B in sub-Saharan Africa. *BMC Infectious Diseases* 2017;17(1):438.
- 787.Bertrais, S., J. Boursier, et al. Prognostic durability of liver fibrosis tests and improvement in predictive performance for mortality by combining tests. *Journal of gastroenterology and hepatology* 2017;32(6):1240-1249.
- 788.Andrade, P., S. Rodrigues, et al. Diagnostic Accuracy of Controlled Attenuation Parameter for Detecting Hepatic Steatosis in Patients with Chronic Liver Disease. *GE Port J Gastroenterol* 2017;24(4):161-168.
- 789.Alempijevic, T., S. Zec, et al. Doppler ultrasonography combined with transient elastography improves the non-invasive assessment of fibrosis in patients with chronic liver diseases. *Medical Ultrasonography Journal* 2017;19(1):7-15.
- 790.Abad, S., A. Vega, et al. Universal Sustained Viral Response to the Combination of Omibitasvir/Paritaprevir/Ritonavir and Dasabuvir with/without Ribavirin in Patients on Hemodialysis Infected with Hepatitis C Virus Genotypes 1 and 4. *Am J Nephrol* 2017;45(3):267-272.
- 791.Agrawal, S., C. L. Hoad, et al. Visual morphometry and three non-invasive markers in the evaluation of

- liver fibrosis in chronic liver disease. Scandinavian journal of gastroenterology 792. 2017;52(1):107-115.
793. Abdel Alem, S., A. Elsharkawy, et al. Improvement of glycemic state among responders to Sofosbuvir-based treatment regimens: Single center experience. Journal of Medical Virology 2017;89(12):2181-2187.
794. Augustin, S., M. Pons, et al. Expanding the Baveno VI criteria for the screening of varices in patients with compensated advanced chronic liver disease. Hepatology 2017;66(6):1980-1988.
795. Andriamananjara, S. F., M. M. Olive, et al. Prevalence of chronic hepatitis B virus infection and infrastructure for its diagnosis in Madagascar: implication for the WHO's elimination strategy. BMC Public Health 2017;17(1):636.
796. Alferink, L. J. M., J. Fittipaldi, et al. Coffee and herbal tea consumption is associated with lower liver stiffness in the general population: The Rotterdam study. Journal of Hepatology 2017;67(2):339-348.
797. Berzigotti, A. Non-invasive evaluation of portal hypertension using ultrasound elastography. Journal of Hepatology 2017;67(2):399-411.
798. Bhati, C., M. O. Idowu, et al. Long-term Outcomes in Patients Undergoing Liver Transplantation for Nonalcoholic Steatohepatitis-Related Cirrhosis. Transplantation 2017;101(8):1867-1874.
799. Bende, F., I. Sporea, et al. Performance of 2D-SWE.GE for predicting different stages of liver fibrosis, using Transient Elastography as the reference method. Medical Ultrasonography Journal 2017;19(2):143-149.
800. Aquil, A., M. M. Jonas, et al. Correlation of Transient Elastography With Severity of Cystic Fibrosis-related Liver Disease. Journal of Pediatric Gastroenterology and Nutrition 2017;64(4):505-511.
801. Bhat, M., M. Tazari, et al. Performance of transient elastography and serum fibrosis biomarkers for non-invasive evaluation of recurrent fibrosis after liver transplantation: A meta-analysis. PLoS ONE 2017;12(9):e0185192.
- 2016**
802. Zheng, Y. X., S. J. Ma, et al. Diagnostic Estimation of Noninvasive Tests for Hepatic Fibrosis in Chronic Hepatitis B Patients Without a Gold Standard. Hepatitis Monthly 2016;16(2):e31983.
803. Wells, M. M. Is transient Elastography a valid screening method for esophageal varices? Saudi Journal of Gastroenterology 2016;22(5):343-344.
804. Voutilainen, S., R. Kivisaari, et al. A Prospective Comparison of Noninvasive Methods in the Assessment of Liver Fibrosis and Esophageal Varices in Pediatric Chronic Liver Diseases. Journal of Clinical Gastroenterology 2016;50(8):658-63.
805. Ying, H. Y., L. G. Lu, et al. Accuracy of transient elastography in the assessment of chronic hepatitis C-related liver cirrhosis. Clinical and investigative medicine 2016;39(5):E150-E160.
806. Weiss, J., M. Rau, et al. Feasibility of liver stiffness measurement in morbidly obese patients undergoing bariatric surgery using XL probe. Scandinavian Journal of Gastroenterology 2016;51(10):1263-8.
807. Yao, J., L. Zhou, et al. Effects of nucleos(t)ide analogs on body composition in HBV-infected men: An age- and BMI-matched, cross-sectional study. Nutrition 2016;32(11-12):1206-10.
808. Udomsinprasert, W., S. Honsawek, et al. Elevation of serum urokinase plasminogen activator receptor and liver stiffness in postoperative biliary atresia. World Journal of Hepatology 2016;8(33):1471-1477.
809. Vuille-Lessard, E., B. Lebouche, et al. Nonalcoholic fatty liver disease diagnosed by transient elastography with controlled attenuation parameter in unselected HIV monoinfected patients. AIDS 2016;30(17):2635-2643.
810. Ulveling, D., S. Le Clerc, et al. A new 3p25 locus is associated with liver fibrosis progression in human immunodeficiency virus/hepatitis C virus-coinfected patients. Hepatology 2016;64(5):1462-1472.
811. Vinikoor, M. J., L. Mulenga, et al. Association between hepatitis B co-infection and elevated liver stiffness among HIV-infected adults in Lusaka, Zambia. Tropical medicine & international health 2016;21(11):1435-1441.
812. Xu, Z. J., J. P. Shi, et al. Evaluating the Relationship Between Metabolic Syndrome and Liver Biopsy-Proven Non-Alcoholic Steatohepatitis in China: A Multicenter Cross-Sectional Study Design. Advances in therapy 2016;33(11):2069-2081.
813. Thiele, M. and A. Krag. Editorial: *in vino veritas* - transient elastography for staging liver fibrosis in alcoholic liver disease. Alimentary Pharmacology & Therapeutics 2016;43(9):1014-5.
814. van der Voort, E. A., E. M. Koehler, et al. Increased Prevalence of Advanced Liver Fibrosis in Patients with Psoriasis: A Cross-sectional Analysis from the Rotterdam Study. Acta dermatovenerologica 2016;96(2):213-7.
815. Thurnheer, M. C., T. R. Schulz, et al. Regional challenges: evaluation of a hepatitis outreach programme using transient elastography (FibroScan) in Victoria. Internal Medicine Journal 2016;46(3):273-81.
816. Yang, N. H., G. S. Yuan, et al. [Entecavir combined with Fufang Biejia Ruangan tablet in treatment of chronic hepatitis B patients with liver fibrosis: 96-week efficacy analyses]. Nan Fang Yi Ke Da Xue Xue Bao 2016;36(6):775-9.
817. Yosry, A., R. Fouad, et al. FibroScan, APRI, FIB4, and GUCI: Role in prediction of fibrosis and response to

- therapy in Egyptian patients with HCV infection. *Arab journal of gastroenterology* 2016;17(2):78-83.
818. Valcour, V. G., L. H. Rubin, et al. Liver Fibrosis Linked to Cognitive Performance in HIV and Hepatitis C. *Journal of Acquired Immune Deficiency Syndromes* 2016;72(3):266-73.
819. Thiele, M., S. Detlefsen, et al. Transient and 2-Dimensional Shear-Wave Elastography Provide Comparable Assessment of Alcoholic Liver Fibrosis and Cirrhosis. *Gastroenterology* 2016;150(1):123-33.
820. Xu, L., P. Li, et al. [Value of controlled attenuation parameter in diagnosis of fatty liver using FibroScan]. *Zhonghua Gan Zang Bing Za Zhi* 2016;24(2):108-13.
821. Wang, J. H., Y. H. Yen, et al. Liver stiffness-based score in hepatoma risk assessment for chronic hepatitis C patients after successful antiviral therapy. *Liver International* 2016;36(12):1793-1799.
822. Wang, Y., Y. Zeng, et al. Hypertension and non-alcoholic fatty liver disease proven by transient elastography. *Hepatology Research* 2016;46(13):1304-1310.
824. Wong, G. L., R. Kwok, et al. Measuring spleen stiffness to predict varices in chronic hepatitis B cirrhotic patients with or without receiving non-selective beta-blockers. *Journal of Digestive Diseases* 2016;17(8):538-546.
825. Yamamoto, A., N. Nishida, et al. Prediction for Improvement of Liver Function after Balloon-Occluded Retrograde Transvenous Obliteration for Gastric Varices to Manage Portosystemic Shunt Syndrome. *Journal of Vascular and Interventional Radiology* 2016;27(8):1160-7.
826. Zhang, R. N., R. D. Zheng, et al. APOC3 rs2070666 Is Associated with the Hepatic Steatosis Independently of PNPLA3 rs738409 in Chinese Han Patients with Nonalcoholic Fatty Liver Diseases. *Digestive Diseases & Sciences* 2016;61(8):2284-93.
827. Van Biervliet, S., H. Verdievel, et al. Longitudinal Transient Elastography Measurements Used in Follow-up for Patients with Cystic Fibrosis. *Ultrasound in Medicine and Biology* 2016;42(4):848-54.
828. Yoshida, S., M. Shimada, et al. Hepatobiliary and Pancreatic: Elastography assessment in AL hepatic amyloidosis with no fibrosis. *Journal of Gastroenterology & Hepatology* 2016;31(4):708.
829. Zhang, D., M. Chen, et al. The Clinical Application of Acoustic Radiation Force Impulse Imaging and Transient Elastography in Chronic Hepatitis B. *Ultrasound in Medicine and Biology* 2016;42(4):1026.
830. Tokuhara, D., Y. Cho, et al. Transient Elastography-Based Liver Stiffness Age-Dependently Increases in Children. *PLoS ONE* 2016;11(11):e0166683.
831. Xue, X. and S. Cai. Comment on "Assessment of Liver Stiffness in Pediatric Fontan Patients Using Transient Elastography". *Canadian Journal of Gastroenterology & Hepatology* 2016;2016:9343960.
832. Tapper, E. B., M. G. Hunink, et al. Cost-Effectiveness Analysis: Risk Stratification of Nonalcoholic Fatty Liver Disease (NAFLD) by the Primary Care Physician Using the NAFLD Fibrosis Score. *PLoS ONE* 2016;11(2):e0147237.
833. Mikolasevic, I., S. Milic, et al. Factors associated with significant liver steatosis and fibrosis as assessed by transient elastography in patients with one or more components of the metabolic syndrome. *Journal of diabetes and its complications* 2016;30(7):1347-53.
834. Qu, Y., T. Li, et al. A Beginning or the End? A Meta-analysis to Assess the Diagnostic Accuracy of Transient Elastography for the Prediction of Esophageal Varices. *Saudi Journal of Gastroenterology* 2016;22(5):345-352.
835. Ramos-Lopez, O., S. Roman, et al. CD36 genetic variation, fat intake and liver fibrosis in chronic hepatitis C virus infection. *World Journal of Hepatology* 2016;8(25):1067-74.
836. Medrano, L. M., N. Rallon, et al. Relationship of TRIM5 and TRIM22 polymorphisms with liver disease and HCV clearance after antiviral therapy in HIV/HCV coinfecting patients. *Journal of translational medicine* 2016;14:257.
837. Lim, T. H., E. Gane, et al. HBsAg loss in a New Zealand community study with 28-year follow-up: rates, predictors and long-term outcomes. *Hepatology International* 2016;10(5):829-37.
838. Maruyama, H., K. Kobayashi, et al. Two-dimensional shear wave elastography with propagation-based reliability assessment for grading hepatic fibrosis and portal hypertension. *Journal of Hepato-biliary-pancreatic Sciences* 2016;23(9):595-602.
839. Mehta, S. H., A. M. McFall, et al. Morbidity and Mortality Among Community-Based People Who Inject Drugs With a High Hepatitis C and Human Immunodeficiency Virus Burden in Chennai, India. *Open Forum Infectious Diseases* 2016;3(3):ofw121.
840. Park, S. H., S. Y. Kim, et al. What we need to know when performing and interpreting US elastography. *Clinical and Molecular Hepatology* 2016;22(3):406-414.
841. Pokora, A., S. Kiciak, et al. Assessing the consistency of liver fibrosis results obtained through biopsy and transient elastography in patients suffering from chronic hepatitis C. *Clin Exp Hepatol* 2016;2(3):109-111.
842. Shen, F. F. and L. G. Lu. Advances in noninvasive methods for diagnosing nonalcoholic fatty liver disease. *Journal of Digestive Diseases* 2016;17(9):565-571.
843. Sultanik, P., L. Kramer, et al. The relationship between liver stiffness measurement and outcome in patients with chronic hepatitis C and cirrhosis: a retrospective longitudinal hospital study.

- Alimentary Pharmacology & Therapeutics 2016;44(5):505-13.
844. Mandorfer, M., K. Kozbial, et al. Sustained virologic response to interferon-free therapies ameliorates HCV-induced portal hypertension. *Journal of Hepatology* 2016;65(4):692-9.
845. Matta, B., T. H. Lee, et al. Use of Non-invasive Testing to Stage Liver Fibrosis in Patients with HIV. *Current HIV and AIDS reports* 2016;13(5):279-88.
846. Njei, B., T. R. McCarty, et al. Use of transient elastography in patients with HIV-HCV coinfection: A systematic review and meta-analysis. *Journal of Gastroenterology & Hepatology* 2016;31(10):1684-1693.
847. Perez-Latorre, L., A. Rivero-Juarez, et al. Prognostic Value of Transient Elastography in Human Immunodeficiency Virus-Infected Patients With Chronic Hepatitis C. *Open Forum Infectious Diseases* 2016;3(4):ofw212.
848. Pons, M., M. Simon-Talero, et al. Basal values and changes of liver stiffness predict the risk of disease progression in compensated advanced chronic liver disease. *Digestive & Liver Disease* 2016;48(10):1214-9.
849. Sasso, M., Y. Liu, et al. AdipoScan: A Novel Transient Elastography-Based Tool Used to Non-Invasively Assess Subcutaneous Adipose Tissue Shear Wave Speed in Obesity. *Ultrasound in Medicine and Biology* 2016;42(10):2401-13.
850. Shan, R., H. Yin, et al. Influencing factors of transient elastography in detecting liver stiffness. *Experimental and therapeutic medicine* 2016;12(4):2302-2306.
851. Srinivasa Babu, A., M. L. Wells, et al. Elastography in Chronic Liver Disease: Modalities, Techniques, Limitations, and Future Directions. *Radiographics* 2016;36(7):1987-2006.
852. Ronit, A., J. Haissman, et al. Copenhagen comorbidity in HIV infection (COCOMO) study: a study protocol for a longitudinal, non-interventional assessment of non-AIDS comorbidity in HIV infection in Denmark. *BMC Infectious Diseases* 2016;16(1):713.
853. Macias, J., L. M. Real, et al. Changes in liver steatosis evaluated by transient elastography with the controlled attenuation parameter in HIV-infected patients. *HIV Medicine* 2016;17(10):766-773.
854. Maurice, J. B., E. Brodkin, et al. Validation of the Baveno VI criteria to identify low risk cirrhotic patients not requiring endoscopic surveillance for varices. *Journal of Hepatology* 2016;65(5):899-905.
855. Montano-Loza, A. J., R. B. Thandassery, et al. Targeting Hepatic Fibrosis in Autoimmune Hepatitis. *Digestive Diseases & Sciences* 2016;61(11):3118-3139.
856. O, E. A., M. Buchter, et al. Performance and Utility of Transient Elastography and Non-Invasive Markers of Liver Fibrosis in Patients with Autoimmune Hepatitis: A Single Centre Experience. *Hepatitis Monthly* 2016;16(11):e40737.
857. San Segundo, D., C. Alonso, et al. De Novo Donor-Specific Anti-Human Leukocyte Antigen Antibody Detection in Long-Term Adult Liver Transplantation. *Transplantation Proceedings* 2016;48(9):2980-2982.
858. Seto, W. K., J. Fung, et al. Body-mass index is associated with fibrosis regression during long-term nucleoside analogue therapy in chronic hepatitis B. *Alimentary Pharmacology & Therapeutics* 2016;44(10):1071-1079.
859. Shan, Q. Y., B. X. Liu, et al. Elastography of shear wave speed imaging for the evaluation of liver fibrosis: A meta-analysis. *Hepatology Research* 2016;46(12):1203-1213.
860. Staugaard, B., P. B. Christensen, et al. Feasibility of transient elastography versus real-time two-dimensional shear wave elastography in difficult-to-scan patients. *Scandinavian Journal of Gastroenterology* 2016;51(11):1354-9.
861. Liang, X. E., L. Dai, et al. Combining routine markers improves the accuracy of transient elastography for hepatitis B cirrhosis detection. *Digestive & Liver Disease* 2016;48(5):512-8.
862. Pavlov, C., G. Casazza, et al. Editorial: in vino veritas - transient elastography for staging liver fibrosis in alcoholic liver disease - authors' reply. *Alimentary Pharmacology & Therapeutics* 2016;43(9):1015-6.
863. Stockdale, A. J., R. O. Phillips, et al. The gamma-glutamyl transpeptidase to platelet ratio (GPR) shows poor correlation with transient elastography measurements of liver fibrosis in HIV-positive patients with chronic hepatitis B in West Africa. Response to: 'The gamma-glutamyl transpeptidase to platelet ratio (GPR) predicts significant liver fibrosis and cirrhosis in patients with chronic HBV infection in West Africa' by Lemoine et al. *Gut* 2016;65(5):882-4.
864. Tapper, E. B., T. Challies, et al. The Performance of Vibration Controlled Transient Elastography in a US Cohort of Patients With Nonalcoholic Fatty Liver Disease. *American Journal of Gastroenterology* 2016;111(5):677-84.
865. Mateos-Munoz, B., E. Garcia-Martin, et al. GC Gene Polymorphism and Unbound Serum Retinol-Binding Protein 4 Are Related to the Risk of Insulin Resistance in Patients With Chronic Hepatitis C: A Prospective Cross-Sectional Study. *Medicine (Baltimore)* 2016;95(10):e3019.
866. Pavlov, C. S., G. Casazza, et al. Systematic review with meta-analysis: diagnostic accuracy of transient elastography for staging of fibrosis in people with alcoholic liver disease. *Alimentary Pharmacology & Therapeutics* 2016;43(5):575-85.
867. Saez-Royuela, F., P. Linares, et al. Evaluation of advanced fibrosis measured by transient elastography after hepatitis C virus protease

- inhibitor-based triple therapy. European Journal of Gastroenterology & Hepatology 2016;28(3):305-12.
- 868.Seo, Y. S., M. N. Kim, et al. Risk Assessment of Hepatocellular Carcinoma Using Transient Elastography Vs. Liver Biopsy in Chronic Hepatitis B Patients Receiving Antiviral Therapy. MEDICINE 2016;95(12):e2985.
- 869.Sporea, I., R. Sirli, et al. Feasibility of Transient Elastography with M and XL probes in real life. Medical Ultrasonography Journal 2016;18(1):7-10.
- 870.Taouli, B. and L. Serfaty. Magnetic Resonance Imaging/Elastography Is Superior to Transient Elastography for Detection of Liver Fibrosis and Fat in Nonalcoholic Fatty Liver Disease. Gastroenterology 2016;150(3):553-6.
- 871.Lewindon, P. J., F. Balouch, et al. Transient liver elastography in unsedated control children: Impact of age and intercurrent illness. Journal of Paediatrics and Child Health 2016;52(6):637-42.
- 872.Papapostoli, I., F. Lammert, et al. Effect of Short-Term Vitamin D Correction on Hepatic Steatosis as Quantified by Controlled Attenuation Parameter (CAP). Journal of Gastrointestinal & Liver Diseases 2016;25(2):175-81.
- 873.Shen, F., R. D. Zheng, et al. [Value of a two-step approach with cytokeratin-18 and controlled attenuation parameter in noninvasive differential diagnosis of nonalcoholic steatohepatitis]. Zhonghua Gan Zang Bing Za Zhi 2016;24(6):429-34.
- 874.Sporea, I., R. Mare, et al. Liver Stiffness Evaluation by Transient Elastography in Type 2 Diabetes Mellitus Patients with Ultrasound-proven Steatosis. Journal of Gastrointestinal & Liver Diseases 2016;25(2):167-74.
- 875.Mobarak, L., M. M. Nabeel, et al. Real-time elastography as a noninvasive assessment of liver fibrosis in chronic hepatitis C Egyptian patients: a prospective study. Annals of gastroenterology 2016;29(3):358-62.
- 876.Mateos-Munoz, B., M. J. Devesa-Medina, et al. The relation of fibrosis stage with nutritional deficiencies and bioelectrical impedance analysis of body composition in patients with chronic hepatitis C. Annals of Hepatology 2016;15(4):492-500.
- 877.Shin, J. W. and N. H. Park. [Clinical Application of Non-invasive Diagnostic Tests for Liver Fibrosis]. Korean Journal of Gastroenterology 2016;68(1):4-9.
- 878.Li, T., Y. Qu, et al. Evaluation of large esophageal varices in cirrhotic patients by transient elastography: a meta-analysis. Revista Española de Enfermedades Digestivas 2016;108.
- 879.Sun, Y. M., X. N. Wu, et al. [A discussion of the methods for the assessment of chronic hepatitis B-related liver fibrosis regression]. Zhonghua Gan Zang Bing Za Zhi 2016;24(7):552-554.
- 880.Leite, N. C., C. A. Villela-Nogueira, et al. Increasing aortic stiffness is predictive of advanced liver fibrosis in patients with type 2 diabetes: the Rio-
- T2DM cohort study. Liver International 2016;36(7):977-85.
- 881.Stasi, C. and S. Milani. Non-invasive assessment of liver fibrosis: Between prediction/prevention of outcomes and cost-effectiveness. World Journal of Gastroenterology 2016;22(4):1711-20.
- 882.Lim, Y. L., E. Choi, et al. Clinical Implications of the Serum Apelin Level on Portal Hypertension and Prognosis of Liver Cirrhosis. Gut Liver 2016;10(1):109-16.
- 883.Lee, H. W., W. Kang, et al. Red cell volume distribution width-to-platelet ratio in assessment of liver fibrosis in patients with chronic hepatitis B. Liver International 2016;36(1):24-30.
- 884.Rosso, C., G. P. Caviglia, et al. Cytokeratin 18-Aspartate396 apoptotic fragment for fibrosis detection in patients with non-alcoholic fatty liver disease and chronic viral hepatitis. Digestive & Liver Disease 2016;48(1):55-61.
- 885.Sasso, M., S. Audiere, et al. Liver Steatosis Assessed by Controlled Attenuation Parameter (CAP) Measured with the XL Probe of the FibroScan: A Pilot Study Assessing Diagnostic Accuracy. Ultrasound in Medicine and Biology 2016;42(1):92-103.
- 886.Strasser, M., A. Stadlmayr, et al. Natural course of subjects with elevated liver tests and normal liver histology. Liver International 2016;36(1):119-25.
- 887.Qu, Y. and L. G. Lu. [New advances in diagnosis of nonalcoholic fatty liver disease]. Zhonghua Gan Zang Bing Za Zhi 2016;24(2):85-8.
- 888.Li, Y., Y. S. Huang, et al. Systematic review with meta-analysis: the diagnostic accuracy of transient elastography for the staging of liver fibrosis in patients with chronic hepatitis B. Alimentary Pharmacology & Therapeutics 2016;43(4):458-69.
- 889.Merli, M., A. Castagna, et al. Diagnostic accuracy of APRI, FIB-4 and Forns for the detection of liver cirrhosis in HIV/HCV-coinfected patients. New Microbiologica 2016;39(2):110-3.
- 890.Nam, T. H., B. J. Park, et al. Non-invasive assessment of liver fibrosis using acoustic structure quantification: comparison with transient elastography in chronic viral hepatitis. Abdominal Radiology 2016;41(2):239-47.
- 891.Perazzo, H., F. F. Fernandes, et al. Learning curve and intra/interobserver agreement of transient elastography in chronic hepatitis C patients with or without HIV co-infection. Clinics and Research in Hepatology and Gastroenterology 2016;40(1):73-82.
- 892.Piscaglia, F., V. Salvatore, et al. Ultrasound Shear Wave Elastography for Liver Disease. A Critical Appraisal of the Many Actors on the Stage. Ultraschall in der Medizin 2016;37(1):1-5.
- 893.Stefanescu, A. C., T. L. Pop, et al. Transient elastography of the liver in children with Wilson's disease: Preliminary results. Journal of Clinical Ultrasound 2016;44(2):65-71.

894. Sultanik, P., P. Sogni, et al. Patients with chronic hepatitis B should be screened for hepatocellular carcinoma regardless of liver stiffness measurement. *Hepatology* 2016;63(2):672.
895. Logan, S., A. Rodger, et al. Prevalence of significant liver disease in human immunodeficiency virus-infected patients exposed to Didanosine: A cross sectional study. *World Journal of Hepatology* 2016;8(36):1623-1628.
896. Rausch, V., T. Peccerella, et al. Primary liver injury and delayed resolution of liver stiffness after alcohol detoxification in heavy drinkers with the PNPLA3 variant I148M. *World Journal of Hepatology* 2016;8(35):1547-1556.
897. Morisco, F., M. Masarone, et al. Impact of Telaprevir in HCV Patients with Cirrhosis and RVR: Real-Life Data from Boceprevir or Telaprevir based "Triple Therapy" Experience in Southern Italy. *Reviews on recent clinical trials* 2016;11(4):306-316.
898. Li, Y., Q. Cai, et al. Development of algorithms based on serum markers and transient elastography for detecting significant fibrosis and cirrhosis in chronic hepatitis B patients: Significant reduction in liver biopsy. *Hepatology Research* 2016;46(13):1367-1379.
899. Lombardi, R., H. Sambatakou, et al. Prevalence and predictors of liver steatosis and fibrosis in unselected patients with HIV mono-infection. *Digestive & Liver Disease* 2016;48(12):1471-1477.
900. Ozturker, C., E. Karagoz, et al. Non-Invasive Evaluation of Liver Fibrosis: 2-D Shear Wave Elastography, Transient Elastography or Acoustic Radiation Force Impulse Imaging? *Ultrasound in Medicine and Biology* 2016;42(12):3052.
901. Rinaldi, L., G. Valente, et al. Serial Liver Stiffness Measurements and Monitoring of Liver-Transplanted Patients in a Real-Life Clinical Practice. *Hepatitis Monthly* 2016;16(12):e41162.
902. Shur, N. F., Y. Tan, et al. Non-viral liver disease burden in HIV-monoinfected individuals: a longitudinal observational retrospective cohort study. *AIDS Care* 2016;28(12):1522-1527.
903. Taneja, S., S. Tohra, et al. Noninvasive Assessment of Liver Fibrosis By Transient Elastography and FIB4/APRI for Prediction of Treatment Response in Chronic Hepatitis C-An Experience from a Tertiary Care Hospital. *Journal of Clinical and Experimental Hepatology* 2016;6(4):282-290.
904. Mikolasevic, I., L. Orlic, et al. Transient elastography (FibroScan((R))) with controlled attenuation parameter in the assessment of liver steatosis and fibrosis in patients with nonalcoholic fatty liver disease - Where do we stand? *World Journal of Gastroenterology* 2016;22(32):7236-51.
905. Mendes, L. C., P. A. Ferreira, et al. Transient elastography and APRI score: looking at false positives and false negatives. Diagnostic performance and association to fibrosis staging in chronic hepatitis C. *Brazilian journal of medical and biological research* 2016;49(9):e5432.
906. Lemoine, M., Y. Shimakawa, et al. The gamma-glutamyl transpeptidase to platelet ratio (GPR) predicts significant liver fibrosis and cirrhosis in patients with chronic HBV infection in West Africa. *Gut* 2016;65(8):1369-76.
907. Lui, G., V. W. Wong, et al. Liver fibrosis and fatty liver in Asian HIV-infected patients. *Alimentary Pharmacology & Therapeutics* 2016;44(4):411-21.
908. Petta, S. and A. Craxi. Response to Eddowas and colleagues. *Hepatology* 2016;64(2):701.
909. Rezende, R. E., S. M. Duarte, et al. Randomized clinical trial: benefits of aerobic physical activity for 24 weeks in postmenopausal women with nonalcoholic fatty liver disease. *Menopause* 2016;23(8):876-83.
910. Shibata, S., T. Umemura, et al. Liver stiffness-spleen size-to-platelet ratio risk score identifies esophageal varices in Japanese patients with chronic hepatitis C. *Hepatology Research* 2016;46(9):884-9.
911. Mandorfer, M., P. Schwabl, et al. Interferon-free treatment with sofosbuvir/daclatasvir achieves sustained virologic response in 100% of HIV/hepatitis C virus-coinfected patients with advanced liver disease. *Aids* 2016;30(7):1039-47.
912. Swanson, S., Y. Ma, et al. Association of HIV, Hepatitis C Virus, and Liver Fibrosis Severity With the Enhanced Liver Fibrosis Score. *The Journal of Infectious Diseases* 2016;213(7):1079-86.
913. Liu, X. Y., L. N. Ma, et al. Combined detection of liver stiffness and C-reactive protein in patients with hepatitis B virus-related liver cirrhosis, with and without hepatocellular carcinoma. *Molecular and Clinical Oncology* 2016;4(4):587-590.
914. Moossavi, S., S. Besharat, et al. Inverse Association of Plasma Level of Glutathione Peroxidase with Liver Fibrosis in Chronic Hepatitis B: Potential Role of Iron. *Middle East journal of digestive diseases* 2016;8(2):122-30.
915. Ozturker, C. and E. Karagoz. Acoustic Radiation Force Impulse Imaging or Transient Elastography in Chronic Hepatitis B? *Ultrasound in Medicine and Biology* 2016;42(4):1026.
916. Mikolasevic, I., S. Milic, et al. Nonalcoholic Fatty Liver Disease (NAFLD)-A New Cardiovascular Risk Factor in Peritoneal Dialysis Patients. *Peritoneal Dialysis International* 2016;36(4):427-432.
917. Lee, H. W., S. Y. Park, et al. Discrimination of Nonalcoholic Steatohepatitis Using Transient Elastography in Patients with Nonalcoholic Fatty Liver Disease. *PLoS ONE* 2016;11(6):e0157358.
918. Lemaitre, C., S. Dominique, et al. Relevance of 3D Cholangiography and Transient Elastography to Assess Cystic Fibrosis-Associated Liver Disease? *Canadian Respiratory Journal* 2016;2016:4592702.

919. Lombardi, R., G. Pisano, et al. Role of Serum Uric Acid and Ferritin in the Development and Progression of NAFLD. *International Journal of Molecular Sciences* 2016;17(4).
920. Ma, X., L. Wang, et al. Spleen Stiffness Is Superior to Liver Stiffness for Predicting Esophageal Varices in Chronic Liver Disease: A Meta-Analysis. *PLoS ONE* 2016;11(11):e0165786.
921. Nguyen Truong, T., D. Laureillard, et al. High Proportion of HIV-HCV Coinfected Patients with Advanced Liver Fibrosis Requiring Hepatitis C Treatment in Haiphong, Northern Vietnam (ANRS 12262). *PLoS ONE* 2016;11(5):e0153744.
922. Ntagirabiri, R., B. Munezero, et al. [Hepatitis B virus genotypes and evolutionary markers in chronic HBsAG patients in Bujumbura]. *The Pan African medical journal* 2016;23:95.
923. Nunez-Torres, R., J. Macias, et al. The PNPLA3 Genetic Variant rs738409 Influences the Progression to Cirrhosis in HIV/Hepatitis C Virus Coinfected Patients. *PLoS ONE* 2016;11(12):e0168265.
924. Papadopoulos, N., M. Deutsch, et al. Simeprevir and Sofosbuvir Combination Treatment in a Patient with HCV Cirrhosis and HbS Beta 0-Thalassemia: Efficacy and Safety despite Baseline Hyperbilirubinemia. *Case Reports in Hematology* 2016;2016:7635128.
925. Pongpit, J., S. Porntharukchareon, et al. Liver Stiffness Measurement in Psoriasis: Do Metabolic or Disease Factors Play the Important Role? *BioMed Research International* 2016;2016:7963972.
926. Poynard, T., T. Pham, et al. Real-Time Shear Wave versus Transient Elastography for Predicting Fibrosis: Applicability, and Impact of Inflammation and Steatosis. A Non-Invasive Comparison. *PLoS ONE* 2016;11(10):e0163276.
927. Shibata, S., S. Joshi, et al. Liver stiffness-spleen size-to-platelet ratio risk score detects esophageal varices in chronic liver disease. *Springerplus* 2016;5(1):998.
928. Solomon, S. S., A. K. Srikrishnan, et al. Burden of Liver Disease among Community-Based People Who Inject Drugs (PWID) in Chennai, India. *PLoS ONE* 2016;11(1):e0147879.
929. Lee, H. W., Y. E. Chon, et al. Predicting Liver-Related Events Using Transient Elastography in Chronic Hepatitis C Patients with Sustained Virological Response. *Gut Liver* 2016;10(3):429-36.
930. Koller, T., Z. Piestanska, et al. [Hepatic transit times and liver elasticity compared with meld in predicting a 1 year adverse clinical outcome of a clinically diagnosed cirrhosis]. *Vnitrni lekarstvi* 2016;62(5):351-9.
931. Giunta, M., D. Conte, et al. Role of spleen elastography in patients with chronic liver diseases. *World Journal of Gastroenterology* 2016;22(35):7857-67.
932. Blake, L., R. V. Duarte, et al. Decision analytic model of the diagnostic pathways for patients with suspected non-alcoholic fatty liver disease using non-invasive transient elastography and multiparametric magnetic resonance imaging. *BMJ Open* 2016;6(9):e010507.
933. Ko, B. J., Y. S. Kim, et al. Relationship between 25-Hydroxyvitamin D Levels and Liver Fibrosis as Assessed by Transient Elastography in Patients with Chronic Liver Disease. *Gut Liver* 2016;10(5):818-25.
934. Bihari, C., A. Rastogi, et al. Quantitative fibrosis estimation by image analysis predicts development of decompensation, composite events and defines event-free survival in chronic hepatitis B patients. *Human Pathology* 2016;55:63-71.
935. Boursier, J., J. Vergniol, et al. Diagnostic accuracy and prognostic significance of blood fibrosis tests and liver stiffness measurement by FibroScan in non-alcoholic fatty liver disease. *Journal of Hepatology* 2016;65(3):570-8.
936. Chen, J., D. Wu, et al. Controlled attenuation parameter for the detection of hepatic steatosis in patients with chronic hepatitis B. *Infectious diseases* 2016;48(9):670-5.
937. Chien, C. H., Y. L. Lin, et al. Transient Elastography for Spleen Stiffness Measurement in Patients With Cirrhosis: Role in Degree of Thrombocytopenia. *Journal of ultrasound in medicine* 2016;35(9):1849-57.
938. Ciocan, D., A. Lebrun, et al. Adaptation of liver stiffness measurement depth in bariatric surgery candidates with suspected nonalcoholic fatty liver disease. *European Journal of Gastroenterology & Hepatology* 2016;28(9):1014-20.
939. de Barros, F., S. Setubal, et al. The Correlation Between Obesity-Related Diseases and Non-alcoholic Fatty Liver Disease in Women in the Pre-operative Evaluation for Bariatric Surgery Assessed by Transient Hepatic Elastography. *Obesity surgery* 2016;26(9):2089-2097.
940. Fujimori, N., N. Tanaka, et al. Controlled attenuation parameter is correlated with actual hepatic fat content in patients with non-alcoholic fatty liver disease with none-to-mild obesity and liver fibrosis. *Hepatology Research* 2016;46(10):1019-27.
941. Iurlo, A., D. Cattaneo, et al. Reply To: Assessment of Liver and Spleen Stiffness in Patients With Myelofibrosis Using FibroScan and Shear Wave Elastography. *Ultrasound Quarterly* 2016;32(3):317.
942. Jimenez-Sousa, M. A., J. Berenguer, et al. IL15 polymorphism is associated with advanced fibrosis, inflammation-related biomarkers and virological response in human immunodeficiency virus/hepatitis C virus coinfection. *Liver International* 2016;36(9):1258-66.
943. Hashimoto, Y., T. Osaka, et al. The relationship between hepatic steatosis and skeletal muscle mass

- index in men with type 2 diabetes. *Endocrine Journal* 2016;63(10):877-884.
944. Lee, H. W. and S. H. Ahn. Prediction models of hepatocellular carcinoma development in chronic hepatitis B patients. *World Journal of Gastroenterology* 2016;22(37):8314-8321.
945. Deorsola, L., E. Aidala, et al. Liver stiffness modifications shortly after total cavopulmonary connection. *Interactive Cardiovascular and Thoracic Surgery* 2016;23(4):513-8.
946. Hartl, J., U. Denzer, et al. Transient elastography in autoimmune hepatitis: Timing determines the impact of inflammation and fibrosis. *Journal of Hepatology* 2016;65(4):769-75.
947. Huang, Y. Q. Recent advances in the diagnosis and treatment of primary biliary cholangitis. *World Journal of Hepatology* 2016;8(33):1419-1441.
948. Berge, E., E. Oton, et al. Predictors of Poor Prognosis in Recurrent Hepatitis C After Liver Transplantation. *Transplantation Proceedings* 2016;48(9):2997-2999.
949. Cheng, J. Y., V. W. Wong, et al. Metabolic syndrome increases cardiovascular events but not hepatic events and death in patients with chronic hepatitis B. *Hepatology* 2016;64(5):1507-1517.
950. Conti, F., R. Vukotic, et al. Transient elastography in healthy subjects and factors influencing liver stiffness in non-alcoholic fatty liver disease: An Italian community-based population study. *Digestive & Liver Disease* 2016;48(11):1357-1363.
951. Frank Wolf, M., D. Peleg, et al. Correlation between changes in liver stiffness and preeclampsia as shown by transient elastography. *Hypertension in Pregnancy* 2016;35(4):536-541.
952. Gathercole, L. L., J. M. Hazlehurst, et al. Advanced non-alcoholic fatty liver disease and adipose tissue fibrosis in patients with Alstrom syndrome. *Liver International* 2016;36(11):1704-1712.
953. Kaya, E., D. Demir, et al. Prevalence of hepatic steatosis in apparently healthy medical students: a transient elastography study on the basis of a controlled attenuation parameter. *European Journal of Gastroenterology & Hepatology* 2016;28(11):1264-7.
954. Keane, M. G., C. Hensher, et al. Improving the identification and monitoring of cirrhosis. *Practitioner* 2016;260(1798):25-9.
955. Kelly, D., H. J. Verkade, et al. Late graft hepatitis and fibrosis in pediatric liver allograft recipients: Current concepts and future developments. *Liver Transplantation* 2016;22(11):1593-1602.
956. Kim, H. S., B. K. Kim, et al. Association Between Level of Fibrosis, Rather Than Antiviral Regimen, and Outcomes of Patients With Chronic Hepatitis B. *Clinical Gastroenterology & Hepatology* 2016;14(11):1647-1656 e6.
957. Ferraioli, G., R. Lissandrini, et al. Liver stiffness assessed by transient elastography in patients with beta thalassaemia major. *Annals of Hepatology* 2016;15(3):410-417.
958. Jonas, M. C., C. V. Rodriguez, et al. Streamlining Screening to Treatment: The Hepatitis C Cascade of Care at Kaiser Permanente Mid-Atlantic States. *Clinical Infectious Diseases* 2016;62(10):1290-6.
959. Duan, S. H., Z. Y. Bao, et al. [Efficacy Observation of Yiguanjian Decoction Combined Adefovir Dipivoxil Tablet in Treating HBeAg Negative Chronic Viral Hepatitis B Active Compensated Liver Cirrhosis Patients]. *Zhongguo Zhong Xi Yi Jie He Za Zhi* 2016;36(5):535-8.
960. Dyvorne, H. A., G. H. Jajamovich, et al. Prospective comparison of magnetic resonance imaging to transient elastography and serum markers for liver fibrosis detection. *Liver International* 2016;36(5):659-66.
961. Huang, R., N. Jiang, et al. Fibroscan improves the diagnosis sensitivity of liver fibrosis in patients with chronic hepatitis B. *Experimental and therapeutic medicine* 2016;11(5):1673-1677.
962. Kaswala, D. H., M. Lai, et al. Fibrosis Assessment in Nonalcoholic Fatty Liver Disease (NAFLD) in 2016. *Digestive Diseases & Sciences* 2016;61(5):1356-64.
963. Kim, S. U., B. K. Kim, et al. Transient Elastography is Superior to FIB-4 in Assessing the Risk of Hepatocellular Carcinoma in Patients With Chronic Hepatitis B. *MEDICINE* 2016;95(20):e3434.
964. Kumagai, E., K. Korenaga, et al. Appropriate use of virtual touch quantification and FibroScan(R) M and XL probes according to the skin capsular distance. *Journal of Gastroenterology & Hepatology* 2016;51(5):496-505.
965. Conti, C. B., F. Cavalcoli, et al. Ultrasound elastographic techniques in focal liver lesions. *World Journal of Gastroenterology* 2016;22(9):2647-56.
966. Bernuth, S., E. Yagmur, et al. Early changes in dynamic biomarkers of liver fibrosis in hepatitis C virus-infected patients treated with sofosbuvir. *Digestive & Liver Disease* 2016;48(3):291-7.
967. Caston, J. J., L. Castells, et al. Impact of Cytomegalovirus Infection on Severe Hepatitis C Recurrence in Patients Undergoing Liver Transplantation. *Transplantation* 2016;100(3):593-9.
968. Cheng, P. N., Y. C. Chiu, et al. The Application of Liver Stiffness Measurement in Residents Without Overt Liver Diseases Through a Community-Based Screening Program. *MEDICINE* 2016;95(12):e3193.
969. Crespo, G., G. Castro-Narro, et al. Usefulness of liver stiffness measurement during acute cellular rejection in liver transplantation. *Liver Transplantation* 2016;22(3):298-304.
970. Fitzpatrick, J. A., J. U. Kim, et al. Changes in Liver Volume in Patients with Chronic Hepatitis C Undergoing Antiviral Therapy. *Journal of Clinical and Experimental Hepatology* 2016;6(1):15-20.

- 971.Hollande, C., V. Mallet, et al. Impact of Obliterative Portal Venopathy Associated With Human Immunodeficiency Virus. *Medicine (Baltimore)* 2016;95(11):e3081.
- 972.Hukkanen, M., R. Kivisaari, et al. Transient elastography and aspartate aminotransferase to platelet ratio predict liver injury in paediatric intestinal failure. *Liver International* 2016;36(3):361-9.
- 973.Imajo, K., T. Kessoku, et al. Magnetic Resonance Imaging More Accurately Classifies Steatosis and Fibrosis in Patients With Nonalcoholic Fatty Liver Disease Than Transient Elastography. *Gastroenterology* 2016;150(3):626-637 e7.
- 974.Kim, J. S., C. W. Kim, et al. Hepatic hydrothorax without ascites as the first sign of liver cirrhosis. *Respirology Case Reports* 2016;4(1):16-8.
- 975.Belei, O., I. Sporea, et al. Comparison of three ultrasound based elastographic techniques in children and adolescents with chronic diffuse liver diseases. *Medical Ultrasonography Journal* 2016;18(2):145-50.
- 976.Cassinotto, C., J. Boursier, et al. Liver stiffness in nonalcoholic fatty liver disease: A comparison of supersonic shear imaging, FibroScan, and ARFI with liver biopsy. *Hepatology* 2016;63(6):1817-27.
- 977.Crespo, G., M. Gambato, et al. Early non-invasive selection of patients at high risk of severe hepatitis C recurrence after liver transplantation. *Transplant Infectious Disease* 2016;18(3):471-9.
- 978.Desai, N. K., S. Harney, et al. Comparison of Controlled Attenuation Parameter and Liver Biopsy to Assess Hepatic Steatosis in Pediatric Patients. *The Journal of pediatrics* 2016;173:160-164 e1.
- 979.Gambato, M., G. Crespo, et al. Simple prediction of long-term clinical outcomes in patients with mild hepatitis C recurrence after liver transplantation. *Transplant International* 2016;29(6):698-706.
- 980.Hannah, W. N., Jr. and S. A. Harrison. Nonalcoholic fatty liver disease and elastography: Incremental advances but work still to be done. *Hepatology* 2016;63(6):1762-4.
- 981.Hezode, C., S. Chevaliez, et al. Retreatment with sofosbuvir and simeprevir of patients with hepatitis C virus genotype 1 or 4 who previously failed a daclatasvir-containing regimen. *Hepatology* 2016;63(6):1809-16.
- 982.Iacob, S., C. Ester, et al. Real-life Perception and Practice Patterns of NAFLD/NASH in Romania: Results of a Survey Completed by 102 Board-certified Gastroenterologists. *Journal of Gastrointestinal & Liver Diseases* 2016;25(2):183-9.
- 983.Jansen, C., C. Bogs, et al. Algorithm to rule out clinically significant portal hypertension combining Shear-wave elastography of liver and spleen: a prospective multicentre study. *Gut* 2016;65(6):1057-8.
- 984.Jayakumar, S., S. A. Harrison, et al. Noninvasive Markers of Fibrosis and Inflammation in Nonalcoholic Fatty Liver Disease. *Curr Hepatol Rep* 2016;15(2):86-95.
- 985.Lee, C. M., W. K. Jeong, et al. Diagnosis of Clinically Significant Portal Hypertension in Patients with Cirrhosis: Splenic Arterial Resistive Index versus Liver Stiffness Measurement. *Ultrasound in Medicine and Biology* 2016;42(6):1312-20.
- 986.Geng, X. X., R. G. Huang, et al. Transient elastography in clinical detection of liver cirrhosis: A systematic review and meta-analysis. *Saudi Journal of Gastroenterology* 2016;22(4):294-303.
- 987.Chang, H. K., E. Y. Chang, et al. Cyclooxygenase-2 Inhibitor Reduces Hepatic Stiffness in Pediatric Chronic Liver Disease Patients Following Kasai Portointerostomy. *Yonsei Medical Journal* 2016;57(4):893-9.
- 988.Chon, Y. E., K. J. Kim, et al. The Relationship between Type 2 Diabetes Mellitus and Non-Alcoholic Fatty Liver Disease Measured by Controlled Attenuation Parameter. *Yonsei Medical Journal* 2016;57(4):885-92.
- 989.Friedrich-Rust, M., T. Poynard, et al. Critical comparison of elastography methods to assess chronic liver disease. *Nature reviews. Gastroenterology & hepatology* 2016;13(7):402-11.
- 990.Girleanu, I., A. Trifan, et al. Platelet Indices and Liver Fibrosis Evaluation in Chronic Hepatitis C. *Revista medico-chirurgicala a Societatii de Medici si Naturalisti din Iasi* 2016;120(1):55-61.
- 991.Kotsiri, I., E. Hadziyannis, et al. Changes in serum transforming growth factor-beta1 levels in chronic hepatitis C patients under antiviral therapy. *Annals of gastroenterology* 2016;29(1):79-84.
- 992.Boyd, A. and K. Lacombe. More Long-term Assessment of Transient Elastography Is Needed for HIV/Hepatitis B Virus-Coinfected Patients Undergoing Treatment With Tenofovir. *Clinical Infectious Diseases* 2016;62(1):128-30.
- 993.Casado, J. L., A. Mena, et al. Liver toxicity and risk of discontinuation in HIV/hepatitis C virus-coinfected patients receiving an etravirine-containing antiretroviral regimen: influence of liver fibrosis. *HIV Medicine* 2016;17(1):62-7.
- 994.El-Asrar, M. A., N. S. Elbarbary, et al. Serum YKL-40 in young patients with beta-thalassemia major: Relation to hepatitis C virus infection, liver stiffness by transient elastography and cardiovascular complications. *Blood Cells, Molecules & Diseases* 2016;56(1):1-8.
- 995.Gao, Q., D. Shao, et al. Noninvasive assessment of liver stiffness by transient elastography (FibroScan) in liver fluke disease. *European Journal of Gastroenterology & Hepatology* 2016;28(1):70-3.
- 996.Guibal, A., G. Renosi, et al. Shear wave elastography: An accurate technique to stage liver fibrosis in

- chronic liver diseases. *Diagnostic and Interventional Imaging* 2016;97(1):91-9.
997. Halasz, T., G. Horvath, et al. Evaluation of Histological and non-Invasive Methods for the Detection of Liver Fibrosis: The Values of Histological and Digital Morphometric Analysis, Liver Stiffness Measurement and APRI Score. *Pathology oncology research* 2016;22(1):1-6.
998. Invernizzi, F., M. Vigano, et al. The prognosis and management of inactive HBV carriers. *Liver International* 2016;36 Suppl 1:100-4.
999. Koehler, E. M., E. P. Plomp, et al. Presence of diabetes mellitus and steatosis is associated with liver stiffness in a general population: The Rotterdam study. *Hepatology* 2016;63(1):138-47.
1000. Lavaill, L., A. Dussaucy, et al. Automatic detection of cirrhosis in hospitalized patients: a pragmatic experience. *European Journal of Gastroenterology & Hepatology* 2016;28(1):74-81.
1001. Chen, Y., W. K. Seto, et al. Relation of Tricuspid Regurgitation to Liver Stiffness Measured by Transient Elastography in Patients With Left-Sided Cardiac Valve Disease. *American Journal of Cardiology* 2016;117(4):640-6.
1002. Chang, P. E., G. B. Goh, et al. Clinical applications, limitations and future role of transient elastography in the management of liver disease. *World Journal of Gastrointestinal Pharmacology and Therapeutics* 2016;7(1):91-106.
1003. Franckenberg, S., C. Gubler, et al. [Radiology Update Ultrasound Elastography - Quintessence for the Primary Care Physician]. *Praxis (Bern 1994)* 2016;105(3):139-45.
1004. Crossan, C., E. A. Tsochatzis, et al. Cost-effectiveness of noninvasive liver fibrosis tests for treatment decisions in patients with chronic hepatitis B in the UK: systematic review and economic evaluation. *Journal of viral hepatitis* 2016;23(2):139-49.
1005. Ding, N. S., T. Nguyen, et al. Liver stiffness plus platelet count can be used to exclude high-risk oesophageal varices. *Liver International* 2016;36(2):240-5.
1006. Jacomet, C., A. Guyot-Lenat, et al. Addressing the challenges of chronic viral infections and addiction in prisons: the PRODEPIST study. *European Journal of Public Health* 2016;26(1):122-8.
1007. Kim, M. N., K. H. Han, et al. Reply to "Patients with chronic hepatitis b should be screened for hepatocellular carcinoma regardless of liver stiffness measurement.". *Hepatology* 2016;63(2):672-3.
1008. Kim, J. M., S. Y. Ha, et al. Predicting Hepatic Steatosis in Living Liver Donors via Noninvasive Methods. *Medicine (Baltimore)* 2016;95(7):e2718.
1009. Hashemi, S. A., S. M. Alavian, et al. Assessment of transient elastography (FibroScan) for diagnosis of fibrosis in non-alcoholic fatty liver disease: A systematic review and meta-analysis. *Caspian J Intern Med* 2016;7(4):242-252.
1010. Bellan, M., G. Pogliani, et al. Gas6 as a putative noninvasive biomarker of hepatic fibrosis. *Biomark Med* 2016;10(12):1241-1249.
1011. Charatcharoenwithaya, P., P. Phisalprapa, et al. Alanine aminotransferase course, serum hepatitis B virus DNA, and liver stiffness measurement for therapeutic decisions in hepatitis B e antigen-negative chronic hepatitis B. *Hepatology Research* 2016;46(13):1347-1357.
1012. Ferraioli, G., C. Tinelli, et al. The clinical value of controlled attenuation parameter for the noninvasive assessment of liver steatosis. *Liver International* 2016;36(12):1860-1866.
1013. Gerber, L. and M. Friedrich-Rust. Re: Non-Invasive Evaluation of Liver Fibrosis: 2-D Shear Wave Elastography, Transient Elastography or Acoustic Radiation Force Impulse Imaging? *Ultrasound in Medicine and Biology* 2016;42(12):3053.
1014. Hannah, W. N., Jr. and S. A. Harrison. Noninvasive imaging methods to determine severity of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. *Hepatology* 2016;64(6):2234-2243.
1015. Harrison, S. A., S. R. Marri, et al. Randomised clinical study: GR-MD-02, a galectin-3 inhibitor, vs. placebo in patients having non-alcoholic steatohepatitis with advanced fibrosis. *Alimentary Pharmacology & Therapeutics* 2016;44(11-12):1183-1198.
1016. Kalsch, J., H. Keskin, et al. Patients with ultrasound diagnosis of hepatic steatosis are at high metabolic risk. *Zeitschrift für Gastroenterologie* 2016;54(12):1312-1319.
1017. Katwesigye, E., E. Seremba, et al. Low seroprevalence of hepatitis delta antibodies in HIV/hepatitis B co-infected patients attending an urban HIV clinic in Uganda. *Afr Health Sci* 2016;16(4):1089-1093.
1018. Knop, V., D. Hoppe, et al. Regression of fibrosis and portal hypertension in HCV-associated cirrhosis and sustained virologic response after interferon-free antiviral therapy. *Journal of viral hepatitis* 2016;23(12):994-1002.
1019. Dai, T., J. Si, et al. Transient Elastography with Serum Hepatitis B Surface Antigen Enhances Liver Fibrosis Detection. *Medical science monitor : international medical journal of experimental and clinical research* 2016;22:2878-85.
1020. Collazos, J., E. Valle-Garay, et al. Factors associated with long-term CD4 cell recovery in HIV-infected patients on successful antiretroviral therapy. *HIV Medicine* 2016;17(7):532-41.
1021. Eddowes, P., M. Sasso, et al. Steatosis and liver stiffness measurements using transient elastography. *Hepatology* 2016;64(2):700.

1022. Ergelen, R., Y. Yilmaz, et al. Comparison of Doppler ultrasound and transient elastography in the diagnosis of significant fibrosis in patients with nonalcoholic steatohepatitis. *Abdominal Radiology* 2016;41(8):1505-10.
1023. Fazel, Y., A. B. Koenig, et al. Epidemiology and natural history of non-alcoholic fatty liver disease. *Metabolism: Clinical and Experimental* 2016;65(8):1017-25.
1024. Fraquelli, M., A. Baccarin, et al. Liver stiffness measurement reliability and main determinants of point shear-wave elastography in patients with chronic liver disease. *Alimentary Pharmacology & Therapeutics* 2016;44(4):356-65.
1025. Karlas, T. and D. Petroff. Is Magnetic Resonance Imaging Really More Accurate for Classifying Steatosis Than Controlled Attenuation Parameter? *Gastroenterology* 2016;151(2):374-5.
1026. Kimer, N., S. Wiese, et al. Advances in the treatment of portal hypertension in cirrhosis. *Expert Review of Gastroenterology & Hepatology* 2016;10(8):961-9.
1027. Kwok, R., K. C. Choi, et al. Screening diabetic patients for non-alcoholic fatty liver disease with controlled attenuation parameter and liver stiffness measurements: a prospective cohort study. *Gut* 2016;65(8):1359-68.
1028. Behairy Bel, S., M. M. Sira, et al. Transient elastography compared to liver biopsy and morphometry for predicting fibrosis in pediatric chronic liver disease: Does etiology matter? *World Journal of Gastroenterology* 2016;22(16):4238-49.
1029. de Ledinghen, V., G. L. Wong, et al. Controlled attenuation parameter for the diagnosis of steatosis in non-alcoholic fatty liver disease. *Journal of Gastroenterology & Hepatology* 2016;31(4):848-55.
1030. Heo, J. Y., S. U. Kim, et al. Use of Wisteria Floribunda Agglutinin-Positive Human Mac-2 Binding Protein in Assessing Risk of Hepatocellular Carcinoma Due to Hepatitis B Virus. *Medicine (Baltimore)* 2016;95(14):e3328.
1031. Buechter, M., A. Kahraman, et al. Spleen and Liver Stiffness Is Positively Correlated with the Risk of Esophageal Variceal Bleeding. *Digestion* 2016;94(3):138-144.
1032. Chazouilleres, O. Novel Aspects in the Management of Cholestatic Liver Diseases. *Digestive Diseases* 2016;34(4):340-6.
1033. Chekuri, S., J. Nickerson, et al. Liver Stiffness Decreases Rapidly in Response to Successful Hepatitis C Treatment and Then Plateaus. *PLoS ONE* 2016;11(7):e0159413.
1034. Chen, B., R. A. Schreiber, et al. Assessment of Liver Stiffness in Pediatric Fontan Patients Using Transient Elastography. *Canadian Journal of Gastroenterology & Hepatology* 2016;2016:7125193.
1035. Colli, A., M. Fraquelli, et al. Deciding on Interferon-Free Treatment for Chronic Hepatitis C: Updating Liver Stiffness Cut-Off Values to Maximize Benefit. *PLoS ONE* 2016;11(10):e0164452.
1036. D'Ambrosio, R., E. Degasperi, et al. Serological Tests Do Not Predict Residual Fibrosis in Hepatitis C Cirrhotics with a Sustained Virological Response to Interferon. *PLoS ONE* 2016;11(6):e0155967.
1037. Doyle, M. A., J. Singer, et al. Improving treatment and liver fibrosis outcomes with metformin in HCV-HIV co-infected and HCV mono-infected patients with insulin resistance: study protocol for a randomized controlled trial. *Trials* 2016;17(1):331.
1038. Ehlik, H., R. Wroblewski, et al. Validation of Transient Elastography and Comparison with Spleen Length Measurement for Staging of Fibrosis and Clinical Prognosis in Primary Sclerosing Cholangitis. *PLoS ONE* 2016;11(10):e0164224.
1039. Ginès, P., I. Graupera, et al. Screening for liver fibrosis in the general population: a call for action. *The Lancet Gastroenterology & Hepatology* 2016;1(3):256-260.
1040. Hametner, S., A. Ferlitsch, et al. The VITRO Score (Von Willebrand Factor Antigen/Thrombocyte Ratio) as a New Marker for Clinically Significant Portal Hypertension in Comparison to Other Non-Invasive Parameters of Fibrosis Including ELF Test. *PLoS ONE* 2016;11(2):e0149230.
1041. Jaquet, A., G. Wandeler, et al. HIV infection, viral hepatitis and liver fibrosis among prison inmates in West Africa. *BMC Infectious Diseases* 2016;16(1):249.
1042. Jeon, S. R. and J. O. Kim. Capsule Endoscopy for Portal Hypertensive Enteropathy. *Gastroenterology Research & Practice* 2016;2016:8501394.
1043. Kubo, K., H. Kawakami, et al. Liver elasticity measurement before and after biliary drainage in patients with obstructive jaundice: a prospective cohort study. *BMC Gastroenterology* 2016;16(1):65.
1044. Kuo, Y. H., M. C. Tsai, et al. Associated Factors for Metabolic Syndrome in the Older Adults with Chronic Virus Hepatitis in the Community. *PLoS ONE* 2016;11(5):e0155544.
1045. Ahn, J. M., Y. H. Paik, et al. Relationship between Controlled Attenuation Parameter and Hepatic Steatosis as Assessed by Ultrasound in Alcoholic or Nonalcoholic Fatty Liver Disease. *Gut Liver* 2016;10(2):295-302.
1046. Arslanow, A., C. S. Stokes, et al. The common PNPLA3 variant p.I148M is associated with liver fat contents as quantified by controlled attenuation parameter (CAP). *Liver International* 2016;36(3):418-26.

1047. Audsley, J., C. Robson, et al. Liver Fibrosis Regression Measured by Transient Elastography in Human Immunodeficiency Virus (HIV)-Hepatitis B Virus (HBV)-Coinfected Individuals on Long-Term HBV-Active Combination Antiretroviral Therapy. *Open Forum Infectious Diseases* 2016;3(1):ofw035.
1048. Alkhouri, N. and A. E. Feldstein. Noninvasive diagnosis of nonalcoholic fatty liver disease: Are we there yet? *Metabolism: Clinical and Experimental* 2016.
1049. Bansal, R. K., M. Kumar, et al. Prospective study of profile of hepatic osteodystrophy in patients with non-cholestatic liver cirrhosis and impact of bisphosphonate supplementation. *United European Gastroenterology Journal* 2016;4(1):77-83.
1050. Abraldes, J. G., C. Bureau, et al. Noninvasive tools and risk of clinically significant portal hypertension and varices in compensated cirrhosis: The "Anticipate" study. *Hepatology* 2016;64(6):2173-2184.
1051. Adler, M., L. Larocca, et al. Evaluating the risk of hepatocellular carcinoma in patients with prominently elevated liver stiffness measurements by FibroScan: a multicentre study. *HPB : the official journal of the International Hepato Pancreato Biliary Association* 2016;18(8):678-83.
1052. Arain, A., J. De Sousa, et al. Pilot Study: Combining Formal and Peer Education with FibroScan to Increase HCV Screening and Treatment in Persons who use Drugs. *Journal of Substance Abuse Treatment* 2016;67:44-9.
1053. Attia, D., H. Bantel, et al. Liver stiffness measurement using acoustic radiation force impulse elastography in overweight and obese patients. *Alimentary Pharmacology & Therapeutics* 2016;44(4):366-79.
1054. Bano, A., L. Chaker, et al. Thyroid Function and the Risk of Nonalcoholic Fatty Liver Disease: The Rotterdam Study. *The Journal of Clinical Endocrinology and Metabolism* 2016;101(8):3204-11.
1055. Andreev, D. N., I. V. Maev, et al. [The efficiency and safety of anti-Helicobacter pylori therapy in patients with concomitant chronic hepatitis C]. *Terapevticheskii Arkhiv* 2016;88(4):75-81.
1056. Arslanow, A., M. Teutsch, et al. Short-Term Hypocaloric High-Fiber and High-Protein Diet Improves Hepatic Steatosis Assessed by Controlled Attenuation Parameter. *Clinical and Translational Gastroenterology* 2016;7(6):e176.
1057. Bakulin, I. G., Y. G. Sandler, et al. [A new non-invasive method for assessing steatosis in chronic liver diseases]. *Terapevticheskii Arkhiv* 2016;88(2):49-57.
1058. Wu, Q. W., X. Zhu, et al. [Expression of Tim-3 on Peripheral CD56(+) NK Cells and Its Correlation with Liver Fibrosis in Patients with Advanced Schistosomiasis]. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* 2015;33(5):346-50.
1059. Xu, X., Y. Su, et al. Performance of transient elastography assessing fibrosis of single hepatitis B virus infection: a systematic review and meta-analysis of a diagnostic test. *Hepatology International* 2015;9(4):558-66.
1060. You, S. C., K. J. Kim, et al. Hepatic fibrosis assessed using transient elastography independently associated with coronary artery calcification. *Journal of Gastroenterology & Hepatology* 2015;30(10):1536-42.
1061. Yada, N., T. Sakurai, et al. A Newly Developed Shear Wave Elastography Modality: With a Unique Reliability Index. *Oncology* 2015;89 Suppl 2:53-9.
1062. Zhang, E., C. Wartelle-Bladou, et al. Cost-utility analysis of nonalcoholic steatohepatitis screening. *European Radiology* 2015;25(11):3282-94.
1063. Zeng, X., C. Xu, et al. The Diagnostic Value of FibroScan in Assessing Significant Liver Fibrosis in Patients with Chronic Hepatitis B. *The West Indian medical journal* 2015;65(1):106-110.
1064. Yo, I. K., O. S. Kwon, et al. The factors associated with longitudinal changes in liver stiffness in patients with chronic hepatitis B. *Clinical and Molecular Hepatology* 2015;21(1):32-40.
1065. Zhang, W., L. Wang, et al. Liver Stiffness Measurement, Better than APRI, Fibroindex, Fib-4, and NBI Gastroscopy, Predicts Portal Hypertension in Patients with Cirrhosis. *Cell biochemistry and biophysics* 2015;71(2):865-73.
1066. Zeng, X., C. Xu, et al. Influence of Hepatic Inflammation on FibroScan Findings in Diagnosing Fibrosis in Patients with Chronic Hepatitis B. *Ultrasound in Medicine and Biology* 2015;41(6):1538-44.
1067. Zeng, X., C. Xu, et al. Performance of several simple, noninvasive models for assessing significant liver fibrosis in patients with chronic hepatitis B. *Croatian medical journal* 2015;56(3):272-9.
1068. You, S. C., K. J. Kim, et al. Factors associated with significant liver fibrosis assessed using transient elastography in general population. *World Journal of Gastroenterology* 2015;21(4):1158-66.
1069. Yoneda, M., K. Imajo, et al. Screening of nonalcoholic fatty liver disease in patients with coronary heart disease by transient elastography: proceed with caution. *Wiener Klinische Wochenschrift* 2015;127(1-2):75-6.
1070. Yoshioka, K., S. Hashimoto, et al. Measurement of liver stiffness as a non-invasive method for diagnosis of non-alcoholic fatty liver disease. *Hepatology Research* 2015;45(2):142-51.
1071. Zhang, D., M. Chen, et al. Comparison of acoustic radiation force impulse imaging and

2015

- transient elastography for non-invasive assessment of liver fibrosis in patients with chronic hepatitis B. *Ultrasound in Medicine and Biology* 2015;41(1):7-14.
1072. Yoneda, M., E. Thomas, et al. Supersonic Shear Imaging and Transient Elastography With the XL Probe Accurately Detect Fibrosis in Overweight or Obese Patients With Chronic Liver Disease. *Clinical Gastroenterology & Hepatology* 2015;13(8):1502-1509 e5.
1073. Zaleska-Dorobisz, U., A. Pawlus, et al. [SWE elastography in assessment of liver fibrosis]. *Postepy higieny i medycyny doswiadczałnej* 2015;69:221-6.
1074. Zhang, Y. P., Q. Zhao, et al. Relationships between transient elastography values and liver fibrosis in chronic liver disease patients with normal or mildly abnormal aminotransferase levels. *Genetics and molecular research* 2015;14(4):18172-80.
1075. Zykus, R., L. Jonaitis, et al. Liver and spleen transient elastography predicts portal hypertension in patients with chronic liver disease: a prospective cohort study. *BMC Gastroenterology* 2015;15(1):183.
1076. Stockdale, A. J., R. O. Phillips, et al. Liver Fibrosis by Transient Elastography and Virologic Outcomes After Introduction of Tenofovir in Lamivudine-Experienced Adults With HIV and Hepatitis B Virus Coinfection in Ghana. *Clinical Infectious Diseases* 2015;61(6):883-91.
1077. Tapper, E. B., N. Sengupta, et al. Cost-Effective Evaluation of Nonalcoholic Fatty Liver Disease With NAFLD Fibrosis Score and Vibration Controlled Transient Elastography. *American Journal of Gastroenterology* 2015;110(9):1298-304.
1078. Tsouchatzis, E. A. and L. Castera. Assessing liver disease in HIV-HCV coinfected patients. *Current Opinion in HIV and AIDS* 2015;10(5):316-22.
1079. Webb, M., O. Shibolet, et al. Assessment of Liver and Spleen Stiffness in Patients With Myelofibrosis Using FibroScan and Shear Wave Elastography. *Ultrasound Quarterly* 2015;31(3):166-9.
1080. Wei, J. L., J. C. Leung, et al. Prevalence and Severity of Nonalcoholic Fatty Liver Disease in Non-Obese Patients: A Population Study Using Proton-Magnetic Resonance Spectroscopy. *American Journal of Gastroenterology* 2015;110(9):1306-14.
1081. Wong, V. W., G. L. Wong, et al. Bacterial endotoxin and non-alcoholic fatty liver disease in the general population: a prospective cohort study. *Alimentary Pharmacology & Therapeutics* 2015;42(6):731-40.
1082. Wong, V. W. and H. L. Janssen. Can we use HCC risk scores to individualize surveillance in chronic hepatitis B infection? *Journal of Hepatology* 2015;63(3):722-32.
1083. Wong, G. L., W. Z. Espinosa, et al. Personalized management of cirrhosis by non-invasive tests of liver fibrosis. *Clinical and Molecular Hepatology* 2015;21(3):200-11.
1084. Wang, C. Y., X. J. Cheng, et al. [Diagnostic value of FibroTest for liver fibrosis in patients with chronic hepatitis B]. *Zhonghua Gan Zang Bing Za Zhi* 2015;23(10):738-41.
1085. Thiele, M., S. Detlefsen, et al. Transient and 2-dimensional Shear-Wave Elastography provide comparable assessment of Alcoholic Liver Fibrosis and Cirrhosis. *Gastroenterology* 2015.
1086. Williams, K. H., K. Burns, et al. An association of large-fibre peripheral nerve dysfunction with non-invasive measures of liver fibrosis secondary to non-alcoholic fatty liver disease in diabetes. *Journal of diabetes and its complications* 2015;29(8):1240-7.
1087. Trovato, F. M., J. M. Tognarelli, et al. Challenges of liver cancer: Future emerging tools in imaging and urinary biomarkers. *World Journal of Gastroenterology* 2015;7(26):2664-75.
1088. Williams, K. H., A. J. Vieira De Ribeiro, et al. Circulating dipeptidyl peptidase-4 activity correlates with measures of hepatocyte apoptosis and fibrosis in non-alcoholic fatty liver disease in type 2 diabetes mellitus and obesity: A dual cohort cross-sectional study. *Journal of diabetes* 2015;7(6):809-819.
1089. Tapper, E. B. and N. H. Afdhal. Vibration-controlled transient elastography: a practical approach to the noninvasive assessment of liver fibrosis. *Current Opinion in Gastroenterology* 2015;31(3):192-8.
1090. Tatsumi, A., S. Maekawa, et al. Liver stiffness measurement for risk assessment of hepatocellular carcinoma. *Hepatology Research* 2015;45(5):523-32.
1091. Wong, V. W., P. Lampertico, et al. Probability-based interpretation of liver stiffness measurement in untreated chronic hepatitis B patients. *Digestive Diseases & Sciences* 2015;60(5):1448-56.
1092. Wong, G. L., H. L. Chan, et al. Noninvasive assessments of liver fibrosis with transient elastography and Hui index predict survival in patients with chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2015;30(3):582-90.
1093. Trovato, F. M., S. Atzori, et al. Liver and spleen transient elastography and Acoustic Radiation Force Impulse Measurements. Performance and comparison of measurements in the same area concurrently assessed for liver fibrosis by biopsy. *Advances in medical sciences* 2015;60(2):300-306.
1094. Sulyok, M., M. Makara, et al. Hepatic steatosis in individuals living with HIV measured by controlled attenuation parameter: a cross-sectional study. *European Journal of Gastroenterology & Hepatology* 2015;27(6):679-85.

1095. Williams, K. H., A. J. Viera de Ribeiro, et al. Lower serum fibroblast activation protein shows promise in the exclusion of clinically significant liver fibrosis due to non-alcoholic fatty liver disease in diabetes and obesity. *Diabetes Research and Clinical Practice* 2015;108(3):466-72.
1096. Wong, G. L., H. L. Chan, et al. Impact of IL28B and PNPLA3 polymorphisms on treatment outcomes in patients infected with genotype 6 hepatitis C virus. *Journal of Gastroenterology & Hepatology* 2015;30(6):1040-8.
1097. Tapper, E. B., L. Castera, et al. FibroScan (Vibration-Controlled Transient Elastography): Where Does It Stand in the United States Practice. *Clinical Gastroenterology & Hepatology* 2015;13(1):27-36.
1098. Wong, V. W., G. L. Wong, et al. Incidence of non-alcoholic fatty liver disease in Hong Kong: A population study with paired proton-magnetic resonance spectroscopy. *Journal of Hepatology* 2015;62(1):182-9.
1099. Wong, G. L. and V. W. Wong. Fat and fiber: how the controlled attenuation parameter complements noninvasive assessment of liver fibrosis. *Digestive Diseases & Sciences* 2015;60(1):9-12.
1100. Stefanescu, H., C. Radu, et al. Non-invasive menage a trois for the prediction of high-risk varices: stepwise algorithm using lok score, liver and spleen stiffness. *Liver International* 2015;35(2):317-25.
1101. Walker, C., R. Thimme, et al. [Fibroscan of the liver]. *Deutsche Medizinische Wochenschrift* 2015;140(4):267-70.
1102. Suarez Ferrer, C., E. Llop Herrera, et al. Idiopathic portal hypertension regarding thiopurine treatment in patients with inflammatory bowel disease. *Revista Española de Enfermedades Digestivas* 2015;108.
1103. Van Gossum, A., L. Pironi, et al. Transient Elastography (FibroScan) Is Not Correlated With Liver Fibrosis but With Cholestasis in Patients With Long-Term Home Parenteral Nutrition. *Journal of Parenteral and Enteral Nutrition* 2015;39(6):719-24.
1104. Tovo, C. V., A. Z. de Mattos, et al. Noninvasive imaging assessment of non-alcoholic fatty liver disease: Focus on liver scintigraphy. *World Journal of Gastroenterology* 2015;21(15):4432-9.
1105. Vuppulanchi, R. and A. J. Sanyal. Myths and mysteries about staging hepatic fibrosis by fibroscan. *Clinical Gastroenterology & Hepatology* 2015;13(4):780-2.
1106. Wong, G. L., R. Kwok, et al. Huge adrenal hemangioma: a rare cause of deceptively high liver stiffness measurement by transient elastography. *Clinical Gastroenterology & Hepatology* 2015;13(4):e37-8.
1107. Sobhonslidsuk, A., A. Pulsombat, et al. Non-alcoholic Fatty Liver Disease (NAFLD) and Significant Hepatic Fibrosis Defined by Non-invasive Assessment in Patients with Type 2 Diabetes. *Asian Pacific Journal of Cancer Prevention* 2015;16(5):1789-94.
1108. Thavorn, K. and D. Coyle. Transient Elastography and Controlled Attenuation Parameter for Diagnosing Liver Fibrosis and Steatosis in Ontario: An Economic Analysis. *Ontario Health Technology Assessment Series* 2015;15(19):1-58.
1109. Wang, Y., Q. Fan, et al. Controlled attenuation parameter for assessment of hepatic steatosis grades: a diagnostic meta-analysis. *International journal of clinical and experimental medicine* 2015;8(10):17654-63.
1110. Sirli, R., I. Sporea, et al. Ultrasound-based elastography for the diagnosis of portal hypertension in cirrhotics. *World Journal of Gastroenterology* 2015;21(41):11542-51.
1111. Seo, Y. S., M. Y. Kim, et al. Accuracy of transient elastography in assessing liver fibrosis in chronic viral hepatitis: A multicentre, retrospective study. *Liver International* 2015;35(10):2246-55.
1112. Serres, L., P. Vasseur, et al. Cardiovascular events in chronic hepatitis C: prognostic value of liver stiffness evolution. *European Journal of Gastroenterology & Hepatology* 2015;27(11):1286-92.
1113. Shen, F., R. D. Zheng, et al. Impact of skin capsular distance on the performance of controlled attenuation parameter in patients with chronic liver disease. *Liver International* 2015;35(11):2392-400.
1114. Sebastiani, G., P. Ghali, et al. Patient preference and willingness to pay for transient elastography versus liver biopsy: A perspective from British Columbia. *Canadian Journal of Gastroenterology & Hepatology* 2015;29(4):219-20.
1115. Shin, S. H., S. U. Kim, et al. Liver stiffness-based model for prediction of hepatocellular carcinoma in chronic hepatitis B virus infection: comparison with histological fibrosis. *Liver International* 2015;35(3):1054-62.
1116. Shen, Q. L., Y. J. Chen, et al. Assessment of liver fibrosis by Fibroscan as compared to liver biopsy in biliary atresia. *World Journal of Gastroenterology* 2015;21(22):6931-6.
1117. Shah, S., Y. Ma, et al. Association of HIV, hepatitis C virus and liver fibrosis severity with interleukin-6 and C-reactive protein levels. *Aids* 2015;29(11):1325-33.
1118. Sekimoto, T., H. Maruyama, et al. Liver Stiffness: A Significant Relationship with the Waveform Pattern in the Hepatic Vein. *Ultrasound in Medicine and Biology* 2015;41(7):1801-7.
1119. Shen, J., G. L. Wong, et al. PNPLA3 gene polymorphism and response to lifestyle modification in patients with nonalcoholic fatty liver disease. *Journal of Gastroenterology & Hepatology* 2015;30(1):139-46.

1120. Schwabl, P., S. Bota, et al. New reliability criteria for transient elastography increase the number of accurate measurements for screening of cirrhosis and portal hypertension. *Liver International* 2015;35(2):381-90.
1121. Sharma, P., R. Bansal, et al. Role of Transient Elastography (Fibroscan) in Differentiating Severe Acute Hepatitis and Acute on Chronic Liver Failure. *Journal of Clinical and Experimental Hepatology* 2015;5(4):303-9.
1122. Seifert, L. L., E. Vorona, et al. Interferon-Free Sofosbuvir-Based Anti-HCV Therapy After Liver Transplantation. *Annals of Transplantation* 2015;20:561-8.
1123. Scheiner, B., M. Mandorfer, et al. The Impact of PNPLA3 rs738409 SNP on Liver Fibrosis Progression, Portal Hypertension and Hepatic Steatosis in HIV/HCV Coinfection. *PLoS ONE* 2015;10(11):e0143429.
1124. Richards, D. B., L. M. Cookson, et al. Therapeutic Clearance of Amyloid by Antibodies to Serum Amyloid P Component. *The New England Journal of Medicine* 2015;373(12):1106-14.
1125. Sagir, A., B. Glaubach, et al. Transient Elastography for the Detection of Liver Damage in Patients with HIV. *Infectious Diseases and Therapy* 2015;4(3):355-64.
1126. Sagnelli, C., S. Martini, et al. Liver fibrosis in human immunodeficiency virus/hepatitis C virus coinfection: Diagnostic methods and clinical impact. *World Journal of Hepatology* 2015;7(24):2510-21.
1127. Sarsekeyeva, N., B. Kosherova, et al. Liver Fibrosis in Combined Course of Chronic Hepatitis C and Hiv Infection. *Georgian Medical News* 2015;--(248):30-4.
1128. Sacchi, P., S. Cima, et al. Liver fibrosis, microbial translocation and immune activation markers in HIV and HCV infections and in HIV/HCV co-infection. *Digestive & Liver Disease* 2015;47(3):218-25.
1129. Ribaldone, D. G., M. Garavagno, et al. Prevalence and prognostic value of hepatic histological alterations in patients with Crohn's disease. *Scandinavian Journal of Gastroenterology* 2015;50(12):1463-8.
1130. Sadler, M. D., P. Crotty, et al. Noninvasive methods, including transient elastography, for the detection of liver disease in adults with cystic fibrosis. *Canadian Journal of Gastroenterology & Hepatology* 2015;29(3):139-44.
1131. Perazzo, H., F. F. Fernandes, et al. Points to be considered when using transient elastography for diagnosis of portal hypertension according to the Baveno's VI consensus. *Journal of Hepatology* 2015;63(4):1048-9.
1132. Petta, S., M. Maida, et al. The severity of steatosis influences liver stiffness measurement in patients with nonalcoholic fatty liver disease. *Hepatology* 2015;62(4):1101-10.
1133. Pawlus, A., D. Sokolowska-Dabek, et al. Ultrasound Elastography--Review of Techniques and Its Clinical Applications in Pediatrics--Part 1. *Advances in Clinical and Experimental Medicine* 2015;24(3):537-43.
1134. Par, A., A. Vincze, et al. [Non-invasive diagnostic methods of fibrosis in chronic hepatitis C virus infection: their role in treatment indication, follow-up and assessment of prognosis]. *Orvosi Hetilap* 2015;156(21):855-61.
1135. Parikh, P., J. Patel, et al. Serum ferritin levels predict histological severity in patients with nonalcoholic fatty liver disease in India. *Indian Journal of Gastroenterology* 2015;34(3):200-8.
1136. Perazzo, H., F. F. Fernandes, et al. Interobserver variability in transient elastography analysis of patients with chronic hepatitis C. *Liver International* 2015;35(5):1533-9.
1137. Petta, S., E. Vanni, et al. The combination of liver stiffness measurement and NAFLD fibrosis score improves the noninvasive diagnostic accuracy for severe liver fibrosis in patients with nonalcoholic fatty liver disease. *Liver International* 2015;35(5):1566-73.
1138. Procopet, B., A. Berzigotti, et al. Real-time shear-wave elastography: applicability, reliability and accuracy for clinically significant portal hypertension. *Journal of Hepatology* 2015;62(5):1068-75.
1139. Procopet, B., V. M. Cristea, et al. Serum tests, liver stiffness and artificial neural networks for diagnosing cirrhosis and portal hypertension. *Digestive & Liver Disease* 2015;47(5):411-6.
1140. Pillon, M., N. S. Carucci, et al. Focal nodular hyperplasia of the liver: an emerging complication of hematopoietic SCT in children. *Bone Marrow Transplantation* 2015;50(3):414-9.
1141. Park, M. S., S. U. Kim, et al. Prognostic value of the combined use of transient elastography and fibrotest in patients with chronic hepatitis B. *Liver International* 2015;35(2):455-62.
1142. Plomp, E. P., S. Darwish Murad, et al. Prothrombotic genetic risk factors are associated with an increased risk of liver fibrosis in the general population: The Rotterdam Study. *Journal of Hepatology* 2015;63(6):1459-65.
1143. Pathik, P., S. Ravindra, et al. Fibroscan versus simple noninvasive screening tools in predicting fibrosis in high-risk nonalcoholic fatty liver disease patients from Western India. *Annals of Gastroenterology* 2015;28(2):281-286.
1144. Paparo, F., G. Cenderello, et al. Diagnostic value of MRI proton density fat fraction for assessing liver steatosis in chronic viral C hepatitis. *BioMed Research International* 2015;2015:758164.

1145. Pavlov, C. S., G. Casazza, et al. Transient elastography for diagnosis of stages of hepatic fibrosis and cirrhosis in people with alcoholic liver disease. Cochrane database of systematic reviews 2015;1:CD010542.
1146. Perazzo, H., V. G. Veloso, et al. Factors That Could Impact on Liver Fibrosis Staging by Transient Elastography. International journal of hepatology 2015;2015:624596.
1147. Poupon, R. Non-Invasive Assessment of Liver Fibrosis Progression and Prognosis in Primary Biliary Cholangitis. Digestive Diseases 2015;33 Suppl 2:115-7.
1148. Ozaras, R., G. Corti, et al. Differences in the availability of diagnostics and treatment modalities for chronic hepatitis B across Europe. Clinical microbiology and infection 2015;21(11):1027-32.
1149. Orlic, L., I. Mikolasevic, et al. Nonalcoholic fatty liver disease and the renin-angiotensin system blockers in the patients with chronic kidney disease. Wiener Klinische Wochenschrift 2015;127(9-10):355-62.
1150. Ozimok, C. J., A. Z. Kielar, et al. Efficacy and Accessibility of a Novel Liver Fibrosis Evaluation Program: Limited Abdominal Ultrasound with Concurrent Transient Elastography. Academic Radiology 2015;22(7):814-9.
1151. Ou, X. J., X. M. Wang, et al. [Comparison of FibroTouch and FibroScan for the assessment of fibrosis in chronic hepatitis B patients]. Zhonghua Gan Zang Bing Za Zhi 2015;23(2):103-6.
1152. Orasan, O. H., M. Iancu, et al. Non-invasive assessment of liver fibrosis in chronic viral hepatitis. European Journal of Clinical Investigation 2015;45(12):1243-51.
1153. Macias, J., A. Rivero-Juarez, et al. Impact of genetic polymorphisms associated with nonalcoholic fatty liver disease on HIV-infected individuals. Aids 2015;29(15):1927-35.
1154. Lutz, H. H., B. Schroeter, et al. Doppler Ultrasound and Transient Elastography in Liver Transplant Patients for Noninvasive Evaluation of Liver Fibrosis in Comparison with Histology: A Prospective Observational Study. Digestive Diseases & Sciences 2015;60(9):2825-31.
1155. Mandorfer, M., K. Kozbial, et al. Interferon-free regimens for chronic hepatitis C overcome the effects of portal hypertension on virological responses. Alimentary Pharmacology & Therapeutics 2015;42(6):707-18.
1156. Marshall, A. D., M. Micallef, et al. Liver disease knowledge and acceptability of non-invasive liver fibrosis assessment among people who inject drugs in the drug and alcohol setting: The LiveRLife Study. International Journal of Drug Policy 2015;26(10):984-91.
1157. Morse, C. G., M. McLaughlin, et al. Transient elastography for the detection of hepatic fibrosis in HIV-monoinfected adults with elevated aminotransferases on antiretroviral therapy. Aids 2015;29(17):2297-302.
1158. Macias, J., M. Mancebo, et al. Low risk of liver decompensation among human immunodeficiency virus/hepatitis C virus-coinfected patients with mild fibrosis in the short term. Hepatology 2015;61(5):1503-11.
1159. Lupsor-Platon, M., D. Feier, et al. Diagnostic Accuracy of Controlled Attenuation Parameter Measured by Transient Elastography for the Non-invasive Assessment of Liver Steatosis: a Prospective Study. Journal of Gastrointestinal & Liver Diseases 2015;24(1):35-42.
1160. Meng, F., Y. Zheng, et al. Noninvasive evaluation of liver fibrosis using real-time tissue elastography and transient elastography (FibroScan). Journal of ultrasound in medicine 2015;34(3):403-10.
1161. Ogawa, E., N. Furusyo, et al. Non-invasive fibrosis assessment predicts sustained virological response to telaprevir with pegylated interferon and ribavirin for chronic hepatitis C. Antiviral Therapy 2015;20(2):185-92.
1162. Mansoor, S., E. Collyer, et al. A comprehensive review of noninvasive liver fibrosis tests in pediatric nonalcoholic Fatty liver disease. Current Gastroenterology Reports 2015;17(6):23.
1163. Mikolasevic, I., L. Orlic, et al. Usefulness of liver test and controlled attenuation parameter in detection of nonalcoholic fatty liver disease in patients with chronic renal failure and coronary heart disease. Wiener Klinische Wochenschrift 2015;127(11-12):451-8.
1164. Mi, Y. Q., Q. Y. Shi, et al. Controlled attenuation parameter for noninvasive assessment of hepatic steatosis using fibroscan((R)): validation in chronic hepatitis B. Digestive Diseases & Sciences 2015;60(1):243-51.
1165. Okada, M., T. Murakami, et al. Comparison between T1 relaxation time of Gd-EOB-DTPA-enhanced MRI and liver stiffness measurement of ultrasound elastography in the evaluation of cirrhotic liver. Journal of Magnetic Resonance Imaging 2015;41(2):329-38.
1166. Nishi, H., K. Toda, et al. Novel Method of Evaluating Liver Stiffness Using Transient Elastography to Evaluate Perioperative Status in Severe Heart Failure. Circulation Journal 2015;79(2):391-397.
1167. Mohr, R., R. Schierwagen, et al. Liver Fibrosis in HIV Patients Receiving a Modern cART: Which Factors Play a Role? Medicine (Baltimore) 2015;94(50):e2127.
1168. Mueller, S., S. Englert, et al. Inflammation-adapted liver stiffness values for improved fibrosis staging in patients with hepatitis C virus and

- alcoholic liver disease. *Liver International* 2015;35(12):2514-21.
1169. Naqvi, A., V. Giordanengo, et al. Virological response and resistance mutations to NS3/4A inhibitors in hepatitis C virus-human immunodeficiency virus coinfection. *World Journal of Gastroenterology* 2015;7(18):2177-83.
1170. Mancia, C., V. Loustaud-Ratti, et al. Controlled Attenuation Parameter and Liver Stiffness Measurements for Steatosis Assessment in the Liver Transplant of Brain Dead Donors. *Transplantation* 2015;99(8):1619-24.
1171. Margariti, A., M. D. Kontogianni, et al. Increased abdominal fat levels measured by bioelectrical impedance are associated with histological lesions of nonalcoholic steatohepatitis. *European Journal of Gastroenterology & Hepatology* 2015;27(8):907-13.
1172. Merat, S., H. Poustchi, et al. PolyPill for Prevention of Cardiovascular Disease in an Urban Iranian Population with Special Focus on Nonalcoholic Steatohepatitis: A Pragmatic Randomized Controlled Trial within a Cohort (PolyIran - Liver) - Study Protocol. *Archives of Iranian medicine* 2015;18(8):515-23.
1173. Matthews, G. V., J. Neuhaus, et al. Baseline prevalence and predictors of liver fibrosis among HIV-positive individuals: a substudy of the INSIGHT Strategic Timing of AntiRetroviral Treatment (START) trial. *HIV Medicine* 2015;16 Suppl 1:129-36.
1174. Nakao, T., H. Ushigome, et al. Evaluation of renal allograft fibrosis by transient elastography (fibro scan). *Transplantation Proceedings* 2015;47(3):640-3.
1175. Nascimbeni, F., P. Lebray, et al. Significant Variations in Elastometry Measurements Made Within Short-term in Patients With Chronic Liver Diseases. *Clinical Gastroenterology & Hepatology* 2015;13(4):763-771 e6.
1176. Maggi, P., S. Altizio, et al. Prevalence and Risk Factors for Significant Liver Fibrosis in Patients with HIV Infection. *In Vivo* 2015;29(6):771-775.
1177. Lupsor-Platon, M. and R. Badea. Noninvasive assessment of alcoholic liver disease using unidimensional transient elastography (Fibroscan((R))). *World Journal of Gastroenterology* 2015;21(42):11914-23.
1178. Liu, Y., C. F. Dong, et al. Optimal linear combination of ARFI, transient elastography and APRI for the assessment of fibrosis in chronic hepatitis B. *Liver International* 2015;35(3):816-25.
1179. Li, C., J. Y. Zhang, et al. FibroScan predicts ascites after liver resection for hepatitis B virus-related hepatocellular carcinoma: A prospective cohort study. *International Journal of Surgery* 2015;20:21-25.
1180. Lim, T. H., E. Gane, et al. Serological and clinical outcomes of horizontally transmitted chronic hepatitis B infection in New Zealand Maori: results from a 28-year follow-up study. *Gut* 2015;64(6):966-72.
1181. Liu, K., K. T. Bui, et al. Longer duration of transient elastography predicts unreliable liver stiffness measurements. *European Journal of Gastroenterology & Hepatology* 2015;27(6):655-9.
1182. Li, W., C. Zhu, et al. Increased Pygo2 expression in liver of patients with hepatitis B virus-related fibrosis. *Liver International* 2015;35(12):2522-9.
1183. Lepida, A., M. Colombo, et al. Final Results of the Telaprevir Access Program: FibroScan Values Predict Safety and Efficacy in Hepatitis C Patients with Advanced Fibrosis or Cirrhosis. *PLoS ONE* 2015;10(9):e0138503.
1184. Liang, L. B., X. Zhu, et al. Serum hepatitis B surface antigen titer and transient elastography in screening for insignificant fibrosis in HBeAg-positive chronic hepatitis B patients. *Therapeutics and Clinical Risk Management* 2015;11:229-35.
1185. Lee, S. W., B. K. Kim, et al. Clinical Predictors of Silent but Substantial Liver Fibrosis in Primary Sjogren's Syndrome. *Modern Rheumatology* 2015;-:1-22.
1186. Legros, L., E. Bardou-Jacquet, et al. Non-invasive assessment of liver fibrosis in C282Y homozygous HFE hemochromatosis. *Liver International* 2015;35(6):1731-8.
1187. Lee, S. W., B. K. Kim, et al. Disease duration and Medsger's severity score are associated with significant liver fibrosis in patients with systemic sclerosis. *Clinical and Experimental Rheumatology* 2015;33(4 Suppl 91):S68-74.
1188. Lee, S. H., S. U. Kim, et al. Use of transient elastography to predict de novo recurrence after radiofrequency ablation for hepatocellular carcinoma. *OncoTargets and therapy* 2015;8:347-56.
1189. Lanthier, N., P. Starkel, et al. New imaging assisted methods for liver fibrosis quantification: Is it really favorable to classical transient elastography? *Journal of Hepatology* 2015;63(3):765-6.
1190. Kikuchi, M., R. Umeda, et al. [Diagnostic accuracy for alcoholic liver disease with controlled Attenuation Parameter (CAP) measured by transient elastography for the non-invasive assessment of liver steatosis]. *Nihon Arukoru Yakubutsu Igakkai Zasshi* 2015;50(5):235-47.
1191. Lai, M. Is liver stiffness measurement to stage fibrosis in patients with nonalcoholic fatty liver disease ready for clinical use? *Hepatology* 2015;62(4):997-8.
1192. Lampertico, P., M. Maini, et al. Optimal management of hepatitis B virus infection - EASL Special Conference. *Journal of Hepatology* 2015;63(5):1238-53.
1193. Lampertico, P., F. Invernizzi, et al. The long-term benefits of nucleos(t)ide analogs in

- compensated HBV cirrhotic patients with no or small esophageal varices: A 12-year prospective cohort study. *Journal of Hepatology* 2015;63(5):1118-25.
1194. Kim, J. K., K. S. Lee, et al. Usefulness of the controlled attenuation parameter for detecting liver steatosis in health checkup examinees. *Gut Liver* 2015;9(3):405-10.
1195. Latt, N. L., F. Araz, et al. Management of hepatitis C infection among patients with renal failure. *Minerva Gastroenterologica e Dietologica* 2015;61(1):39-49.
1196. Kim, M. N., S. U. Kim, et al. Increased risk of hepatocellular carcinoma in chronic hepatitis B patients with transient elastography-defined subclinical cirrhosis. *Hepatology* 2015;61(6):1851-9.
1197. Kucharska, M., M. Inglot, et al. Current Possibilities to Assess the Degree of Liver Fibrosis in Patients with Haemophilia Infected with HCV--Review. *Advances in Clinical and Experimental Medicine* 2015;24(4):671-7.
1198. Kobayashi, K., H. Nakao, et al. Diagnostic accuracy of real-time tissue elastography for the staging of liver fibrosis: a meta-analysis. *European Radiology* 2015;25(1):230-8.
1199. Karlas, T., J. Kollmeier, et al. Noninvasive characterization of graft steatosis after liver transplantation. *Scandinavian Journal of Gastroenterology* 2015;50(2):224-32.
1200. Lammers, W. J., G. M. Hirschfield, et al. Development and Validation of a Scoring System to Predict Outcomes of Patients With Primary Biliary Cirrhosis Receiving Ursodeoxycholic Acid Therapy. *Gastroenterology* 2015;149(7):1804-1812 e4.
1201. Kim, H. J., H. K. Lee, et al. Quantitative comparison of transient elastography (TE), shear wave elastography (SWE) and liver biopsy results of patients with chronic liver disease. *Journal of Physical Therapy Science* 2015;27(8):2465-8.
1202. Kassaye, S., Y. Li, et al. Direct and Indirect Serum Markers of Liver Fibrosis Compared with Transient Elastography among Women in the Women's Interagency HIV Study. *Journal of AIDS & clinical research* 2015;6(4).
1203. Kim, J. H., M. N. Kim, et al. Clinical application of transient elastography in patients with chronic viral hepatitis receiving antiviral treatment. *Liver International* 2015;35(4):1103-15.
1204. Kitson, M. T., S. K. Roberts, et al. Liver stiffness and the prediction of clinically significant portal hypertension and portal hypertensive complications. *Scandinavian Journal of Gastroenterology* 2015;50(4):462-9.
1205. Khunpakdee, N., K. Jayanama, et al. Transient Elastography in End-Stage Renal Disease Patients on Hemodialysis: The Effect of Net Fluid Withdrawal. *Blood purification* 2015;40(3):256-9.
1206. Labarga, P., J. V. Fernandez-Montero, et al. Liver fibrosis progression despite HCV cure with antiviral therapy in HIV-HCV-coinfected patients. *Antiviral Therapy* 2015;20(3):329-34.
1207. Kan, V. Y., V. Marquez Azalgara, et al. Patient preference and willingness to pay for transient elastography versus liver biopsy: A perspective from British Columbia. *Canadian Journal of Gastroenterology & Hepatology* 2015;29(2):72-6.
1208. Kaffenberger, B. H., J. A. Kaffenberger, et al. Magnetic resonance elastography and transient elastography as non-invasive analyses for liver fibrosis: can they obviate the need for liver biopsy in psoriasis patients treated with methotrexate? *International journal of dermatology* 2015;54(7):752-6.
1209. Jung, K. S., S. U. Kim, et al. Validation of hepatitis B virus-related hepatocellular carcinoma prediction models in the era of antiviral therapy. *Hepatology* 2015;62(6):1757-1766.
1210. Karlas, T., J. Berger, et al. Estimating steatosis and fibrosis: Comparison of acoustic structure quantification with established techniques. *World Journal of Gastroenterology* 2015;21(16):4894-902.
1211. Karlas, T., A. Dietrich, et al. Evaluation of Transient Elastography, Acoustic Radiation Force Impulse Imaging (ARFI), and Enhanced Liver Function (ELF) Score for Detection of Fibrosis in Morbidly Obese Patients. *PLoS ONE* 2015;10(11):e0141649.
1212. Huet, N., I. Denis, et al. Ultrasonographic assessment of liver fibrosis with computer-assisted analysis of liver surface irregularities. *Diagnostic and Interventional Imaging* 2015;96(9):941-6.
1213. Iurlo, A., D. Cattaneo, et al. Transient elastography spleen stiffness measurements in primary myelofibrosis patients: a pilot study in a single centre. *British Journal of Haematology* 2015;170(6):890-2.
1214. Jalal, Z., X. Iriart, et al. Liver stiffness measurements for evaluation of central venous pressure in congenital heart diseases. *Heart* 2015;101(18):1499-504.
1215. Jaroszewicz, J., M. Flisiak-Jackiewicz, et al. Current drugs in early development for treating hepatitis C virus-related hepatic fibrosis. Expert opinion on investigational drugs 2015;24(9):1229-39.
1216. Herzer, K., A. Papadopoulos-Kohn, et al. Management of telaprevir-based triple therapy for hepatitis C virus recurrence post liver transplant. *World Journal of Hepatology* 2015;7(9):1287-96.
1217. Ichikawa, S., U. Motosugi, et al. Hepatitis activity should be considered a confounder of liver stiffness measured with MR elastography. *Journal of Magnetic Resonance Imaging* 2015;41(5):1203-8.
1218. Hedenstierna, M., O. Weiland, et al. Long-term follow-up of successful hepatitis C virus therapy:

- waning immune responses and disappearance of liver disease are consistent with cure. *Alimentary Pharmacology & Therapeutics* 2015;41(6):532-43.
1219. Honsawek, S., W. Udomsinprasert, et al. Elevated serum periostin is associated with liver stiffness and clinical outcome in biliary atresia. *Biomarkers* 2015;20(2):157-61.
1220. Hu, Z., Y. Li, et al. Using Ultrasonic Transient Elastometry (FibroScan) to Predict Esophageal Varices in Patients with Viral Liver Cirrhosis. *Ultrasound in Medicine and Biology* 2015;41(6):1530-7.
1221. Juarez-Hernandez, E., M. H. Uribe-Ramos, et al. Factors Associated with the Quality of Transient Elastography. *Digestive Diseases & Sciences* 2015;60(7):2177-82.
1222. Ichikawa, S., U. Motosugi, et al. Comparison of the diagnostic accuracies of magnetic resonance elastography and transient elastography for hepatic fibrosis. *Magnetic resonance imaging* 2015;33(1):26-30.
1223. Hua, J., G. Q. Liu, et al. The role of liver stiffness measurement in the evaluation of liver function and esophageal varices in cirrhotic patients. *Journal of Digestive Diseases* 2015;16(2):98-103.
1224. Hou, J. L., D. Xu, et al. Long-Term Telbivudine Treatment Results in Resolution of Liver Inflammation and Fibrosis in Patients with Chronic Hepatitis B. *Advances in therapy* 2015;32(8):727-41.
1225. Hodowanec, A. C., R. D. Lee, et al. A matched cross-sectional study of the association between circulating tissue factor activity, immune activation and advanced liver fibrosis in hepatitis C infection. *BMC Infectious Diseases* 2015;15(1):190.
1226. Jia, J., J. Hou, et al. Transient elastography compared to serum markers to predict liver fibrosis in a cohort of Chinese patients with chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2015;30(4):756-62.
1227. Hernandez-Gea, V. and A. Berzigotti. Clinical Evaluation and Prognosis. *Digestive Diseases* 2015;33(4):515-23.
1228. Ichikawa, K., Y. Narita, et al. Transient elastography-derived liver stiffness measurements were found to be useful for predicting liver infiltration in a case of mature T-cell neoplasm involving liver dysfunction. *International journal of clinical and experimental pathology* 2015;8(4):4220-6.
1229. Jansen, C., T. Reiberger, et al. Circulating MiRNA-122 Levels Are Associated with Hepatic Necroinflammation and Portal Hypertension in HIV/HCV Coinfection. *PLoS ONE* 2015;10(2):e0116768.
1230. Gerber, L., D. Kasper, et al. Assessment of Liver Fibrosis with 2-D Shear Wave Elastography in Comparison to Transient Elastography and Acoustic Radiation Force Impulse Imaging in Patients with Chronic Liver Disease. *Ultrasound in Medicine and Biology* 2015;41(9):2350-9.
1231. Hagan, M., S. K. Asrani, et al. Non-invasive assessment of liver fibrosis and prognosis. *Expert Review of Gastroenterology & Hepatology* 2015;9(10):1251-60.
1232. He, Y., J. Zhu, et al. Advanced glycation end product (AGE)-induced hepatic stellate cell activation via autophagy contributes to hepatitis C-related fibrosis. *Acta Diabetologica* 2015;52(5):959-69.
1233. Hartman, J., K. Bichoupan, et al. Re-treatment of hepatitis C virus: Eight patients who relapsed twice after direct-acting-antiviral drugs. *World Journal of Gastroenterology* 2015;21(43):12430-8.
1234. Harkissoen, S., J. E. Arends, et al. Low serum hyaluronic acid levels associated with spontaneous HBsAg clearance. *European Journal of Clinical Microbiology & Infectious Diseases* 2015;34(11):2183-9.
1235. Harrison, S. A. Utilization of FibroScan Testing in Hepatitis C Virus Management. *Gastroenterology & Hepatology* 2015;11(3):187-9.
1236. Gherlan, G. S. Liver ultrasound elastography: More than staging the disease. *World Journal of Gastroenterology* 2015;7(12):1595-600.
1237. Guechot, J. Non-invasive evaluation of liver fibrosis: More well-validated tests available for patient management. *Liver International* 2015;35(6):1643-5.
1238. Halasz, T., G. Horvath, et al. miR-122 negatively correlates with liver fibrosis as detected by histology and FibroScan. *World Journal of Gastroenterology* 2015;21(25):7814-23.
1239. Hasemi, F., A. Fathi Ashtiani, et al. Impact of Hepatitis C Virus Infection on Cognitive Function in Patients With Covert Hepatic Encephalopathy. *Hepatitis Monthly* 2015;15(7):e30507.
1240. He, Y., L. Jin, et al. Mechanisms of fibrosis in acute liver failure. *Liver International* 2015;35(7):1877-85.
1241. Gobel, T., J. Schadewaldt-Tummers, et al. Transient elastography improves detection of liver cirrhosis compared to routine screening tests. *World Journal of Gastroenterology* 2015;21(3):953-60.
1242. Gouvea Ade, F., D. M. Machado, et al. [Noncirrhotic portal hypertension in a human immunodeficiency virus (HIV) infected adolescent]. *Revista Paulista de Pediatria* 2015;33(2):246-50.
1243. Gill, U. S. and P. T. Kennedy. New insights in the management of chronic hepatitis B. *Clinical Medicine* 2015;15(2):191-6.
1244. Gonzalez, F. A., E. Van den Eynde, et al. Liver stiffness and aspartate aminotransferase levels predict the risk for liver fibrosis progression in

- hepatitis C virus/HIV-coinfected patients. *HIV Medicine* 2015;16(4):211-8.
1245. Goldschmidt, I., A. Di Nanni, et al. Improvement of BMI after Lifestyle Intervention Is Associated with Normalisation of Elevated ELF Score and Liver Stiffness in Obese Children. *BioMed Research International* 2015;2015:457473.
1246. Harman, D. J., S. D. Ryder, et al. Direct targeting of risk factors significantly increases the detection of liver cirrhosis in primary care: a cross-sectional diagnostic study utilising transient elastography. *BMJ Open* 2015;5(4):e007516.
1247. Festi, D., R. Schiumerini, et al. Non-invasive diagnostic approach to non-alcoholic fatty liver disease: current evidence and future perspectives. *Expert Review of Gastroenterology & Hepatology* 2015;9(8):1039-53.
1248. Ferraioli, G., C. Tinelli, et al. Correlation of the controlled attenuation parameter with indices of liver steatosis in overweight or obese individuals: a pilot study. *European Journal of Gastroenterology & Hepatology* 2015;27(3):305-312.
1249. Friedrich-Rust, M., M. Lupsor, et al. Point Shear Wave Elastography by Acoustic Radiation Force Impulse Quantification in Comparison to Transient Elastography for the Noninvasive Assessment of Liver Fibrosis in Chronic Hepatitis C: A Prospective International Multicenter Study. *Ultraschall in der Medizin* 2015;36(3):239-47.
1250. Gaia, S., D. Campion, et al. Non-invasive score system for fibrosis in chronic hepatitis: proposal for a model based on biochemical, FibroScan and ultrasound data. *Liver International* 2015;35(8):2027-35.
1251. Fernandez, M., E. Trepo, et al. Transient elastography using Fibroscan is the most reliable noninvasive method for the diagnosis of advanced fibrosis and cirrhosis in alcoholic liver disease. *European Journal of Gastroenterology & Hepatology* 2015;27(9):1074-9.
1252. Fernandes, F. F., M. L. Ferraz, et al. Enhanced liver fibrosis panel as a predictor of liver fibrosis in chronic hepatitis C patients. *Journal of Clinical Gastroenterology* 2015;49(3):235-41.
1253. Ergelen, R., U. Akyuz, et al. Measurements of serum procollagen-III peptide and M30 do not improve the diagnostic accuracy of transient elastography for the detection of hepatic fibrosis in patients with nonalcoholic fatty liver disease. *European Journal of Gastroenterology & Hepatology* 2015;27(6):667-71.
1254. El-Haggar, S. M. and T. M. Mostafa. Comparative clinical study between the effect of fenofibrate alone and its combination with pentoxifylline on biochemical parameters and liver stiffness in patients with non-alcoholic fatty liver disease. *Hepatology International* 2015;9(3):471-9.
1255. Egresi, A., G. Lengyel, et al. [Options for non-invasive assessment of liver fibrosis based on clinical data]. *Orvosi Hetilap* 2015;156(2):43-52.
1256. Emamgholipour, S., N. Moradi, et al. The association of circulating levels of complement-C1q TNF-related protein 5 (CTRP5) with nonalcoholic fatty liver disease and type 2 diabetes: a case-control study. *Diabetology & metabolic syndrome* 2015;7:108.
1257. Dincses, E. and Y. Yilmaz. Diagnostic usefulness of FibroMeter VCTE for hepatic fibrosis in patients with nonalcoholic fatty liver disease. *European Journal of Gastroenterology & Hepatology* 2015;27(10):1149-53.
1258. Dugum, M. and A. McCullough. Diagnosis and Management of Alcoholic Liver Disease. *Journal of Clinical and Translational Hepatology* 2015;3(2):109-16.
1259. Dessi, C., G. Leoni, et al. Thalassemia major between liver and heart: Where we are now. *Blood Cells, Molecules & Diseases* 2015;55(1):82-8.
1260. Dong, D. R., M. N. Hao, et al. Acoustic radiation force impulse elastography, FibroScan(R), Forns' index and their combination in the assessment of liver fibrosis in patients with chronic hepatitis B, and the impact of inflammatory activity and steatosis on these diagnostic methods. *Molecular Medicine Reports* 2015;11(6):4174-82.
1261. Deffieux, T., J. L. Gennisson, et al. Investigating liver stiffness and viscosity for fibrosis, steatosis and activity staging using shear wave elastography. *Journal of Hepatology* 2015;62(2):317-24.
1262. Ding, D., H. Li, et al. FibroScan, aspartate aminotransferase and alanine aminotransferase ratio (AAR), aspartate aminotransferase to platelet ratio index (APRI), fibrosis index based on the 4 factor (FIB-4), and their combinations in the assessment of liver fibrosis in patients with hepatitis B. *International journal of clinical and experimental medicine* 2015;8(11):20876-82.
1263. Dajani, A. I., A. M. Abu Hammour, et al. Essential phospholipids as a supportive adjunct in the management of patients with NAFLD. *Arab journal of gastroenterology* 2015;16(3-4):99-104.
1264. Cohort, A. C. H. Regression of liver stiffness after sustained hepatitis C virus (HCV) virological responses among HIV/HCV-coinfected patients. *Aids* 2015;29(14):1821-30.
1265. Curran, A., R. Marti, et al. Darunavir and ritonavir total and unbound plasmatic concentrations in HIV-HCV-coinfected patients with hepatic cirrhosis compared to those in HIV-monoinfected patients. *Antimicrobial Agents and Chemotherapy* 2015;59(11):6782-90.
1266. Cohen-Ezra, O. and Z. Ben-Ari. [Non-invasive assessment of liver fibrosis]. *Harefuah* 2015;154(3):204-7, 210, 209.

1267. Chon, Y. E., K. S. Jung, et al. Normal controlled attenuation parameter values: a prospective study of healthy subjects undergoing health checkups and liver donors in Korea. *Digestive & Liver Disease* 2015;60(1):234-42.
1268. Besharat, S., H. Poustchi, et al. Central Obesity and Advanced Liver Stiffness in Hepatitis B: Result from Golestan Hepatitis B Cohort Study. *Archives of Iranian medicine* 2015;18(9):562-6.
1269. Bota, S., R. Paternostro, et al. Performance of 2-D Shear Wave Elastography in Liver Fibrosis Assessment Compared with Serologic Tests and Transient Elastography in Clinical Routine. *Ultrasound in Medicine and Biology* 2015;41(9):2340-9.
1270. Cassinotto, C. and V. de Ledinghen. Reply to: "New imaging assisted methods for liver fibrosis quantification: Is it really favorable to classical transient elastography?". *Journal of Hepatology* 2015;63(3):767.
1271. Berzigotti, A., T. D. Boyer, et al. Reply to "Points to be considered when using transient elastography for diagnosis of portal hypertension according to the Baveno's VI consensus". *Journal of Hepatology* 2015;63(4):1049-50.
1272. Chan, W. K., N. R. Nik Mustapha, et al. A novel 2-step approach combining the NAFLD fibrosis score and liver stiffness measurement for predicting advanced fibrosis. *Hepatology International* 2015;9(4):594-602.
1273. Chen, T., R. Wong, et al. Occult cirrhosis diagnosed by transient elastography is a frequent and under-monitored clinical entity. *Liver International* 2015;35(10):2285-93.
1274. Cardoso, A. C., M. Beaugrand, et al. Diagnostic performance of controlled attenuation parameter for predicting steatosis grade in chronic hepatitis B. *Annals of Hepatology* 2015;14(6):826-36.
1275. Barone, M., A. Iannone, et al. Liver stiffness and portal blood flow modifications induced by a liquid meal consumption: pathogenetic mechanisms and clinical relevance. *Scandinavian Journal of Gastroenterology* 2015;50(5):560-6.
1276. Borgia, S. M. FibroScan(R) access in Canada: Time for reform, a call for universal access. *Canadian Journal of Gastroenterology & Hepatology* 2015;29(4):221-2.
1277. Cabibi, D., V. Calvaruso, et al. Comparison of Histochemical Staining Methods and Correlation with Transient Elastography in Acute Hepatitis. *Pathobiology* 2015;82(1):48-52.
1278. Bintintan, A., R. I. Chira, et al. Value of hepatic elastography and Doppler indexes for predictions of esophageal varices in liver cirrhosis. *Medical Ultrasonography Journal* 2015;17(1):5-11.
1279. Bonnard, P., A. Elsharkawy, et al. Comparison of liver biopsy and noninvasive techniques for liver fibrosis assessment in patients infected with HCV- genotype 4 in Egypt. *Journal of viral hepatitis* 2015;22(3):245-53.
1280. Carrion, J. A., M. Puigvehi, et al. Applicability and accuracy improvement of transient elastography using the M and XL probes by experienced operators. *Journal of viral hepatitis* 2015;22(3):297-306.
1281. Chavez-Tapia, N., J. Torres-Sanchez, et al. Prevalence in vulnerable population of liver fibrosis identified by transient elastography. *Annals of Hepatology* 2015;14(4):524-30.
1282. Cales, P., J. Boursier, et al. Cirrhosis Diagnosis and Liver Fibrosis Staging: Transient Elastometry Versus Cirrhosis Blood Test. *Journal of Clinical Gastroenterology* 2015;49(6):512-9.
1283. Besharat, S., H. Poustchi, et al. Association of Mutations in the Basal Core Promoter and Pre-core Regions of the Hepatitis B Viral Genome and Longitudinal Changes in HBV Level in HBeAg Negative Individuals: Results From a Cohort Study in Northern Iran. *Hepatitis Monthly* 2015;15(2):e23875.
1284. Chin, J. L., G. Chan, et al. Spleen stiffness can non-invasively assess resolution of portal hypertension after liver transplantation. *Liver International* 2015;35(2):518-23.
1285. Castera, L. Is it really worth adapting liver stiffness cut-offs according to AST levels? *Liver International* 2015;35(12):2495-7.
1286. Cassinotto, C., M. Feldis, et al. MR relaxometry in chronic liver diseases: Comparison of T1 mapping, T2 mapping, and diffusion-weighted imaging for assessing cirrhosis diagnosis and severity. *European Journal of Radiology* 2015;84(8):1459-65.
1287. Cassinotto, C., A. Charrie, et al. Liver and spleen elastography using supersonic shear imaging for the non-invasive diagnosis of cirrhosis severity and oesophageal varices. *Digestive & Liver Disease* 2015;47(8):695-701.
1288. Berzigotti, A., M. Reig, et al. Value of transient elastography measured with fibroscan in predicting the outcome of hepatic resection for hepatocellular carcinoma. *Annals of Surgery* 2015;261(4):e105.
1289. Cescon, M., A. Cucchetti, et al. Reply to letter: "value of transient elastography measured with fibroscan in predicting the outcome of hepatic resection for hepatocellular carcinoma". *Annals of Surgery* 2015;261(4):e105-6.
1290. Barone, M., M. T. Viggiani, et al. Endothelial dysfunction correlates with liver fibrosis in chronic HCV infection. *Gastroenterology Research & Practice* 2015;2015:682174.
1291. Bocchino, M., G. Brancaccio, et al. Increased liver stiffness in idiopathic pulmonary fibrosis: a pilot study. *Sarcoidosis, Vasculitis, and Diffuse Lung Diseases* 2015;32(2):176-80.

1292. Buzzetti, E., R. Lombardi, et al. Noninvasive Assessment of Fibrosis in Patients with Nonalcoholic Fatty Liver Disease. *International Journal of Endocrinology* 2015;2015:343828.
1293. Cabibi, D., F. Bronte, et al. Comparison of Histochemical Stainings in Evaluation of Liver Fibrosis and Correlation with Transient Elastography in Chronic Hepatitis. *Analytical Cellular Pathology* 2015;2015:431750.
1294. Castera, L. Noninvasive Assessment of Liver Fibrosis. *Digestive Diseases & Sciences* 2015;33(4):498-503.
1295. Chan, R., V. W. Wong, et al. Higher estimated net endogenous Acid production may be associated with increased prevalence of nonalcoholic Fatty liver disease in chinese adults in Hong Kong. *PLoS ONE* 2015;10(4):e0122406.
1296. Cheng, J., J. Hou, et al. Validation of Ten Noninvasive Diagnostic Models for Prediction of Liver Fibrosis in Patients with Chronic Hepatitis B. *PLoS ONE* 2015;10(12):e0144425.
1297. Cho, Y., D. Tokuhara, et al. Transient Elastography-Based Liver Profiles in a Hospital-Based Pediatric Population in Japan. *PLoS ONE* 2015;10(9):e0137239.
1298. Cho, E. J., M. Y. Kim, et al. Diagnostic and Prognostic Values of Noninvasive Predictors of Portal Hypertension in Patients with Alcoholic Cirrhosis. *PLoS ONE* 2015;10(7):e0133935.
1299. Aube, C. Imaging modalities for the diagnosis of hepatic fibrosis and cirrhosis. *Clinics and Research in Hepatology and Gastroenterology* 2015;39(1):38-44.
1300. Asrani, S. K. Incorporation of Noninvasive Measures of Liver Fibrosis Into Clinical Practice: Diagnosis and Prognosis. *Clinical Gastroenterology & Hepatology* 2015;13(12):2190-204.
1301. Aljawad, M., S. Sirpal, et al. Transient elastography in Canada: current state and future directions. *Canadian Journal of Gastroenterology & Hepatology* 2015;29(7):373-6.
1302. Al-Hamoudi, W. K., A. A. Abdelrahman, et al. The role of Fibroscan in predicting the presence of varices in patients with cirrhosis. *European Journal of Gastroenterology & Hepatology* 2015;27(11):1307-12.
1303. Alsebaey, A., N. Allam, et al. Normal liver stiffness: A study in living donors with normal liver histology. *World Journal of Hepatology* 2015;7(8):1149-53.
1304. AlShaalan, R., M. Aljiffry, et al. Nonalcoholic fatty liver disease: Noninvasive methods of diagnosing hepatic steatosis. *Saudi Journal of Gastroenterology* 2015;21(2):64-70.
1305. Arasaradnam, R. P., M. McFarlane, et al. Non-invasive distinction of non-alcoholic fatty liver disease using urinary volatile organic compound analysis: early results. *Journal of Gastrointestinal & Liver Diseases* 2015;24(2):197-201.
1306. Andreasson, K., R. Hesselstrand, et al. Cartilage oligomeric matrix protein: a new promising biomarker of liver fibrosis in chronic hepatitis C. *Infectious diseases* 2015;47(12):915-8.
1307. Alvarez, D., F. Orozco, et al. Meal ingestion markedly increases liver stiffness suggesting the need for liver stiffness determination in fasting conditions. *Gastroenterologia y hepatología* 2015;38(7):431-5.
1308. Afdhal, N. H., B. R. Bacon, et al. Accuracy of fibroscan, compared with histology, in analysis of liver fibrosis in patients with hepatitis B or C: a United States multicenter study. *Clinical Gastroenterology & Hepatology* 2015;13(4):772-779 e3.
1309. Agrawal, S., A. Duseja, et al. Obstructive sleep apnea is an important predictor of hepatic fibrosis in patients with nonalcoholic fatty liver disease in a tertiary care center. *Hepatology International* 2015;9(2):283-91.

2014

1310. Yuan, L., J. Shao, et al. [Correlation between liver hardness testing results obtained by FibroTouch and FibroScan and liver pathological stage]. *Zhonghua Gan Zang Bing Za Zhi* 2014;22(6):425-9.
1311. Zeuzem, S., I. M. Jacobson, et al. Retreatment of HCV with ABT-450/r-ombitasvir and dasabuvir with ribavirin. *The New England Journal of Medicine* 2014;370(17):1604-14.
1312. Yoo, B. W., J. Y. Choi, et al. Congestive hepatopathy after Fontan operation and related factors assessed by transient elastography. *Journal of thoracic and cardiovascular surgery* 2014;148(4):1498-505.
1313. Yoon, J. H., J. M. Lee, et al. Hepatic fibrosis: prospective comparison of MR elastography and US shear-wave elastography for evaluation. *Radiology* 2014;273(3):772-82.
1314. Wu, F. M., A. R. Opotowsky, et al. Transient Elastography May Identify Fontan Patients with Unfavorable Hemodynamics and Advanced Hepatic Fibrosis. *Congenital Heart Disease* 2014;9(5):438-447.
1315. Wong, G. L. Prediction of fibrosis progression in chronic viral hepatitis. *Clinical and Molecular Hepatology* 2014;20(3):228-236.
1316. Wang, J. H., S. K. Chuah, et al. Baseline and serial liver stiffness measurement in prediction of portal hypertension progression for patients with compensated cirrhosis. *Liver International* 2014;34(9):1340-8.
1317. Yilmaz, Y. and F. Eren. A Bayesian approach to an integrated multimodal noninvasive diagnosis of

- definitive nonalcoholic steatohepatitis in the spectrum of nonalcoholic fatty liver disease. European Journal of Gastroenterology & Hepatology 2014;26(11):1292-5.
1318. Yilmaz, Y., A. Yesil, et al. Detection of hepatic steatosis using the Controlled Attenuation Parameter: a comparative study with liver biopsy. Scandinavian Journal of Gastroenterology 2014;49(5):611-6.
1319. Wu, T., L. Li, et al. Transient elastography in hepatitis B virus infection. Liver stiffness discrepancy due to sampling location. Saudi medical journal 2014;35(6):554-60.
1320. Whitty, J. A., C. Tallis, et al. Cost and time savings from a rapid access model of care using transient elastography to screen and triage patients with chronic Hepatitis C infection. Journal of Medical Economics 2014;17(2):159-65.
1321. Wong, G. L., H. L. Chan, et al. Liver stiffness-based optimization of hepatocellular carcinoma risk score in patients with chronic hepatitis B. Journal of Hepatology 2014;60(2):339-45.
1322. Wang, C. Y., W. Lu, et al. Diagnostic value of controlled attenuation parameter for liver steatosis in patients with chronic hepatitis B. World Journal of Gastroenterology 2014;20(30):10585-90.
1323. Wu, G., H. He, et al. [Clinical effect of combination therapy with Fufang Biejia Ruangan tablet and entecavir in patients with hepatitis B virus-related cirrhosis]. Zhonghua Gan Zang Bing Za Zhi 2014;22(8):604-8.
1324. Vuppalanchi, R. and K. Lindor. Magic wand and crystal ball for primary sclerosing cholangitis. Gastroenterology 2014;146(4):890-2.
1325. Wilder, J. and K. Patel. The clinical utility of FibroScan as a noninvasive diagnostic test for liver disease. Journal of Medical Devices 2014;7:107-114.
1326. Yada, N., T. Sakurai, et al. Ultrasound elastography correlates treatment response by antiviral therapy in patients with chronic hepatitis C. Oncology 2014;87 Suppl 1:118-23.
1327. Vukobrat-Bijedic, Z., A. Husic-Selimovic, et al. Analysis of effect of antiviral therapy on regression of liver fibrosis in patient with HCV infection. Materia socio-medica 2014;26(3):172-6.
1328. Vonighia, L., P. Michielsen, et al. Diagnostic challenges in alcohol use disorder and alcoholic liver disease. World Journal of Gastroenterology 2014;20(25):8024-32.
1329. Thin, L. W., I. C. Lawrence, et al. Detection of liver injury in IBD using transient elastography. Journal of Crohn's & Colitis 2014;8(7):671-7.
1330. Vergniol, J., J. Boursier, et al. The evolution of non-invasive tests of liver fibrosis is associated with prognosis in patients with chronic hepatitis C. Hepatology 2014;60(1):65-76.
1331. Trabut, J. B., V. Thepot, et al. [Prognosis assessment of alcoholic liver disease: how and why?]. La Presse Médicale 2014;43(2):124-34.
1332. Thielsen, P., R. Wilkens, et al. [Liver fibrosis can be assessed by non-invasive ultrasound elastography.]. Ugeskrift for laeger 2014;176(15).
1333. Tapper, E. B., L. Castera, et al. FibroScan (Vibration Controlled Elastography): Where Does It Stand in the US Practice. Clin Gastroenterol Hepatol. 2014;13(1):27-36.
1334. Muir, A. J. Cirrhosis in hepatitis C virus-infected patients: a review for practitioners new to hepatitis C care. Topics in antiviral medicine 2014;22(4):685-9.
1335. Licinio, R., M. Principi, et al. Liver fibrosis in primary intestinal lymphangiectasia: An undervalued topic. World Journal of Hepatology 2014;6(9):685-7.
1336. Leeming, D. J., E. Anadol, et al. Combined antiretroviral therapy attenuates hepatic extracellular matrix remodeling in HIV patients assessed by novel protein fingerprint markers. Aids 2014;28(14):2081-90.
1337. Lupsor-Platon, M., H. Stefanescu, et al. Noninvasive assessment of liver steatosis using ultrasound methods. Medical Ultrasonography Journal 2014;16(3):236-245.
1338. Macaluso, F. S., M. Maida, et al. Steatosis affects the performance of liver stiffness measurement for fibrosis assessment in patients with genotype 1 chronic hepatitis C. Journal of Hepatology 2014;61(3):523-9.
1339. Morling, J. R., J. A. Fallowfield, et al. Non-invasive hepatic biomarkers (ELF and CK18) in people with type 2 diabetes: the Edinburgh type 2 diabetes study. Journal of Hepatology 2014;34(8):1267-77.
1340. Ou, X. [Clinical application of transient elastography for assessment of chronic liver diseases]. Zhonghua Gan Zang Bing Za Zhi 2014;22(9):650-3.
1341. Sharma, P., S. Dhawan, et al. The usefulness of transient elastography by FibroScan for the evaluation of liver fibrosis. Indian Journal of Gastroenterology 2014;33(5):445-51.
1342. Shen, F. and J. Fan. [Current status of transient elastography for assessing patients with fatty liver disease]. Zhonghua Gan Zang Bing Za Zhi 2014;22(9):643-6.
1343. Morosan, E. and M. S. Mihailovici. Fibrosis assessment in chronic hepatitis C--is the liver biopsy still necessary? The pathologist point of view. Revista medico-chirurgicala a Societatii de Medici si Naturalisti din Iasi 2014;118(4):992-9.
1344. Mueller, S., H. K. Seitz, et al. Non-invasive diagnosis of alcoholic liver disease. World Journal of Gastroenterology 2014;20(40):14626-14641.

1345. Soriano, V., E. Vispo, et al. Efficacy of prolonged tenofovir therapy on hepatitis delta in HIV-infected patients. *Aids* 2014;28(16):2389-94.
1346. Naveau, S., K. Lamouri, et al. The diagnostic accuracy of transient elastography for the diagnosis of liver fibrosis in bariatric surgery candidates with suspected NAFLD. *Obesity surgery* 2014;24(10):1693-1701.
1347. Park, J. S., M. C. Park, et al. Concurrent use of methotrexate and celecoxib increases risk of silent liver fibrosis in rheumatoid arthritis patients with subclinical reduced kidney function. *Clinical Rheumatology* 2014;33(10):1415-23.
1348. Tangpradabkiet, W., S. Praneenararat, et al. Influence of meal intake on liver stiffness in patients with chronic hepatitis B and C. *Journal of the Medical Association of Thailand* 2014;97(10):1033-9.
1349. Liu, J. H., S. Yan, et al. [Application of transient elastography in early prognosis after liver transplantation]. *Zhejiang Da Xue Xue Bao Yi Xue Ban* 2014;43(6):678-82.
1350. Mazzocato, S., E. Orsetti, et al. Comparison of liver fibrosis progression in HIV/HCV co-infected and HCV mono-infected patients by transient elastometry. *Scandinavian Journal of Infectious Diseases* 2014;46(11):797-802.
1351. Oda, K., H. Uto, et al. Impact of a single nucleotide polymorphism upstream of the IL28B gene in patients positive for anti-HCV antibody in an HCV hyperendemic area in Japan. *Journal of Medical Virology* 2014;86(11):1877-85.
1352. Poynard, T., J. Vergniol, et al. Staging chronic hepatitis B into seven categories, defining inactive carriers and assessing treatment impact using a fibrosis biomarker (FibroTest(R)) and elastography (FibroScan(R)). *Journal of Hepatology* 2014;61(5):994-1003.
1353. Macaluso, F. S., M. Maida, et al. Body mass index and liver stiffness affect accuracy of ultrasonography in detecting steatosis in patients with chronic hepatitis C virus genotype 1 infection. *Clinical Gastroenterology & Hepatology* 2014;12(5):878-884 e1.
1354. Patel, K. and N. A. Shackel. Current status of fibrosis markers. *Current Opinion in Gastroenterology* 2014;30(3):253-9.
1355. Shin, N. Y., M. J. Kim, et al. Transient elastography and sonography for prediction of liver fibrosis in infants with biliary atresia. *Journal of ultrasound in medicine* 2014;33(5):853-64.
1356. Lemoine, M., Y. Shimakawa, et al. Commentary: how long does one need to fast before a Fibroscan examination? Authors' reply. *Alimentary Pharmacology & Therapeutics* 2014;39(5):541-2.
1357. Orlic, L., I. Mikolasevic, et al. Chronic kidney disease and Nonalcoholic Fatty Liver Disease - is there a link? *Gastroenterology Research & Practice* 2014;2014:6.
1358. Pang, J. X., F. Pradhan, et al. The feasibility and reliability of transient elastography using FibroScan(R): a practice audit of 2335 examinations. *Canadian Journal of Gastroenterology & Hepatology* 2014;28(3):143-9.
1359. Piscaglia, F., S. Marinelli, et al. The role of ultrasound elastographic techniques in chronic liver disease: current status and future perspectives. *European Journal of Radiology* 2014;83(3):450-5.
1360. Sporea, I., S. Bota, et al. Which are the cut-off values of 2D-Shear Wave Elastography (2D-SWE) liver stiffness measurements predicting different stages of liver fibrosis, considering Transient Elastography (TE) as the reference method? *European Journal of Radiology* 2014;83(3):e118-22.
1361. Macias, J., J. Gonzalez, et al. Prevalence and factors associated with liver steatosis as measured by transient elastography with controlled attenuation parameter in HIV-infected patients. *Aids* 2014;28(9):1279-1287.
1362. McCormick, P. A., E. Elrayah, et al. Letter: single liver stiffness measurement predicts liver-related mortality in chronic hepatitis C. *Alimentary Pharmacology & Therapeutics* 2014;39(12):1438-9.
1363. Mikolasevic, I., S. Racki, et al. Nonalcoholic Fatty liver disease in renal transplant recipients proven by transient elastography. *Transplantation Proceedings* 2014;46(5):1347-52.
1364. Parra-Ruiz, J., C. Sanjuan, et al. Letter: accuracy of liver stiffness measurement - a comparison of two different FibroScan devices. *Alimentary Pharmacology & Therapeutics* 2014;39(12):1434-5.
1365. Shi, K. Q., J. Z. Tang, et al. Controlled attenuation parameter for the detection of steatosis severity in chronic liver disease: A meta-analysis of diagnostic accuracy. *Journal of Gastroenterology & Hepatology* 2014;29(6):1149-58.
1366. Singh, S., J. E. Eaton, et al. Accuracy of spleen stiffness measurement in detection of esophageal varices in patients with chronic liver disease: systematic review and meta-analysis. *Clinical Gastroenterology & Hepatology* 2014;12(6):935-45 e4.
1367. Sirli, R., I. Sporea, et al. Comparison between the M and XL probes for liver fibrosis assessment by transient elastography. *Medical Ultrasonography Journal* 2014;16(2):119-22.
1368. Mikolasevic, I., L. Orlic, et al. Non-alcoholic fatty liver disease proven by transient elastography in hemodialysis patients: is it a new risk factor for adverse cardiovascular events? *Blood purification* 2014;37(4):259-65.
1369. Lukenda, V., I. Mikolasevic, et al. Transient elastography: a new noninvasive diagnostic tool for assessment of chronic allograft nephropathy. *International Urology and Nephrology* 2014;46(7):1435-40.

1370. Papatheodoridis, G. V., S. Manolakopoulos, et al. The usefulness of transient elastography in the assessment of patients with HBeAg-negative chronic hepatitis B virus infection. *Journal of viral hepatitis* 2014;21(7):517-24.
1371. Park, M. S., K. H. Han, et al. Non-invasive prediction of development of hepatocellular carcinoma using transient elastography in patients with chronic liver disease. *Expert Review of Gastroenterology & Hepatology* 2014;8(5):501-11.
1372. Schenk, J. P., G. Alzen, et al. Measurement of real-time tissue elastography in a phantom model and comparison with transient elastography in pediatric patients with liver diseases. *Diagnostic and Interventional Radiology* 2014;20(1):90-9.
1373. Stasi, C., E. Triboli, et al. Assessment of liver stiffness in patients with HCV and mixed cryoglobulinemia undergoing rituximab treatment. *Journal of translational medicine* 2014;12(1):21.
1374. Lee, S. and Y. Kim do. Non-invasive diagnosis of hepatitis B virus-related cirrhosis. *World Journal of Gastroenterology* 2014;20(2):445-59.
1375. Lemoine, M., Y. Shimakawa, et al. Food intake increases liver stiffness measurements and hampers reliable values in patients with chronic hepatitis B and healthy controls: the PROLIFICA experience in The Gambia. *Alimentary Pharmacology & Therapeutics* 2014;39(2):188-96.
1376. Mena, A., J. D. Pedreira, et al. Metabolic syndrome association with fibrosis development in chronic hepatitis B virus inactive carriers. *Journal of Gastroenterology & Hepatology* 2014;29(1):173-8.
1377. Narita, Y., T. Genda, et al. Prediction of liver stiffness hepatocellular carcinoma in chronic hepatitis C patients on interferon-based anti-viral therapy. *Journal of Gastroenterology & Hepatology* 2014;29(1):137-43.
1378. Sebastiani, G., P. Ghali, et al. Physicians' practices for diagnosing liver fibrosis in chronic liver diseases: A nationwide, Canadian survey. *Canadian Journal of Gastroenterology* 2014;28(1):23-30.
1379. Par, G., A. Trotsis, et al. [Transient elastography as a predictor of oesophageal varices in patients with liver cirrhosis]. *Orvosi Hetilap* 2014;155(7):270-6.
1380. Taniguchi, T., Y. Sakata, et al. Usefulness of transient elastography for noninvasive and reliable estimation of right-sided filling pressure in heart failure. *American Journal of Cardiology* 2014;113(3):552-8.
1381. Lin, L. J., G. F. Chen, et al. [Performance of FibroScan in evaluating the curative effects of traditional Chinese medicine on liver fibrosis]. *Zhonghua Gan Zang Bing Za Zhi* 2014;22(2):113-7.
1382. Mikolasevic, I., S. Racki, et al. Nonalcoholic fatty liver disease (NAFLD): a new risk factor for adverse cardiovascular events in dialysis patients. *Medical hypotheses* 2014;82(2):205-8.
1383. Praneenararat, S., N. Chamroonkul, et al. HBV DNA level to predict liver fibrosis in HBeAg negative chronic hepatitis B patients with biopsy indication. *BMC Gastroenterology* 2014;14(1):218.
1384. Sharma, S., K. Khalili, et al. Non-invasive diagnosis of advanced fibrosis and cirrhosis. *World Journal of Gastroenterology* 2014;20(45):16820-30.
1385. Stefanescu, H. and B. Procopet. Noninvasive assessment of portal hypertension in cirrhosis: liver stiffness and beyond. *World Journal of Gastroenterology* 2014;20(45):16811-9.
1386. Salzl, P., T. Reiberger, et al. Evaluation of portal hypertension and varices by acoustic radiation force impulse imaging of the liver compared to transient elastography and AST to platelet ratio index. *Ultraschall in der Medizin* 2014;35(6):528-33.
1387. Shehab, H., I. Elattar, et al. CUFA algorithm: assessment of liver fibrosis using routine laboratory data. *Journal of viral hepatitis* 2014;21(12):956-64.
1388. Shen, F., R. Zheng, et al. [A multi-center clinical study of a novel controlled attenuation parameter for assessment of fatty liver]. *Zhonghua Gan Zang Bing Za Zhi* 2014;22(12):926-31.
1389. Sebastiani, G., K. Gkouvatsos, et al. Chronic hepatitis C and liver fibrosis. *World Journal of Gastroenterology* 2014;20(32):11033-53.
1390. Tahiri, M., A. Drighil, et al. Chronic permanent hypoxemia predisposes to mild elevation of liver stiffness. *World Journal of Gastroenterology* 2014;20(30):10564-9.
1391. Lee, H. W., E. J. Yoo, et al. Prediction of development of liver-related events by transient elastography in hepatitis B patients with complete virological response on antiviral therapy. *American Journal of Gastroenterology* 2014;109(8):1241-9.
1392. Li, F., T. Yan, et al. [Clinical study of FibroScan efficiency for diagnosing size of oesophageal varices in liver cirrhosis patients]. *Zhonghua Gan Zang Bing Za Zhi* 2014;22(8):600-3.
1393. Lynch, M., E. Higgins, et al. The use of transient elastography and FibroTest for monitoring hepatotoxicity in patients receiving methotrexate for psoriasis. *JAMA dermatology* 2014;150(8):856-62.
1394. Shen, F., R. D. Zheng, et al. Controlled attenuation parameter for non-invasive assessment of hepatic steatosis in Chinese patients. *World Journal of Gastroenterology* 2014;20(16):4702-11.
1395. Malekzadeh, M. M., A. R. Radmard, et al. Juvenile Hemochromatosis, Genetic Study and Long-term Follow up after Therapy. *Middle East journal of digestive diseases* 2014;6(2):87-92.
1396. Poynard, T., J. Vergniol, et al. Staging chronic hepatitis C in seven categories using fibrosis biomarker (FibroTest) and transient elastography

- (FibroScan(R)). *Journal of Hepatology* 2014;60(4):706-14.
1397. Sanmartin, R., J. Tor, et al. Progression of liver fibrosis in HIV/hepatitis C virus-coinfected individuals on antiretroviral therapy with early stages of liver fibrosis at baseline. *HIV Medicine* 2014;15(4):203-12.
1398. Meng-fei, Z., L. Jing, et al. Comparative analysis of controlled attenuation parameters in evaluation of liver steatosis using FibroTouch and FibroScan. *Acta Metallurgica Sinica* 2014;34(6):589-593.
1399. Maev, I. V., A. N. Kaziulin, et al. [Possibilities of using the noninvasive methods of investigating the morphofunctional changes in the liver in patients with non alcoholic steatohepatitis and type 2 diabetes mellitus]. *Eksperimental Klinical Gastroenterology* 2014;-(3):38-45.
1400. Mikolasevic, I., S. Racki, et al. Nonalcoholic Fatty Liver Disease (NAFLD) And Cardiovascular Risk In Renal Transplant Recipients. *Kidney & blood pressure research* 2014;39(4):308-14.
1401. Pang, J. X., S. Zimmer, et al. Liver stiffness by transient elastography predicts liver-related complications and mortality in patients with chronic liver disease. *PLoS ONE* 2014;9(4):e95776.
1402. Lahmek, P., N. Meunier, et al. [Using transient elastography as a screening tool for liver fibrosis in addiction service]. *La Presse Médicale* 2014;43(3):e17-31.
1403. Lam, K. L., A. J. Hui, et al. Commentary: how long does one need to fast before a Fibroscan examination? *Alimentary Pharmacology & Therapeutics* 2014;39(5):540-1.
1404. Labarga, P., J. V. Fernandez-Montero, et al. Changes in liver fibrosis in HIV/HCV-coinfected patients following different outcomes with peginterferon plus ribavirin therapy. *Journal of viral hepatitis* 2014;21(7):475-9.
1405. Labarga, P., J. V. Fernandez-Montero, et al. Progression to advanced liver fibrosis in HIV-HCV-coinfected patients and prioritization of new hepatitis C therapies. *Antiviral Therapy* 2014;19(8):799-803.
1406. Kwok, R., Y. K. Tse, et al. Systematic review with meta-analysis: non-invasive assessment of non-alcoholic fatty liver disease--the role of transient elastography and plasma cytokeratin-18 fragments. *Alimentary Pharmacology & Therapeutics* 2014;39(3):254-69.
1407. Kontogianni, M. D., N. Tileli, et al. Adherence to the Mediterranean diet is associated with the severity of non-alcoholic fatty liver disease. *Clinical Nutrition* 2014;33(4):678-83.
1408. Kuo, Y. H., S. N. Lu, et al. The changes of liver stiffness and its associated factors for chronic hepatitis B patients with entecavir therapy. *PLoS ONE* 2014;9(3):e93160.
1409. Klauser, A. S., H. Miyamoto, et al. Sonoelastography: musculoskeletal applications. *Radiology* 2014;272(3):622-33.
1410. Kim, S., Y. Kang, et al. Points to Be Considered When Applying FibroScan S Probe in Children With Biliary Atresia. *Journal of Pediatric Gastroenterology and Nutrition* 2014;59(5):624-8.
1411. Karlas, T., D. Petroff, et al. Non-invasive assessment of hepatic steatosis in patients with NAFLD using controlled attenuation parameter and 1H-MR spectroscopy. *PLoS ONE* 2014;9(3):e91987.
1412. Karlas, T., J. Weber, et al. Value of liver elastography and abdominal ultrasound for detection of complications of allogeneic hemopoietic SCT. *Bone Marrow Transplantation* 2014;49(6):806-11.
1413. Khanna, R. and S. K. Sarin. Non-cirrhotic portal hypertension - diagnosis and management. *Journal of Hepatology* 2014;60(2):421-41.
1414. Kao, J. H. Risk stratification of HBV infection in Asia-Pacific region. *Clinical and Molecular Hepatology* 2014;20(3):223-227.
1415. Jerkeman, A., J. Westin, et al. Chronic hepatitis C in Swedish subjects receiving opiate substitution therapy--factors associated with advanced fibrosis. *Scandinavian Journal of Infectious Diseases* 2014;46(5):340-7.
1416. Kang, W., S. U. Kim, et al. Non-invasive prediction of forthcoming cirrhosis-related complications. *World Journal of Hepatology* 2014;20(10):2613-23.
1417. Jalli, R., M. Kazem Hosseini-Asl, et al. Ultrasound based techniques and transient elastography may not be precise methods for the detection of esophageal varices in liver cirrhosis. *Medical Ultrasonography Journal* 2014;16(1):78-9.
1418. Jeong, W. K., H. K. Lim, et al. Principles and clinical application of ultrasound elastography for diffuse liver disease. *Ultrasonography* 2014;33(3):149-60.
1419. Hu, X., X. Xu, et al. Indirect prediction of liver fibrosis by quantitative measurement of spleen stiffness using the FibroScan system. *Journal of ultrasound in medicine* 2014;33(1):73-81.
1420. Jang, W., S. I. Yu, et al. Longitudinal change of liver stiffness by transient elastography in chronic hepatitis B patients treated with nucleos(t)ide analogue. *Clinics and Research in Hepatology and Gastroenterology* 2014;38(2):195-200.
1421. Hong, M. Z., R. M. Zhang, et al. Liver stiffness measurement-based scoring system for significant inflammation related to chronic hepatitis B. *PLoS ONE* 2014;9(10):e111641.
1422. Ji, D., Q. Shao, et al. The frequency and determinants of liver stiffness measurement failure: a retrospective study of "real-life" 38,464 examinations. *PLoS ONE* 2014;9(8):e105183.

1423. Jung, K. S., B. K. Kim, et al. Factors Affecting the Accuracy of Controlled Attenuation Parameter (CAP) in Assessing Hepatic Steatosis in Patients with Chronic Liver Disease. *PLoS ONE* 2014;9(6):e98689.
1424. Jung, K. S., J. H. Kim, et al. Liver stiffness value-based risk estimation of late recurrence after curative resection of hepatocellular carcinoma: development and validation of a predictive model. *PLoS ONE* 2014;9(6):e99167.
1425. Herrero, J. I., M. Inarrairaegui, et al. [Comparison of the M and XL FibroScan((R)) probes to estimate liver stiffness by transient elastography]. *Gastroenterología y hepatología* 2014;37(4):233-9.
1426. Enomoto, M., H. Morikawa, et al. Noninvasive assessment of liver fibrosis in patients with chronic hepatitis B. *World Journal of Gastroenterology* 2014;20(34):12031-8.
1427. Ferraioli, G., C. Tinelli, et al. Interobserver reproducibility of the controlled attenuation parameter (CAP) for quantifying liver steatosis. *Hepatology International* 2014;8(4):576-81.
1428. Elkrief, L., P. E. Rautou, et al. Prospective Comparison of Spleen and Liver Stiffness by Using Shear-Wave and Transient Elastography for Detection of Portal Hypertension in Cirrhosis. *Radiology* 2014;141210.
1429. Eslamparast, T., H. Poustchi, et al. Synbiotic supplementation in nonalcoholic fatty liver disease: a randomized, double-blind, placebo-controlled pilot study. *American Journal of Clinical Nutrition* 2014;99(3):535-42.
1430. Gentile, I., A. R. Buonomo, et al. Is it possible to predict HCV-related liver cirrhosis non-invasively through routine laboratory parameters? *Le Infezioni in Medicina* 2014;22(1):11-8.
1431. Gervain, J. [The diagnosis of hepatitis C viral infection]. *Orvosi Hetilap* 2014;155(26):1019-23.
1432. Ferraioli, G., C. Tinelli, et al. Controlled attenuation parameter for evaluating liver steatosis in chronic viral hepatitis. *World Journal of Gastroenterology* 2014;20(21):6626-31.
1433. Expert Panel on Liver Stiffness, M. Clinical Application of Transient Elastography in the Diagnosis of Liver Fibrosis: an Expert Panel Review and Opinion. *Journal of Clinical and Translational Hepatology* 2014;2(2):110-6.
1434. Feuth, T., D. van Baarle, et al. CD4/CD8 ratio is a promising candidate for non-invasive measurement of liver fibrosis in chronic HCV-monoinfected patients. *European Journal of Clinical Microbiology & Infectious Diseases* 2014;33(7):1113-7.
1435. Garcia-Samaniego, J. and M. Romero. [The value of noninvasive serological markers in hepatitis B]. *Gastroenterología y hepatología* 2014;37 Suppl 2:22-5.
1436. Fransen van de Putte, D. E., M. Makris, et al. Long-term follow-up of hepatitis C infection in a large cohort of patients with inherited bleeding disorders. *Journal of Hepatology* 2014;60(1):39-45.
1437. Gatserelia, L., L. Shar vadze, et al. Occurrence of occult HCV infection among HIV infected patients in Georgia. *Georgian Medical News* 2014;-(226):37-41.
1438. Fraquelli, M., M. Giunta, et al. Feasibility and reproducibility of spleen transient elastography and its role in combination with liver transient elastography for predicting the severity of chronic viral hepatitis. *Journal of viral hepatitis* 2014;21(2):90-8.
1439. Hamidieh, A. A., B. Shazad, et al. Noninvasive measurement of liver fibrosis using transient elastography in pediatric patients with major thalassemia who are candidates for hematopoietic stem cell transplantation. *Biology of Blood and Marrow Transplantation* 2014;20(12):1912-7.
1440. Harkisoen, S., J. E. Arends, et al. Historic and current hepatitis B viral DNA and quantitative HBsAg level are not associated with cirrhosis in non-Asian women with chronic hepatitis B. *International Journal of Infectious Diseases* 2014;29:133-8.
1441. Harkisoen, S., G. J. Boland, et al. ELF-test less accurately identifies liver cirrhosis diagnosed by liver stiffness measurement in non-Asian women with chronic hepatitis B. *Journal of Clinical Virology* 2014;61(4):503-8.
1442. Fitzpatrick, E. and A. Dhawan. Noninvasive biomarkers in non-alcoholic fatty liver disease: current status and a glimpse of the future. *World Journal of Gastroenterology* 2014;20(31):10851-63.
1443. Eslamparast, T., F. Zamani, et al. Effects of synbiotic supplementation on insulin resistance in subjects with the metabolic syndrome: a randomised, double-blind, placebo-controlled pilot study. *British Journal of Nutrition* 2014;112(3):438-45.
1444. Gerstenmaier, J. F. and R. N. Gibson. Ultrasound in chronic liver disease. *Insights into Imaging* 2014;5(4):441-55.
1445. Goldschmidt, I., C. Brauch, et al. Spleen stiffness measurement by transient elastography to diagnose portal hypertension in children. *Journal of Pediatric Gastroenterology and Nutrition* 2014;59(2):197-203.
1446. Hassan, S., S. Syed, et al. Review of diagnostic techniques of hepatic fibrosis. *Journal of Pakistan Medical Association* 2014;64(8):941-5.
1447. Ferraioli, G., C. Tinelli, et al. Point shear wave elastography method for assessing liver stiffness. *World Journal of Gastroenterology* 2014;20(16):4787-96.
1448. Fernandez-Montero, J. V., P. Barreiro, et al. Liver fibrosis progression in HIV-HCV-coinfected patients treated with distinct antiretroviral drugs

- and impact of pegylated interferon/ribavirin therapy. *Antiviral Therapy* 2014;19(3):287-92.
1449. Frulio, N., H. Trillaud, et al. Acoustic Radiation Force Impulse (ARFI) and Transient Elastography (TE) for evaluation of liver fibrosis in HIV-HCV co-infected patients. *BMC Infectious Diseases* 2014;14(1):405.
1450. de Ledinghen, V., J. Vergniol, et al. Controlled attenuation parameter (CAP) for the diagnosis of steatosis: a prospective study of 5323 examinations. *Journal of Hepatology* 2014;60(5):1026-31.
1451. De Robertis, R., M. D'Onofrio, et al. Noninvasive diagnosis of cirrhosis: a review of different imaging modalities. *World Journal of Gastroenterology* 2014;20(23):7231-41.
1452. Colecchia, A., R. Schiumerini, et al. Prognostic factors for hepatocellular carcinoma recurrence. *World Journal of Gastroenterology* 2014;20(20):5935-50.
1453. Colecchia, A., A. Colli, et al. Spleen stiffness measurement can predict clinical complications in compensated HCV-related cirrhosis: a prospective study. *Journal of Hepatology* 2014;60(6):1158-64.
1454. Coccilillo, S., G. Parruti, et al. CEUS and Fibroscan in non-alcoholic fatty liver disease and non-alcoholic steatohepatitis. *World Journal of Hepatology* 2014;6(7):496-503.
1455. Crespo, G., S. Lens, et al. Liver stiffness 1 year after transplantation predicts clinical outcomes in patients with recurrent hepatitis C. *American Journal of Transplantation* 2014;14(2):375-83.
1456. Corpechot, C., F. Gaouar, et al. Baseline values and changes in liver stiffness measured by transient elastography are associated with severity of fibrosis and outcomes of patients with primary sclerosing cholangitis. *Gastroenterology* 2014;146(4):970-9; quiz e15-6.
1457. Christiansen, K. M., B. K. Mossner, et al. Liver stiffness measurement among patients with chronic hepatitis B and C: results from a 5-year prospective study. *PLoS ONE* 2014;9(11):e111912.
1458. Chon, Y. E., S. U. Kim, et al. Dynamics of the liver stiffness value using transient elastography during the perioperative period in patients with valvular heart disease. *PLoS ONE* 2014;9(3):e92795.
1459. Cassinero, E., N. Orofino, et al. Combination of deferasirox and deferoxamine in clinical practice: an alternative scheme of chelation in thalassemia major patients. *Blood Cells, Molecules & Diseases* 2014;53(3):164-7.
1460. Cassinotto, C., B. Lapuyade, et al. Non-invasive assessment of liver fibrosis with impulse elastography: comparison of Supersonic Shear Imaging with ARFI and FibroScan(R). *Journal of Hepatology* 2014;61(3):550-7.
1461. Cequera, A. and M. C. Garcia de Leon Mendez. [Biomarkers for liver fibrosis: advances, advantages and disadvantages]. *Revista de Gastroenterologia de Mexico* 2014;79(3):187-99.
1462. Castellano, G. and M. L. Manzano. [The value of FibroScan(R) in the follow-up of patients with chronic hepatitis B virus infection without indication for treatment]. *Gastroenterología y Hepatología* 2014;37 Suppl 2:15-21.
1463. Chan, W. K., N. R. Nik Mustapha, et al. Controlled attenuation parameter for the detection and quantification of hepatic steatosis in nonalcoholic fatty liver disease. *Journal of Gastroenterology & Hepatology* 2014;29(7):1470-6.
1464. Castera, L., M. Winnock, et al. Comparison of transient elastography (FibroScan), FibroTest, APRI and two algorithms combining these non-invasive tests for liver fibrosis staging in HIV/HCV coinfect ed patients: ANRS CO13 HEPAVIH and FIBROSTIC collaboration. *HIV Medicine* 2014;15(1):30-9.
1465. Chon, Y. E., K. S. Jung, et al. Controlled attenuation parameter (CAP) for detection of hepatic steatosis in patients with chronic liver diseases: a prospective study of a native Korean population. *Liver International* 2014;34(1):102-9.
1466. Bonnet, D., M. Guivarch, et al. Telaprevir- and boceprevir-based tritherapies in real practice for F3-F4 pretreated hepatitis C virus patients. *World Journal of Hepatology* 2014;6(9):660-9.
1467. Azmi, A. N., S. S. Tan, et al. Practical approach in hepatitis B e antigen-negative individuals to identify treatment candidates. *World Journal of Gastroenterology* 2014;20(34):12045-55.
1468. Avihingsanon, A., S. Jitmitraparp, et al. Advanced liver fibrosis by transient elastography, fibrosis 4, and alanine aminotransferase/platelet ratio index among Asian hepatitis C with and without human immunodeficiency virus infection: role of vitamin D levels. *Journal of Gastroenterology & Hepatology* 2014;29(9):1706-14.
1469. Bouscaillou, J., J. Champagnat, et al. Hepatitis C among people who inject drugs in Tbilisi, Georgia: an urgent need for prevention and treatment. *International Journal of Drug Policy* 2014;25(5):871-8.
1470. Branchi, F., C. B. Conti, et al. Non-invasive assessment of liver fibrosis in chronic hepatitis B. *World Journal of Gastroenterology* 2014;20(40):14568-14580.
1471. Aykut, U. E., U. Akyuz, et al. A comparison of FibroMeter NAFLD Score, NAFLD fibrosis score, and transient elastography as noninvasive diagnostic tools for hepatic fibrosis in patients with biopsy-proven non-alcoholic fatty liver disease. *Scandinavian Journal of Gastroenterology* 2014;49(11):1343-8.
1472. Barr, R. G. Elastography in clinical practice. *Radiologic Clinics of North America* 2014;52(6):1145-62.

1473. Berzigotti, A. Getting closer to a point-of-care diagnostic assessment in patients with chronic liver disease: controlled attenuation parameter for steatosis. *Journal of Hepatology* 2014;60(5):910-2.
1474. Bohte, A. E., A. de Niet, et al. Non-invasive evaluation of liver fibrosis: a comparison of ultrasound-based transient elastography and MR elastography in patients with viral hepatitis B and C. *European Radiology* 2014;24(3):638-48.
1475. Bota, S. and M. Peck-Radosavljevic. Non-invasive evaluation of patients with viral hepatitis. *Minerva Gastroenterologica e Dietologica* 2014;60(1):39-54.
1476. Boursier, J., V. de Ledinghen, et al. Precise evaluation of liver histology by computerized morphometry shows that steatosis influences liver stiffness measured by transient elastography in chronic hepatitis C. *Journal of Gastroenterology* 2014;49(3):527-37.
1477. Buck, M., G. Garcia-Tsao, et al. Novel inflammatory biomarkers of portal pressure in compensated cirrhosis patients. *Hepatology* 2014;59(3):1052-9.
1478. Cales, P., J. Boursier, et al. Improved fibrosis staging by elastometry and blood test in chronic hepatitis C. *Liver International* 2014;34(6):907-17.
1479. Carvalhana, S., J. Leitao, et al. How good is controlled attenuation parameter and fatty liver index for assessing liver steatosis in general population: correlation with ultrasound. *Liver International* 2014;34(6):e111-7.
1480. Bonder, A. and N. Afdhal. Utilization of FibroScan in Clinical Practice. *Current Gastroenterology Reports* 2014;16(2):372.
1481. Berzigotti, A. Non-Invasive Assessment of Non-Alcoholic Fatty Liver Disease: Ultrasound and Transient Elastography. *Reviews on recent clinical trials* 2014;9(3):170-177.
1482. Bessone, F. Re-appraisal of old and new diagnostic tools in the current management of chronic hepatitis B. *Liver International* 2014;34(7):991-1000.
1483. Brescini, L., E. Orsetti, et al. Evaluating Liver Fibrosis by Transient Elastometry in Patients With HIV-HCV Coinfection and Monoinfection. *Hepatitis Monthly* 2014;14(8):e15426.
1484. Avihingsanon, A., T. Apornpong, et al. Decline in serum 25 hydroxyvitamin D levels in HIV-HBV-coinfected patients after long-term antiretroviral therapy. *Antiviral Therapy* 2014;19(1):41-9.
1485. Assal, A., D. Mackie, et al. Transient elastography, APRI, and ultrasound have minimal utility in chronic low-replicative hepatitis B infection. *European Journal of Gastroenterology & Hepatology* 2014;26(9):1010-4.
1486. Augustin, S., L. Millan, et al. Detection of early portal hypertension with routine data and liver stiffness in patients with asymptomatic liver disease: A prospective study. *Journal of Hepatology* 2014;60(3):561-9.
1487. Agrawal, S. and A. Duseja. Nonalcoholic Fatty Liver Disease--The Clinician's Perspective. *Tropical Gastroenterology* 2014;35(4):212-21.
1488. Abdennour, M., S. Reggio, et al. Association of Adipose Tissue and Liver Fibrosis with Tissue Stiffness in Morbid Obesity: Links with Diabetes and BMI Loss after Gastric Bypass. *The Journal of Clinical Endocrinology and Metabolism* 2014;99(3):898-907.
1489. Ahn, J. M., Y. H. Paik, et al. Relationship of liver stiffness and controlled attenuation parameter measured by transient elastography with diabetes mellitus in patients with chronic liver disease. *Journal of Korean medical science* 2014;29(8):1113-9.
- 2013**
1490. Zarski, J. P., N. Sturm, et al. Contribution of the ELFG test in algorithms of non-invasive markers towards the diagnosis of significant fibrosis in chronic hepatitis C. *PLoS ONE* 2013;8(3):e59088.
1491. Yoshioka, K. How to adjust the inflammation-induced overestimation of liver fibrosis using transient elastography? *Hepatology Research* 2013;43(2):182-4.
1492. Yoneda, M. Utility of controlled attenuation parameter measurement for assessing liver steatosis in Japanese patients with chronic liver diseases. *Hepatology Research* 2013;43(11):1125-6.
1493. Wong, V. W., R. S. Chan, et al. Community-based lifestyle modification programme for non-alcoholic fatty liver disease: a randomized controlled trial. *Journal of Hepatology* 2013;59(3):536-42.
1494. Yilmaz, Y. Biomarkers for early detection of non-alcoholic steatohepatitis: implications for drug development and clinical trials. *Current Drug Targets* 2013;14(11):1357-66.
1495. Wong, G. L., H. L. Chan, et al. Liver fibrosis progression in chronic hepatitis B patients positive for hepatitis B e antigen: a prospective cohort study with paired transient elastography examination. *Journal of gastroenterology and hepatology* 2013;28(11):1762-9.
1496. Yilmaz, Y., R. Ergelen, et al. Noninvasive detection of hepatic steatosis in patients without ultrasonographic evidence of fatty liver using the controlled attenuation parameter evaluated with transient elastography. *European Journal of Gastroenterology & Hepatology* 2013;25(11):1330-4.
1497. Wong, G. L. Transient elastography: Kill two birds with one stone? *World Journal of Hepatology* 2013;5(5):264-74.

1498. Wang, H. M., C. H. Hung, et al. Liver stiffness measurement as an alternative to fibrotic stage in risk assessment of hepatocellular carcinoma incidence for chronic hepatitis C patients. *Liver International* 2013;33(5):756-61.
1499. von Volkman, H. L., R. F. Havre, et al. Quantitative measurement of ultrasound attenuation and Hepato-Renal Index in Non-Alcoholic Fatty Liver Disease. *Medical Ultrasonography Journal* 2013;15(1):16-22.
1500. Wong, G. L., H. L. Chan, et al. Association between anthropometric parameters and measurements of liver stiffness by transient elastography. *Clinical Gastroenterology & Hepatology* 2013;11(3):295-302 e1-3.
1501. Wong, G. L., J. Vergniol, et al. Non-invasive assessment of liver fibrosis with transient elastography (FibroScan(R)): applying the cut-offs of M probe to XL probe. *Annals of Hepatology* 2013;12(4):570-80.
1502. Wong, G. L. Update of liver fibrosis and steatosis with transient elastography (Fibroscan). *Gastroenterology Report* 2013;1(1):19-26.
1503. Wong, V. W., G. L. Wong, et al. Liver stiffness measurement: simplicity is prerequisite for reliability. *Gastroenterology* 2013;144(1):243-4.
1504. Wiegand, J. and T. Berg. The etiology, diagnosis and prevention of liver cirrhosis: part 1 of a series on liver cirrhosis. *Deutsches Ärzteblatt international* 2013;110(6):85-91.
1505. Yan, L. B., X. Zhu, et al. Impact of mild to moderate elevations of alanine aminotransferase on liver stiffness measurement in chronic hepatitis B patients during antiviral therapy. *Hepatology Research* 2013;43(2):185-91.
1506. Wei, H., B. Li, et al. Serum GP73, a Marker for Evaluating Progression in Patients with Chronic HBV Infections. *PLoS ONE* 2013;8(2):e53862.
1507. Ydreborg, M., J. Westin, et al. Impact of IL28b-related single nucleotide polymorphisms on liver transient elastography in chronic hepatitis C infection. *PLoS ONE* 2013;8(11):e80172.
1508. Suzuki, K., M. Yoneda, et al. Transient elastography for monitoring the fibrosis of non-alcoholic fatty liver disease for 4 years. *Hepatology Research* 2013;43(9):979-83.
1509. Sporea, I., S. Bota, et al. Acoustic Radiation Force Impulse and Supersonic Shear Imaging Versus Transient Elastography for Liver Fibrosis Assessment. *Ultrasound in Medicine and Biology* 2013;39(11):1933-41.
1510. Sherman, D. I., M. Jagtiani, et al. Noninvasive predictors of portal hypertension. *Gastroenterology* 2013;144(5):1150-1.
1511. Sommerer, C., M. Scharf, et al. Assessment of renal allograft fibrosis by transient elastography. *Transplant International* 2013;26(5):545-51.
1512. Steadman, R., R. P. Myers, et al. A health technology assessment of transient elastography in adult liver disease. *Canadian Journal of Gastroenterology* 2013;27(3):149-58.
1513. Sporea, I., I. Ratiu, et al. Are different cut-off values of liver stiffness assessed by transient elastography according to the etiology of liver cirrhosis for predicting significant esophageal varices? *Medical Ultrasonography Journal* 2013;15(2):111-5.
1514. Sun, J., X. Liang, et al. Commentary: prognostication of chronic hepatitis B--are Fibrotest and Fibroscan the final answers? *Alimentary Pharmacology & Therapeutics* 2013;37(11):1113.
1515. Tomono, W., M. Yoneda, et al. Evaluation of the Liver Fibrosis Index calculated by using real-time tissue elastography for the non-invasive assessment of liver fibrosis in chronic liver diseases. *Hepatology Research* 2013;43(7):735-42.
1516. Shi, K. Q., Y. C. Fan, et al. Transient elastography: a meta-analysis of diagnostic accuracy in evaluation of portal hypertension in chronic liver disease. *Clinical Gastroenterology & Hepatology* 2013;33(1):62-71.
1517. Theile, D., W. E. Haefeli, et al. Association of liver stiffness with hepatic expression of pharmacokinetically important genes in alcoholic liver disease. *Alcoholism, clinical and experimental research* 2013;37 Suppl 1:E17-22.
1518. Singh, S., L. L. Fujii, et al. Liver stiffness is associated with risk of decompensation, liver cancer, and death in patients with chronic liver diseases: a systematic review and meta-analysis. *Clinical Gastroenterology & Hepatology* 2013;11(12):1573-84 e1-2; quiz e88-9.
1519. Sporea, I., O. H. Gilja, et al. Liver elastography - an update. *Medical Ultrasonography Journal* 2013;15(4):304-14.
1520. Sirli, R., I. Sporea, et al. Factors influencing reliability of liver stiffness measurements using transient elastography (M-probe)-monocentric experience. *European Journal of Radiology* 2013;82(8):e313-6.
1521. Tangpradabkiet, W., N. Chamroonkul, et al. Influence of Food Intake on Measurement of Liver Stiffness by Transient Elastography in Patients with Chronic Viral Hepatitis B and C. *Thai Journal of Gastroenterology* 2013;14(1):16-22.
1522. Ricci, P., C. Marigliano, et al. Ultrasound evaluation of liver fibrosis: preliminary experience with acoustic structure quantification (ASQ) software. *Radiologia Medica* 2013;118(6):995-1010.
1523. Saad, Y., M. Said, et al. Liver stiffness measurement by fibroscan predicts the presence and size of esophageal varices in egyptian patients with HCV related liver cirrhosis. *Journal of Clinical and Diagnostic Research* 2013;7(10):2253-7.

1524. Saiz de la Hoya Zamacola, P., C. Alia Alia, et al. [The progression of liver fibrosis in prison inmates co-infected by HIV and HCV who started on boosted protease inhibitor therapy]. *Revista Española de Sanidad Penitenciaria* 2013;15(2):54-62.
1525. Rudler, M., T. Poynard, et al. Liver stiffness, platelets, and spleen size is reliable in nondecompensated cirrhotic patients. *Gastroenterology* 2013;144(5):1150.
1526. Schieren, G., E. Bolke, et al. Severe everolimus-induced steatohepatitis: a case report. *European Journal of Medical Research* 2013;18(1):22.
1527. Sharma, P., V. Kirnake, et al. Spleen stiffness in patients with cirrhosis in predicting esophageal varices. *American Journal of Gastroenterology* 2013;108(7):1101-7.
1528. Ruiz-Artacho, P., M. J. Tellez-Molina, et al. HIV/HBV coinfection: serological control and therapeutic strategies. *Revista clinica espanola* 2013;213(6):285-90.
1529. Risum, M., T. S. Barfod, et al. [Noninvasive estimation of fibrosis in chronic viral hepatitis]. *Ugeskrift for Laeger* 2013;175(34):1877-82.
1530. Roitberg, G. E., L. B. Lazebnik, et al. [Non-invasive techniques in diagnosis of steatosis and fibrosis in nonalcoholic fatty liver disease]. *Eksperimental Klinical Gastroenterology* 2013;2013(7):3-9.
1531. Poynard, T., J. Moussalli, et al. Slow regression of liver fibrosis presumed by repeated biomarkers after virological cure in patients with chronic hepatitis C. *Journal of Hepatology* 2013;59(4):675-83.
1532. Poynard, T., M. Munteanu, et al. Liver fibrosis evaluation using real-time shear wave elastography: applicability and diagnostic performance using methods without a gold standard. *Journal of Hepatology* 2013;58(5):928-35.
1533. Pournik, O., S. M. Alavian, et al. Can we accurately assess liver fibrosis with FibroScan(R) using fewer valid measurements? *Yonsei Medical Journal* 2013;54(2):541-2.
1534. Procopet, B., M. Tantau, et al. Are there any alternative methods to hepatic venous pressure gradient in portal hypertension assessment? *Journal of Gastrointestinal & Liver Diseases* 2013;22(1):73-8.
1535. Pradhan, F., F. Ladak, et al. Feasibility and reliability of the FibroScan S2 (pediatric) probe compared with the M probe for liver stiffness measurement in small adults with chronic liver disease. *Annals of Hepatology* 2013;12(1):100-7.
1536. Poustchi, H., M. Eslami, et al. Transient elastography in hepatitis C virus-infected patients with beta-thalassemia for assessment of fibrosis. *Hepatology Research* 2013;43(12):1276-83.
1537. Potthoff, A., D. Attia, et al. Influence of different frequencies and insertion depths on the diagnostic accuracy of liver elastography by acoustic radiation force impulse imaging (ARFI). *European Journal of Radiology* 2013;82(8):1207-12.
1538. Rocio, E., C. Cifuentes, et al. Interobserver concordance in controlled attenuation parameter measurement, a novel tool for the assessment of hepatic steatosis on the basis of transient elastography. *European Journal of Gastroenterology & Hepatology* 2013;25(8):905-11.
1539. Peters, L., A. Mocroft, et al. Hyaluronic acid levels predict risk of hepatic encephalopathy and liver-related death in HIV/viral hepatitis coinfecting patients. *PLoS ONE* 2013;8(5):e64283.
1540. Park, Y., S. U. Kim, et al. Compensated heart failure does not always alleviate overestimation of liver stiffness by transient elastography. *European Journal of Gastroenterology & Hepatology* 2013;25(3):385-7.
1541. Mikolasevic, I., S. Racki, et al. Chronic Kidney Disease and Nonalcoholic Fatty Liver Disease Proven by Transient Elastography. *Kidney & blood pressure research* 2013;37(4-5):305-310.
1542. Nascimbeni, F., R. Pais, et al. From NAFLD in clinical practice to answers from guidelines. *Journal of Hepatology* 2013;59(4):859-71.
1543. Macias, J., M. Marquez, et al. Risk of liver decompensation among HIV/hepatitis C virus-coinfected individuals with advanced fibrosis: implications for the timing of therapy. *Clinical Infectious Diseases* 2013;57(10):1401-8.
1544. Mahadeva, S., A. S. Mahfudz, et al. Performance of transient elastography (TE) and factors associated with discordance in non-alcoholic fatty liver disease. *Journal of Digestive Diseases* 2013;14(11):604-610.
1545. Masaki, K., S. Takaki, et al. Utility of controlled attenuation parameter measurement for assessing liver steatosis in Japanese patients with chronic liver diseases. *Hepatology Research* 2013;43(11):1182-1189.
1546. Northcott, M. J., W. L. Ong, et al. Prevalence of transfusion-acquired hepatitis C in an Australian bleeding disorders population. *Haemophilia* 2013;19(6):847-52.
1547. Morisco, F., R. Granata, et al. Sustained virological response: a milestone in the treatment of chronic hepatitis C. *World Journal of Gastroenterology* 2013;19(18):2793-8.
1548. Nalankilli, K. and J. Lubel. Elevated serum aminotransferases secondary to rippling muscle disease. *Case Reports in Gastroenterology* 2013;7(2):234-9.
1549. Margariti, A., M. Deutsch, et al. The severity of histologic liver lesions is independent of body mass index in patients with nonalcoholic fatty liver disease. *Journal of Clinical Gastroenterology* 2013;47(3):280-6.

1550. Mak, T. M., Y. P. Huang, et al. Liver fibrosis assessment using transient elastography guided with real-time B-mode ultrasound imaging: a feasibility study. *Ultrasound in Medicine and Biology* 2013;39(6):956-66.
1551. Par, G., G. Horvath, et al. [Non-alcoholic fatty liver disease and steatohepatitis]. *Orvosi Hetilap* 2013;154(29):1124-34.
1552. Moon, K. M., G. Kim, et al. Ultrasonographic scoring system score versus liver stiffness measurement in prediction of cirrhosis. *Clinical and Molecular Hepatology* 2013;19(4):389-98.
1553. Paparo, F., L. Cevasco, et al. Diagnostic value of real-time elastography in the assessment of hepatic fibrosis in patients with liver iron overload. *European Journal of Radiology* 2013;82(12):e755-61.
1554. Park, C. H., S. H. Bae, et al. A pilot study of autologous CD34-depleted bone marrow mononuclear cell transplantation via the hepatic artery in five patients with liver failure. *Cytotherapy* 2013;15(12):1571-9.
1555. Macias, J., A. Rivero, et al. Sustained virological response to pegylated interferon plus ribavirin leads to normalization of liver stiffness in hepatitis C virus-infected patients. *Enfermedades Infecciosas y Microbiología Clínica* 2013;31(7):424-9.
1556. Mikolasevic, I., L. Orlic, et al. Nonalcoholic Fatty Liver Disease (NAFLD) Proven By Transient Elastography In Patients With Coronary Heart Disease. *Wiener Klinische Wochenschrift* 2013;126(15-16):474-9.
1557. Macias, J., A. Camacho, et al. Liver stiffness measurement versus liver biopsy to predict survival and decompensations of cirrhosis among HIV/hepatitis C virus-coinfected patients. *Aids* 2013;27(16):2541-9.
1558. Lee, C. K., A. R. Perez-Atayde, et al. Serum biomarkers and transient elastography as predictors of advanced liver fibrosis in a United States cohort: the Boston children's hospital experience. *The Journal of pediatrics* 2013;163(4):1058-1064 e2.
1559. Lee, D. H., J. M. Lee, et al. MR elastography of healthy liver parenchyma: Normal value and reliability of the liver stiffness value measurement. *Journal of Magnetic Resonance Imaging* 2013;38(5):1215-23.
1560. Liu, F., T. H. Li, et al. [Non-invasive assessment of portal hypertension in patients with liver cirrhosis using FibroScan transient elastography]. *Zhonghua Gan Zang Bing Za Zhi* 2013;21(11):840-4.
1561. Kirk, G. D., S. H. Mehta, et al. HIV, age, and the severity of hepatitis C virus-related liver disease: a cohort study. *Annals of Internal Medicine* 2013;158(9):658-66.
1562. Kitson, M. T., W. W. Kemp, et al. Utility of transient elastography in the non-invasive evaluation of cystic fibrosis liver disease. *Liver International* 2013;33(5):698-705.
1563. Koh, C., T. Heller, et al. Long-term outcome of chronic hepatitis C after sustained virological response to interferon-based therapy. *Alimentary Pharmacology & Therapeutics* 2013;37(9):887-94.
1564. Machado, M. V. and H. Cortez-Pinto. Non-invasive diagnosis of non-alcoholic fatty liver disease. A critical appraisal. *Journal of Hepatology* 2013;58(5):1007-19.
1565. Kim, B. K., J. Fung, et al. Clinical application of liver stiffness measurement using transient elastography in chronic liver disease from longitudinal perspectives. *World Journal of Gastroenterology* 2013;19(12):1890-900.
1566. Kellner, P., E. Anadol, et al. The effect of hemodialysis on liver stiffness measurement: a single-center series. *European Journal of Gastroenterology & Hepatology* 2013;25(3):368-72.
1567. Lupsor Platon, M., H. Stefanescu, et al. Performance of unidimensional transient elastography in staging chronic hepatitis C. Results from a cohort of 1,202 biopsied patients from one single center. *Journal of Gastrointestinal & Liver Diseases* 2013;22(2):157-66.
1568. Lens, S. and X. Forns. [Usefulness of liver stiffness measurement after antiviral response in chronic hepatitisC]. *Enfermedades Infecciosas y Microbiología Clínica* 2013;31(7):421-423.
1569. Kemp, W. and S. Roberts. FibroScan(R) and transient elastography. *Australian Family Physician* 2013;42(7):468-71.
1570. Kumar, M., A. Rastogi, et al. Controlled attenuation parameter for non-invasive assessment of hepatic steatosis: does etiology affect performance? *Journal of Gastroenterology & Hepatology* 2013;28(7):1194-201.
1571. Lee, J. Y., D. C. Lee, et al. Serum carcinoembryonic antigen is associated with non-alcoholic fatty liver disease in healthy Korean non-smokers. *Clinical Chemistry and Laboratory Medicine* 2013;51(7):1499-504.
1572. Kim, M. S. and J. R. Lee. Assessment of liver stiffness measurement: novel intraoperative blood loss predictor? *World Journal of Surgery* 2013;37(1):185-91.
1573. Kim, E., Y. Kang, et al. The efficacy of aspartate aminotransferase-to-platelet ratio index for assessing hepatic fibrosis in childhood nonalcoholic steatohepatitis for medical practice. *Korean Journal of Pediatrics* 2013;56(1):19-25.
1574. Kumar, R., A. Rastogi, et al. Liver stiffness measurements in patients with different stages of nonalcoholic fatty liver disease: diagnostic performance and clinicopathological correlation. *Digestive Diseases & Sciences* 2013;58(1):265-74.
1575. Lannerstedt, H., Z. Konopski, et al. Combining transient elastography with FIB4 enhances

- sensitivity in detecting advanced fibrosis of the liver. *Scandinavian Journal of Gastroenterology* 2013;48(1):93-100.
1576. Kim, B. K., H. J. Oh, et al. Early on-treatment change in liver stiffness predicts development of liver-related events in chronic hepatitis B patients receiving antiviral therapy. *Liver International* 2013;33(2):180-9.
1577. Kim, S. U., B. K. Kim, et al. Clinical Application of Liver Stiffness Measurement Using Transient Elastography: A Surgical Perspective. *Digestion* 2013;88(4):258-265.
1578. Karlas, T., M. Neuschulz, et al. ARFI and transient elastography for characterization of cystic fibrosis related liver disease: first longitudinal follow-up data in adult patients. *Journal of Cystic Fibrosis* 2013;12(6):826-7.
1579. Leung, V. Y., J. Shen, et al. Quantitative elastography of liver fibrosis and spleen stiffness in chronic hepatitis B carriers: comparison of shear-wave elastography and transient elastography with liver biopsy correlation. *Radiology* 2013;269(3):910-8.
1580. Leclerc, G. E., F. Charleux, et al. Analysis of liver viscosity behavior as a function of multifrequency magnetic resonance elastography (MMRE) postprocessing. *Journal of Magnetic Resonance Imaging* 2013;38(2):422-8.
1581. Kim, D. Y., K. J. Song, et al. Transient elastography-based risk estimation of hepatitis B virus-related occurrence of hepatocellular carcinoma: development and validation of a predictive model. *Journal of OncoTargets and Therapy* 2013;6:1463-9.
1582. Islam, M. M., A. Taylor, et al. General health of opioid substitution therapy clients. *Internal Medicine Journal* 2013;43(12):1335-8.
1583. Honsawek, S., W. Udomsinprasert, et al. Correlation of connective tissue growth factor with liver stiffness measured by transient elastography in biliary atresia. *Hepatology Research* 2013;43(7):795-800.
1584. Hong, W. K., M. Y. Kim, et al. The usefulness of non-invasive liver stiffness measurements in predicting clinically significant portal hypertension in cirrhotic patients: Korean data. *Clinical and Molecular Hepatology* 2013;19(4):370-5.
1585. Han, S. H., S. U. Kim, et al. Abnormal Liver Stiffness Assessed Using Transient Elastography (FibroScan(R)) in HIV-Infected Patients without HBV/HCV Coinfection Receiving Combined Antiretroviral Treatment. *PLoS ONE* 2013;8(1):e52720.
1586. Hahn, S. M., S. Kim, et al. Clinical benefit of liver stiffness measurement at 3 months after Kasai hepatoportoenterostomy to predict the liver related events in biliary atresia. *PLoS ONE* 2013;8(11):e80652.
1587. Esmat, G., A. Elsharkawy, et al. Fibroscan of chronic HCV patients coinfected with schistosomiasis. *Arab journal of gastroenterology* 2013;14(3):109-12.
1588. Feier, D., M. Lupsor Platon, et al. Transient elastography for the detection of hepatocellular carcinoma in viral C liver cirrhosis. Is there something else than increased liver stiffness? *Journal of Gastrointestinal & Liver Diseases* 2013;22(3):283-9.
1589. Feuth, T., J. E. Arends, et al. Impact of transient elastography on clinical decision-making in patients with chronic viral hepatitis. *Scandinavian Journal of Gastroenterology* 2013;48(9):1074-81.
1590. Friedrich-Rust, M., N. Schlueter, et al. Non-invasive measurement of liver and pancreas fibrosis in patients with cystic fibrosis. *Journal of Cystic Fibrosis* 2013;12(5):431-9.
1591. Goldschmidt, I., H. Stieghorst, et al. The use of transient elastography and non-invasive serum markers of fibrosis in pediatric liver transplant recipients. *Pediatric transplantation* 2013;17(6):525-34.
1592. Grgurevic, I., I. Hrstic, et al. [Assessment of liver disease severity in patients with chronic viral hepatitis]. *Acta Medica Croatica* 2013;67(4):291-301.
1593. Fabrellas, N., M. Alemany, et al. Using transient elastography to detect chronic liver diseases in a primary care nurse consultancy. *Nursing research* 2013;62(6):450-4.
1594. Goyal, R., S. R. Mallick, et al. Fibroscan can avoid liver biopsy in Indian patients with chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2013;28(11):1738-45.
1595. Frulio, N. and H. Trillaud. Ultrasound elastography in liver. *Diagnostic and Interventional Imaging* 2013;94(5):515-34.
1596. Gara, N., X. Zhao, et al. Discordance among transient elastography, aspartate aminotransferase to platelet ratio index, and histologic assessments of liver fibrosis in patients with chronic hepatitis C. *Clinical Gastroenterology & Hepatology* 2013;11(3):303-8 e1.
1597. Goldschmidt, I., C. Streckenbach, et al. Application and limitations of transient liver elastography in children. *Journal of Pediatric Gastroenterology and Nutrition* 2013;57(1):109-13.
1598. Ferraioli, G., C. Tinelli, et al. Performance of liver stiffness measurements by transient elastography in chronic hepatitis. *World Journal of Gastroenterology* 2013;19(1):49-56.
1599. Fitzpatrick, E., A. Quaglia, et al. Transient elastography is a useful noninvasive tool for the evaluation of fibrosis in paediatric chronic liver disease. *Journal of Pediatric Gastroenterology and Nutrition* 2013;56(1):72-6.
1600. Friedrich-Rust, M. and J. Vermehren. [Non-invasive methods for the evaluation of liver fibrosis

- in clinical practice]. Zeitschrift für Gastroenterologie 2013;51(1):43-54.
1601. Festi, D., R. Schiumerini, et al. Review article: the diagnosis of non-alcoholic fatty liver disease -- availability and accuracy of non-invasive methods. *Alimentary Pharmacology & Therapeutics* 2013;37(4):392-400.
1602. Frossard, J. L., E. Giostra, et al. The role of transient elastography in the detection of liver disease in patients with chronic pancreatitis. *Liver International* 2013;33(7):1121-7.
1603. Fernandez-Montero, J. V., P. Barreiro, et al. Liver stiffness predicts liver-related complications and mortality in HIV patients with chronic hepatitis C on antiretroviral therapy. *Aids* 2013;27(7):1129-34.
1604. Fung, J., C. K. Lee, et al. Defining Normal Liver Stiffness Range in a Normal Healthy Chinese Population without Liver Disease. *PLoS ONE* 2013;8(12):e85067.
1605. Barrault, C., F. Roudot-Thoraval, et al. Non-invasive assessment of liver graft fibrosis by transient elastography after liver transplantation. *Clinics and Research in Hepatology and Gastroenterology* 2013;37(4):347-52.
1606. Bota, S., H. Herkner, et al. Meta-analysis: ARFI elastography versus transient elastography for the evaluation of liver fibrosis. *Liver International* 2013;33(8):1138-47.
1607. Canavan, C., J. Eisenburg, et al. Ultrasound elastography for fibrosis surveillance is cost effective in patients with chronic hepatitis C virus in the UK. *Digestive Diseases & Sciences* 2013;58(9):2691-704.
1608. Cassinotto, C., B. Lapuyade, et al. Liver fibrosis: noninvasive assessment with acoustic radiation force impulse elastography - comparison with FibroScan M and XL probes and FibroTest in patients with chronic liver disease. *Radiology* 2013;269(1):283-92.
1609. Aljawad, M., E. M. Yoshida, et al. Percutaneous liver biopsy practice patterns among Canadian hepatologists. *Canadian Journal of Gastroenterology* 2013;27(11):e31-4.
1610. Castera, L., V. Vilgrain, et al. Noninvasive evaluation of NAFLD. *Nature reviews Gastroenterology & hepatology* 2013;10(11):666-75.
1611. Bensamoun, S. F., G. E. Leclerc, et al. Cutoff values for alcoholic liver fibrosis using magnetic resonance elastography technique. *Alcoholism, clinical and experimental research* 2013;37(5):811-7.
1612. Berenguer, M. and D. Schuppan. Progression of liver fibrosis in post-transplant hepatitis C: mechanisms, assessment and treatment. *Journal of Hepatology* 2013;58(5):1028-41.
1613. Berzigotti, A., J. C. Garcia-Pagan, et al. Reply. *Gastroenterology* 2013;144(5):1153-4.
1614. Calvaruso, V., V. Di Marco, et al. Fibrosis evaluation by transient elastography in patients with long-term sustained HCV clearance. *Hepatitis Monthly* 2013;13(5):e7176.
1615. Chin, J. L., G. Chan, et al. Spleen stiffness: the new kid on the block for diagnosing portal hypertension? *Gastroenterology* 2013;144(5):1152-3.
1616. de Ledinghen, V., J. Vergniol, et al. Non-invasive tests for fibrosis and liver stiffness predict 5-year survival of patients chronically infected with hepatitis B virus. *Alimentary Pharmacology & Therapeutics* 2013;37(10):979-88.
1617. de Mattos, A. Z. and A. A. de Mattos. Noninvasive methods for esophageal varices prediction in cirrhotic patients: are we there yet? *Gastroenterology* 2013;144(5):1151-2.
1618. Carbone, M., D. Mutimer, et al. Hepatitis C virus and nonliver solid organ transplantation. *Transplantation* 2013;95(6):779-86.
1619. Abraldes, J. G., E. Reverter, et al. Spleen stiffness: Towards a non-invasive portal sphygmomanometer? *Hepatology* 2013;57(3):1278-80.
1620. Bonder, A. and N. H. Afdhal. Biopsy no more; changing the screening and diagnostic algorithm for hepatitis C. *Clinical Gastroenterology & Hepatology* 2013;11(3):309-10.
1621. Boursier, J., J. P. Zarski, et al. Determination of reliability criteria for liver stiffness evaluation by transient elastography. *Hepatology* 2013;57(3):1182-91.
1622. Abd El Rihim, A. Y., R. F. Omar, et al. Role of fibroscan and APRI in detection of liver fibrosis: a systematic review and meta-analysis. *Arab journal of gastroenterology* 2013;14(2):44-50.
1623. Berenguer, J., J. Alvarez-Pellicer, et al. Clinical effects of viral relapse after interferon plus ribavirin in patients co-infected with human immunodeficiency virus and hepatitis C virus. *Journal of Hepatology* 2013;58(6):1104-12.
1624. Chen, Y. P., J. Peng, et al. Non-invasive assessment of liver fibrosis in patients with chronic hepatitis B. *Hepatology International* 2013;7(2):356-68.
1625. Chung, J. H., H. S. Ahn, et al. The usefulness of transient elastography, acoustic-radiation-force impulse elastography, and real-time elastography for the evaluation of liver fibrosis. *Clinical and Molecular Hepatology* 2013;19(2):156-64.
1626. de Ledinghen, V. and J. Vergniol. Commentary: prognostication of chronic hepatitis B - are Fibrotest and Fibroscan the final answers? Authors' reply. *Alimentary Pharmacology & Therapeutics* 2013;37(11):1114.
1627. Durango, E., C. Dietrich, et al. Direct comparison of the FibroScan XL and M probes for assessment of liver fibrosis in obese and nonobese

- patients. *Hepatic Medicine: Evidence and Research* 2013;5:43-52.
1628. Arena, U., M. Lupsor Platon, et al. Liver stiffness is influenced by a standardized meal in patients with chronic hepatitis C virus at different stages of fibrotic evolution. *Hepatology* 2013;58(1):65-72.
1629. Elalfy, M. S., G. Esmat, et al. Liver fibrosis in young Egyptian beta-thalassemia major patients: relation to hepatitis C virus and compliance with chelation. *Annals of Hepatology* 2013;12(1):54-61.
1630. Bardou-Jacquet, E., L. Legros, et al. Effect of alcohol consumption on liver stiffness measured by transient elastography. *World Journal of Gastroenterology* 2013;19(4):516-22.
1631. Bosselut, N., L. Taibi, et al. Including osteoprotegerin and collagen IV in a score-based blood test for liver fibrosis increases diagnostic accuracy. *Clinica Chimica Acta* 2013;415:63-8.
1632. Alkhouri, N., E. Sedki, et al. Combined paediatric NAFLD fibrosis index and transient elastography to predict clinically significant fibrosis in children with fatty liver disease. *Liver International* 2013;33(1):79-85.
1633. Berzigotti, A., S. Seijo, et al. Elastography, spleen size, and platelet count identify portal hypertension in patients with compensated cirrhosis. *Gastroenterology* 2013;144(1):102-111 e1.
1634. Bureau, C., V. Di Martino, et al. A major new step in non-invasive evaluation of portal hypertension: elastography. *Liver International* 2013;33(1):4-6.
1635. Castera, L. and G. Garcia-Tsao. When the spleen gets tough, the varices get going. *Gastroenterology* 2013;144(1):19-22.
1636. Carton, J. A., J. Collazos, et al. Course of liver fibrosis in HIV-hepatitis C virus-coinfected patients depending on the response to hepatitis C therapy. *AIDS Research and Human Retroviruses* 2013;29(2):215-22.
1637. Dolman, G. E., D. Nieboer, et al. The performance of transient elastography compared to clinical acumen and routine tests - what is the incremental diagnostic value? *Liver International* 2013;33(2):172-9.
1638. Bailony, M. R., R. Scherzer, et al. Association of HIV infection, hepatitis C virus infection, and metabolic factors with liver stiffness measured by transient elastography. *The Journal of Infectious Diseases* 2013;208(11):1776-83.
1639. Armstrong, M. J., C. Corbett, et al. Operator training requirements and diagnostic accuracy of FibroScan in routine clinical practice. *Postgraduate medical journal* 2013;89(1058):685-92.
1640. Awad Mel, D., G. E. Shiha, et al. Evaluation of liver stiffness measurement by FibroScan as compared to liver biopsy for assessment of hepatic fibrosis in children with chronic hepatitis C. *Journal of the Egyptian Society of Parasitology* 2013;43(3):805-19.
1641. Calvaruso, V., F. Bronte, et al. Modified spleen stiffness measurement by transient elastography is associated with presence of large oesophageal varices in patients with compensated hepatitis C virus cirrhosis. *Journal of viral hepatitis* 2013;20(12):867-74.
1642. Afdhal, N. H. FibroScan in the Diagnosis of Hepatitis C Virus Infection. *Gastroenterology & Hepatology* 2013;9(8):533-5.
1643. D'Ambrosio, R., A. Aghemo, et al. The diagnostic accuracy of Fibroscan for cirrhosis is influenced by liver morphometry in HCV patients with a sustained virological response. *Journal of Hepatology* 2013;59(2):251-6.
1644. Elalfy, M. S., A. A. Adly, et al. Effect of antioxidant therapy on hepatic fibrosis and liver iron concentrations in beta-thalassemia major patients. *Hemoglobin* 2013;37(3):257-76.
1645. Bartres, C. and S. Lens. [Elastography of the liver (FibroScan(R)) in hepatology]. *Revista Española de Enfermedades Digestivas* 2013;105(4):235.
1646. Berzigotti, A., A. De Gottardi, et al. Effect of meal ingestion on liver stiffness in patients with cirrhosis and portal hypertension. *PLoS ONE* 2013;8(3):e58742.
1647. Charpentier, C., K. Champenois, et al. Predictive value of liver enzymes and inflammatory biomarkers for the severity of liver fibrosis stage in HIV/HCV co-infected patients. *PLoS ONE* 2013;8(3):e59205.
1648. Chesnokova, L. V., I. M. Petrov, et al. [Proinflammatory cytokine levels depending on the stage of hepatic fibrosis in patients with metabolic syndrome and non-alcoholic fatty liver disease]. *Klinicheskaya Meditsina* 2013;91(12):30-5.

2012

1649. Sharma, P. and A. Kumar. Minimal hepatic encephalopathy in patients with cirrhosis by measuring liver stiffness and hepatic venous pressure gradient. *Saudi Journal of Gastroenterology* 2012;18(5):316-21.
1650. Rath, T., K. M. Menendez, et al. TIMP-1/-2 and transient elastography allow non invasive diagnosis of cystic fibrosis associated liver disease. *Digestive & Liver Disease* 2012;44(9):780-7.
1651. Sirli, R., I. Sporea, et al. Liver elastography for the diagnosis of portal hypertension in patients with liver cirrhosis. *Medical Ultrasonography Journal* 2012;14(3):225-30.
1652. Trifan, A., C. Cojocaru, et al. Liver biopsy or transient elastography? First, do not harm! *Journal of Gastrointestinal & Liver Diseases* 2012;21(3):325-6.

1653. Poynard, T., V. de Ledinghen, et al. Performances of Elasto-FibroTest®, a combination between FibroTest® and liver stiffness measurements for assessing the stage of liver fibrosis in patients with chronic hepatitis C. *Clinics and Research in Hepatology and Gastroenterology* 2012;36(5):455-463.
1654. Resino, S., M. Sanchez-Conde, et al. Coinfection by human immunodeficiency virus and hepatitis C virus: noninvasive assessment and staging of fibrosis. *Current Opinion in Infectious Diseases* 2012;25(5):564-9.
1655. Rizzo, L., G. Nunnari, et al. Acoustic Radial Force Impulse as an effective tool for a prompt and reliable diagnosis of hepatocellular carcinoma - preliminary data. *European Review for Medical and Pharmacological Sciences* 2012;16(11):1596-8.
1656. Roca, B., E. Resino, et al. Interobserver discrepancy in liver fibrosis using transient elastography. *Journal of viral hepatitis* 2012;19(10):711-5.
1657. Seijo, S., E. Reverter, et al. Role of hepatic vein catheterisation and transient elastography in the diagnosis of idiopathic portal hypertension. *Digestive & Liver Disease* 2012;44(10):855-60.
1658. Suarez-Zarracina, T., E. Valle-Garay, et al. Didanosine (ddl) associates with increased liver fibrosis in adult HIV-HCV coinfected patients. *Journal of viral hepatitis* 2012;19(10):685-93.
1659. Venturi, C., C. Sempoux, et al. Novel histologic scoring system for long-term allograft fibrosis after liver transplantation in children. *American Journal of Transplantation* 2012;12(11):2986-96.
1660. Tamano, M., K. Kojima, et al. The usefulness of measuring liver stiffness by transient elastography for assessing hepatic fibrosis in patients with various chronic liver diseases. *Hepatogastroenterology* 2012;59(115):826-30.
1661. Vermehren, J., A. Polta, et al. Comparison of acoustic radiation force impulse imaging with transient elastography for the detection of complications in patients with cirrhosis. *Liver International* 2012;32(5):852-8.
1662. Poynard, T., V. de Ledinghen, et al. Relative performances of FibroTest, Fibroscan and biopsy for assessing the stage of liver fibrosis in patients with chronic hepatitis C: a step toward the truth in the absence of a gold standard. *Journal of Hepatology* 2012;56(3):541-548.
1663. Wong, V. W., W. C. Chu, et al. Prevalence of non-alcoholic fatty liver disease and advanced fibrosis in Hong Kong Chinese: a population study using proton-magnetic resonance spectroscopy and transient elastography. *Gut* 2012;61(3):409-15.
1664. Yoshioka, K. and S. Hashimoto. Can non-invasive assessment of liver fibrosis replace liver biopsy? *Hepatology Research* 2012;42(3):233-40.
1665. Restellini, S. and L. Spahr. [Are non-invasive tests going to replace liver biopsy for diagnosis of liver fibrosis?]. *Revue Médicale Suisse* 2012;8(347):1411-5.
1666. Pesce, A., R. Scilletta, et al. Does transient elastography (FibroScan(R)) have a role in decision making in hepatocellular carcinoma? *HPB (Oxford)* 2012;14(6):403-8.
1667. Reiberger, T., A. Ferlitsch, et al. Noninvasive screening for liver fibrosis and portal hypertension by transient elastography--a large single center experience. *Wiener Klinische Wochenschrift* 2012;124(11-12):395-402.
1668. Sharma, P. and A. Arora. Is the M probe really necessary to measure liver stiffness by FibroScan? *Hepatology* 2012;55(6):2043-4; author reply 199-208.
1669. Sini, M., O. Sorbello, et al. Non-invasive assessment of hepatic fibrosis in a series of patients with Wilson's Disease. *Digestive & Liver Disease* 2012;44(6):487-91.
1670. Sporea, I. Is there a real future for liver elastography? *Journal of Gastrointestinal & Liver Diseases* 2012;21(2):129-31.
1671. Yoon, K. T., S. M. Lim, et al. Liver stiffness measurement using acoustic radiation force impulse (ARFI) elastography and effect of necroinflammation. *Digestive Diseases & Sciences* 2012;57(6):1682-91.
1672. Poynard, T., Y. Ngo, et al. Prognostic value of liver fibrosis biomarkers: a meta-analysis. *Gastroenterology & Hepatology* 2012;7(7):445-54.
1673. Wang, J. H., S. K. Chuah, et al. Transient elastography and simple blood markers in the diagnosis of esophageal varices for compensated patients with hepatitis B virus-related cirrhosis. *Journal of Gastroenterology & Hepatology* 2012;27(7):1213-8.
1674. Zoulim, F. and F. Bailly. New approaches to the management of hepatitis C in haemophilia in 2012. *Haemophilia* 2012;18 Suppl 4:28-33.
1675. Trifan, A., C. Cojocariu, et al. [Non-invasive evaluation of liver fibrosis in chronic hepatitis C]. *Revista medico-chirurgicala a Societății de Medici și Naturaliști din Iași* 2012;116(1):135-8.
1676. Zarski, J. P., N. Sturm, et al. Comparison of nine blood tests and transient elastography for liver fibrosis in chronic hepatitis C: the ANRS HCEP-23 study. *Journal of Hepatology* 2012;56(1):55-62.
1677. Vogel, M., E. Page, et al. Liver fibrosis progression after acute hepatitis C virus infection in HIV-positive individuals. *Clinical Infectious Diseases* 2012;54(4):556-9.
1678. Pineda, J. A., K. Neukam, et al. Hepatic safety of efavirenz in HIV/hepatitis C virus-coinfected patients with advanced liver fibrosis. *The Journal of Infection* 2012;64(2):204-11.

1679. Sasso, M., V. Miette, et al. The controlled attenuation parameter (CAP): A novel tool for the non-invasive evaluation of steatosis using Fibroscan((R)). *Clinics and Research in Hepatology and Gastroenterology* 2012;36(1):13-20.
1680. Sebastiani, G. and A. Alberti. Diagnostic algorithms for liver fibrosis in hepatitis C: are they ready to avoid liver biopsy? *Hepatology* 2012;55(2):652-3; author reply 653.
1681. Stevenson, M., M. Lloyd-Jones, et al. Non-invasive diagnostic assessment tools for the detection of liver fibrosis in patients with suspected alcohol-related liver disease: a systematic review and economic evaluation. *Health Technology Assessment* 2012;16(4):1-174.
1682. Pariente, A. [Management of non alcoholic fatty liver diseases in adults]. *La Revue du Praticien* 2012;62(10):1421-5.
1683. Ramirez, R., S. Valero, et al. A cross-sectional epidemiological study of chronic HCV liver disease stage in HIV-coinfected patients using noninvasive techniques: CoTrans- GREENI Study. *Current HIV Research* 2012;10(8):688-93.
1684. Sporea, I. and R. L. Sirli. Hepatic elastography for the assessment of liver fibrosis--present and future. *Ultraschall in der Medizin* 2012;33(6):550-8.
1685. Wong, V. W., J. Vergniol, et al. Liver stiffness measurement using XL probe in patients with nonalcoholic fatty liver disease. *American Journal of Gastroenterology* 2012;107(12):1862-71.
1686. Xu, H. W., S. N. Lu, et al. Liver stiffness measurement in cirrhotic patient -- implications of disease activity and treatment efficacy. *The Kaohsiung journal of medical sciences* 2012;28(12):641-8.
1687. Park, H., S. U. Kim, et al. Optimal time for restoring the reliability of liver stiffness measurement in patients with chronic hepatitis B experiencing acute exacerbation. *Journal of Clinical Gastroenterology* 2012;46(7):602-7.
1688. Sporea, I., R. Sirli, et al. Comparative study concerning the value of acoustic radiation force impulse elastography (ARFI) in comparison with transient elastography (TE) for the assessment of liver fibrosis in patients with chronic hepatitis B and C. *Ultrasound in Medicine and Biology* 2012;38(8):1310-6.
1689. Talwalkar, J. A. Chronic inflammation, liver stiffness, and clinical decision making: an unavoidable partnership. *Clinical Gastroenterology & Hepatology* 2012;10(8):938-40.
1690. Tapper, E. B., E. B. Cohen, et al. Levels of alanine aminotransferase confound use of transient elastography to diagnose fibrosis in patients with chronic hepatitis C virus infection. *Clinical Gastroenterology & Hepatology* 2012;10(8):932-937 e1.
1691. Torres, D. M., C. D. Williams, et al. Features, diagnosis, and treatment of nonalcoholic fatty liver disease. *Clinical Gastroenterology & Hepatology* 2012;10(8):837-58.
1692. Trabut, J. B., V. Thepot, et al. Rapid decline of liver stiffness following alcohol withdrawal in heavy drinkers. *Alcoholism, clinical and experimental research* 2012;36(8):1407-11.
1693. Wang, H. M., G. H. Lo, et al. Efficacy of transient elastography in screening for large esophageal varices in patients with suspicious or proven liver cirrhosis. *Journal of Digestive Diseases* 2012;13(8):430-8.
1694. Petta, S. and A. Craxi. Assessment by Fibroscan of fibrosis in nonalcoholic fatty liver disease: XL versus M probe? *Hepatology* 2012;55(4):1309; author reply 1309-10.
1695. Salmon, D., F. Bani-Sadr, et al. Insulin resistance is associated with a higher risk of hepatocellular carcinoma in cirrhotic HIV/HCV-co-infected patients: results from ANRS CO13 HEPAVIH. *Journal of Hepatology* 2012;56(4):862-8.
1696. Sasso, M., I. Tengher-Barna, et al. Novel controlled attenuation parameter for noninvasive assessment of steatosis using Fibroscan((R)): validation in chronic hepatitis C. *Journal of viral hepatitis* 2012;19(4):244-53.
1697. Sebagh, M., D. Samuel, et al. 20-year protocol liver biopsies: invasive but useful for the management of liver recipients. *Journal of Hepatology* 2012;56(4):840-7.
1698. Verveer, C., P. E. Zondervan, et al. Evaluation of transient elastography for fibrosis assessment compared with large biopsies in chronic hepatitis B and C. *Liver International* 2012;32(4):622-8.
1699. Reiberger, T., B. A. Payer, et al. A prospective evaluation of pulmonary, systemic and hepatic haemodynamics in HIV-HCV-coinfected patients before and after antiviral therapy with pegylated interferon and ribavirin. *Antiviral Therapy* 2012;17(7):1327-34.
1700. Rivero-Juarez, A., J. Morgaz, et al. Liver stiffness using transient elastography is applicable to canines for hepatic disease models. *PLoS ONE* 2012;7(7):e41557.
1701. Ruhwald, M., E. S. Andersen, et al. IP-10 can be measured in dried plasma spots in patients with chronic hepatitis C infection. *PLoS ONE* 2012;7(9):e45181.
1702. Schwarze-Zander, C., C. Boesecke, et al. Boceprevir in combination with HIV protease inhibitors in patients with advanced fibrosis-altered drug-drug interactions? *Journal of the International AIDS Society* 2012;15(6):18418.
1703. Wahl, K., W. Rosenberg, et al. Biopsy-controlled liver fibrosis staging using the enhanced liver fibrosis (ELF) score compared to transient elastography. *PLoS ONE* 2012;7(12):e51906.

1704. Montes Ramirez, M. L., J. F. Pascual-Pareja, et al. Transient elastography to rule out esophageal varices and portal hypertensive gastropathy in HIV-infected individuals with liver cirrhosis. *Aids* 2012;26(14):1807-12.
1705. Moreno, C. and P. Deltenre. [Management of chronic hepatitis C in 2012 in Belgium]. *Revue Médicale de Bruxelles* 2012;33(4):223-7.
1706. Masuzaki, R., R. Tateishi, et al. Assessment of disease progression in patients with transfusion-associated chronic hepatitis C using transient elastography. *World Journal of Gastroenterology* 2012;18(12):1385-90.
1707. Motosugi, U., T. Ichikawa, et al. Cross-validation of MR elastography and ultrasound transient elastography in liver stiffness measurement: discrepancy in the results of cirrhotic liver. *Journal of Magnetic Resonance Imaging* 2012;35(3):607-10.
1708. Myers, R. P., G. Pomier-Layrargues, et al. Discordance in fibrosis staging between liver biopsy and transient elastography using the FibroScan XL probe. *Journal of Hepatology* 2012;56(3):564-70.
1709. Myoung, S., J. H. Chang, et al. A mixture of experts model for the diagnosis of liver cirrhosis by measuring the liver stiffness. *Healthcare Informatics Research* 2012;18(1):29-34.
1710. Myers, R. P. Reply. *Hepatology* 2012;55(6):2044-2044.
1711. Massironi, S., R. E. Rossi, et al. Transient elastography in patients with celiac disease: a noninvasive method to detect liver involvement associated with celiac disease. *Scandinavian Journal of Gastroenterology* 2012;47(6):640-8.
1712. Merchante, N., A. Rivero-Juarez, et al. Liver stiffness predicts clinical outcome in human immunodeficiency virus/hepatitis C virus-coinfected patients with compensated liver cirrhosis. *Hepatology* 2012;56(1):228-38.
1713. Myers, R. P., A. Pollett, et al. Controlled Attenuation Parameter (CAP): a noninvasive method for the detection of hepatic steatosis based on transient elastography. *Liver International* 2012;32(6):902-10.
1714. Myers, R. P., G. Pomier-Layrargues, et al. Feasibility and diagnostic performance of the FibroScan XL probe for liver stiffness measurement in overweight and obese patients. *Hepatology* 2012;55(1):199-208.
1715. Myers, R. P. Reply: Assessment by Fibroscan of fibrosis in nonalcoholic fatty liver disease: XL versus M probe? *Hepatology* 2012;55(4):1309-1310.
1716. Lutz, H. H., N. Gassler, et al. Doppler ultrasound of hepatic blood flow for noninvasive evaluation of liver fibrosis compared with liver biopsy and transient elastography. *Digestive Diseases & Sciences* 2012;57(8):2222-30.
1717. McCorry, R. B., N. Palaniyappan, et al. Development and evaluation of a nurse-led transient elastography service for the staging of hepatic fibrosis in patients with suspected chronic liver disease. *QJM: Monthly journal of the Association of Physicians* 2012;105(8):749-54.
1718. Martinez, S. M., J. Foucher, et al. Longitudinal liver stiffness assessment in patients with chronic hepatitis C undergoing antiviral therapy. *PLoS ONE* 2012;7(10):e47715.
1719. Mehta, S. H., G. D. Kirk, et al. Stability of liver fibrosis among HCV-infected injection drug users. *Antiviral Therapy* 2012;17(5):813-21.
1720. Hagara, D., H. R. Schwarzenbach, et al. [New aspects in sonography of the liver]. *Praxis (Bern 1994)* 2012;101(18):1161-6.
1721. Jackson, B. D., J. S. Doyle, et al. Non-cirrhotic portal hypertension in HIV mono-infected patients. *Journal of Gastroenterology & Hepatology* 2012;27(9):1512-9.
1722. Kumar, R., M. K. Sharma, et al. Does reference value of "Normal Liver Stiffness" using fibroscan have a regional variation? *Hepatology* 2012;56(3):1183-5.
1723. Lee, S. W., H. J. Park, et al. Leflunomide increases the risk of silent liver fibrosis in patients with rheumatoid arthritis receiving methotrexate. *Arthritis Research & Therapy* 2012;14(5):R232.
1724. Kovac, J. D., M. Dakovic, et al. Diffusion-weighted MRI versus transient elastography in quantification of liver fibrosis in patients with chronic cholestatic liver diseases. *European Journal of Radiology* 2012;81(10):2500-6.
1725. Karlas, T., M. Hempel, et al. Noninvasive evaluation of hepatic manifestation in Wilson disease with transient elastography, ARFI, and different fibrosis scores. *Scandinavian Journal of Gastroenterology* 2012;47(11):1353-61.
1726. Li Vecchi, V., M. Soresi, et al. Prospective evaluation of hepatic steatosis in HIV-infected patients with or without hepatitis C virus co-infection. *International Journal of Infectious Diseases* 2012;16(5):e397-402.
1727. Kircheis, G., A. Sagir, et al. Evaluation of acoustic radiation force impulse imaging for determination of liver stiffness using transient elastography as a reference. *World Journal of Gastroenterology* 2012;18(10):1077-84.
1728. Hopper, I., W. Kemp, et al. Impact of heart failure and changes to volume status on liver stiffness: non-invasive assessment using transient elastography. *European Journal of Heart Failure* 2012;14(6):621-7.
1729. Jung, K. S. and S. U. Kim. Clinical applications of transient elastography. *Clinical and Molecular Hepatology* 2012;18(2):163-73.
1730. Jean J. Santiago-Hernández, Francisco A. Villegas-López, et al. Utilidad de la elastografía de transición (Fibroscan®) en hepatopatías crónicas. *Medicina Interna de México* 2012;28(4):360-364.

1731. Kim, S. U., J. K. Kim, et al. Discordance between Liver Biopsy and FibroScan® in Assessing Liver Fibrosis in Chronic Hepatitis B: Risk Factors and Influence of Necroinflammation. *PLoS ONE* 2012;7(2).
1732. Klibansky, D. A., S. H. Mehta, et al. Transient elastography for predicting clinical outcomes in patients with chronic liver disease. *Journal of viral hepatitis* 2012;19(2):e184-93.
1733. Kumar, R., A. Rastogi, et al. Unconjugated hyperbilirubinemia in patients with non-alcoholic fatty liver disease: a favorable endogenous response. *Clinical Biochemistry* 2012;45(3):272-4.
1734. Lindvig, K., B. K. Mossner, et al. Liver stiffness and 30-day mortality in a cohort of patients admitted to hospital. *European Journal of Clinical Investigation* 2012;42(2):146-52.
1735. Jung, K. S., S. U. Kim, et al. Prediction of recurrence after curative resection of hepatocellular carcinoma using liver stiffness measurement (FibroScan(R)). *Annals of Surgical Oncology* 2012;19(13):4278-86.
1736. Li, F., T. Yan, et al. [FibroScan can be used to diagnose the size of oesophageal varices in patients with HBV-related cirrhosis]. *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 2012;26(6):470-3.
1737. Kuroda, H., K. Kakisaka, et al. Non-invasive determination of hepatic steatosis by acoustic structure quantification from ultrasound echo amplitude. *World Journal of Gastroenterology* 2012;18(29):3889-95.
1738. Lu, L. G. [Significance of FibroScan in the diagnosis of liver fibrosis]. *Zhonghua Gan Zang Bing Za Zhi* 2012;20(8):571-2.
1739. Kemp, W. and S. Roberts. Feasibility and performance of the FibroScan XL probe. *Hepatology* 2012;55(4):1308-9; author reply 1309-10.
1740. Leroy, V. and S. U. Kim. Can transient elastography be used for the management of chronic hepatitis B patients? *Liver International* 2012;32(4):528-30.
1741. Karlas, T., M. Neuschulz, et al. Non-Invasive Evaluation of Cystic Fibrosis Related Liver Disease in Adults with ARFI, Transient Elastography and Different Fibrosis Scores. *PLoS ONE* 2012;7(7):e42139.
1742. Kim, S. U., J. H. Lee, et al. Prediction of liver-related events using fibroscan in chronic hepatitis B patients showing advanced liver fibrosis. *PLoS ONE* 2012;7(5):e36676.
1743. Kim, B. K., H. S. Kim, et al. Prospective validation of ELF test in comparison with FibroScan and FibroTest to predict liver fibrosis in Asian subjects with chronic hepatitis B. *PLoS ONE* 2012;7(7):e41964.
1744. Colecchia, A., L. Montrone, et al. Measurement of spleen stiffness to evaluate portal hypertension and the presence of esophageal varices in patients with HCV-related cirrhosis. *Gastroenterology* 2012;143(3):646-54.
1745. Ferraioli, G., R. Lissandrini, et al. Assessment of liver stiffness with transient elastography by using S and M probes in healthy children. *European Journal of Pediatrics* 2012;171(9):1415; author reply 1417.
1746. Fraquelli, M., C. Rigamonti, et al. Spleen stiffness measured by transient elastography accurately predicts esophageal varices in liver cirrhosis. *Gastroenterology* 2012;143(4):e23; author reply e23-4.
1747. Du, D., X. Zhu, et al. Comparison of LecT-Hepa and FibroScan for assessment of liver fibrosis in hepatitis B virus infected patients with different ALT levels. *Clinica Chimica Acta* 2012;413(21-22):1796-9.
1748. Cescon, M., A. Colecchia, et al. Value of transient elastography measured with FibroScan in predicting the outcome of hepatic resection for hepatocellular carcinoma. *Annals of Surgery* 2012;256(5):706-12; discussion 712-3.
1749. de Ledinghen, V., J. Vergniol, et al. Screening for liver fibrosis by using FibroScan((R)) and FibroTest in patients with diabetes. *Digestive & Liver Disease* 2012;44(5):413-8.
1750. Fransen van de Putte, D. E., K. Fischer, et al. Beneficial effect of successful HCV treatment in patients with inherited bleeding disorders, assessed by liver stiffness measurements. *Haemophilia* 2012;18(3):e266-72.
1751. Chowdhury, A. and K. Das. Reply: Does reference value of "Normal Liver Stiffness" using fibroscan have a regional variation? *Hepatology* 2012;56(3):1184-1185.
1752. Chen, Y. P., Q. Zhang, et al. Is transient elastography valuable for high-risk esophageal varices prediction in patients with hepatitis-B-related cirrhosis? *Journal of Gastroenterology & Hepatology* 2012;27(3):533-9.
1753. Crisan, D., C. Radu, et al. Two or more synchronous combination of noninvasive tests to increase accuracy of liver fibrosis assessment in chronic hepatitis C; results from a cohort of 446 patients. *Hepatitis Monthly* 2012;12(3):177-84.
1754. Dang, H. W., S. U. Kim, et al. How many valid measurements are necessary to assess liver fibrosis using FibroScan(R) in patients with chronic viral hepatitis? An analysis of subjects with at least 10 valid measurements. *Yonsei Medical Journal* 2012;53(2):337-45.
1755. Friedrich-Rust, M., D. Romen, et al. Acoustic radiation force impulse-imaging and transient elastography for non-invasive assessment of liver fibrosis and steatosis in NAFLD. *European Journal of Radiology* 2012;81(3):e325-31.

1756. Garcia-Jurado, L., I. Oyaguez, et al. [Evaluation of the costs of Transient Elastography (FibroScan) in the diagnosis of liver fibrosis in HIV patients with hepatitis C virus.]. *Enfermedades Infecciosas y Microbiología Clínica* 2012;30(6):294-9.
1757. Chen, Y. P., X. E. Liang, et al. Larger biopsies evaluation of transient elastography for detecting advanced fibrosis in patients with compensated chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2012;27(7):1219-26.
1758. Chon, Y. E., E. S. Jung, et al. The accuracy of noninvasive methods in predicting the development of hepatocellular carcinoma and hepatic decompensation in patients with chronic hepatitis B. *Journal of Clinical Gastroenterology* 2012;46(6):518-25.
1759. Chon, Y. E., S. U. Kim, et al. Transient elastography in real practice: a time for action? *Journal of Gastroenterology & Hepatology* 2012;27(7):1134-5.
1760. Corpechot, C., F. Carrat, et al. Noninvasive elastography-based assessment of liver fibrosis progression and prognosis in primary biliary cirrhosis. *Hepatology* 2012;56(1):198-208.
1761. de Ledinghen, V., J. Vergniol, et al. Non-invasive diagnosis of liver steatosis using controlled attenuation parameter (CAP) and transient elastography. *Liver International* 2012;32(6):911-8.
1762. Ferraioli, G., C. Tinelli, et al. Performance of Real-Time Strain Elastography, Transient Elastography, and aspartate-to-platelet ratio index in the assessment of fibrosis in chronic Hepatitis C. *American journal of roentgenology* 2012;199(1):19-25.
1763. Chen, Y. P., X. E. Liang, et al. Improving transient elastography performance for detecting hepatitis B cirrhosis. *Digestive & Liver Disease* 2012;44(1):61-6.
1764. Das, K., R. Sarkar, et al. "Normal" liver stiffness measure (LSM) values are higher in both lean and obese individuals: a population-based study from a developing country. *Hepatology* 2012;55(2):584-93.
1765. Ding, H., T. Wu, et al. Noninvasive measurement of liver fibrosis by transient elastography and influencing factors in patients with chronic hepatitis B-A single center retrospective study of 466 patients. *Journal of Huazhong University of Science and Technology. Medical sciences* 2012;32(1):69-74.
1766. Engelmann, G., C. Gebhardt, et al. Feasibility study and control values of transient elastography in healthy children. *European Journal of Pediatrics* 2012;171(2):353-60.
1767. Friedrich-Rust, M., J. Nierhoff, et al. Performance of Acoustic Radiation Force Impulse imaging for the staging of liver fibrosis: a pooled meta-analysis. *Journal of viral hepatitis* 2012;19(2):e212-9.
1768. Collazos, J., V. Asensi, et al. The effect of unsuccessful treatment with peginterferon and ribavirin on the liver fibrosis course of HIV/HCV-coinfected patients. *Current HIV Research* 2012;10(8):679-87.
1769. Crisan, D., C. Radu, et al. Prospective non-invasive follow-up of liver fibrosis in patients with chronic hepatitis C. *Journal of Gastrointestinal & Liver Diseases* 2012;21(4):375-82.
1770. Fernandez-Varo, G. [Serological markers of fibrosis]. *Gastroenterologia y hepatologia* 2012;35 Suppl 2:10-6.
1771. Forestier, N., A. Gaus, et al. Acoustic radiation force impulse imaging for evaluation of antiviral treatment response in chronic hepatitis C. *Journal of Gastrointestinal & Liver Diseases* 2012;21(4):367-73.
1772. Fraquelli, M. and R. Pozzi. The accuracy of noninvasive methods in the prediction of clinically relevant outcomes in patients with chronic liver disease. *Expert Review of Gastroenterology & Hepatology* 2012;6(6):679-82.
1773. Crespo, G., G. Fernandez-Varo, et al. ARFI, Fibroscan, ELF, and their combinations in the assessment of liver fibrosis: a prospective study. *Journal of Hepatology* 2012;57(2):281-7.
1774. Colombo, S., M. Buonocore, et al. Head-to-head comparison of transient elastography (TE), real-time tissue elastography (RTE), and acoustic radiation force impulse (ARFI) imaging in the diagnosis of liver fibrosis. *Journal of Gastroenterology* 2012;47(4):461-9.
1775. de Ledinghen, V., V. W. Wong, et al. Diagnosis of liver fibrosis and cirrhosis using liver stiffness measurement: comparison between M and XL probe of FibroScan(R). *Journal of Hepatology* 2012;56(4):833-9.
1776. Chon, Y. E., E. H. Choi, et al. Performance of transient elastography for the staging of liver fibrosis in patients with chronic hepatitis B: a meta-analysis. *PLoS ONE* 2012;7(9):e44930.
1777. Friedrich-Rust, M., P. Buggisch, et al. Acoustic radiation force impulse imaging for non-invasive assessment of liver fibrosis in chronic hepatitis B. *Journal of viral hepatitis* 2012;19(12).
1778. Castera, L. Noninvasive methods to assess liver disease in patients with hepatitis B or C. *Gastroenterology* 2012;142(6):1293-1302 e4.
1779. Castera, L., M. Pinzani, et al. Non invasive evaluation of portal hypertension using transient elastography. *Journal of Hepatology* 2012;56(3):696-703.
1780. Casey, S. P., W. W. Kemp, et al. A prospective evaluation of the role of transient elastography for the detection of hepatic fibrosis in type 2 diabetes without overt liver disease. *Scandinavian Journal of Gastroenterology* 2012;47(7):836-41.

1781. Barbero-Villares, A., J. Mendoza Jimenez-Ridruejo, et al. Evaluation of liver fibrosis by transient elastography (Fibroscan(R)) in patients with inflammatory bowel disease treated with methotrexate: a multicentric trial. *Scandinavian Journal of Gastroenterology* 2012;47(5):575-9.
1782. Barreiro, P., E. Vispo, et al. Management and treatment of chronic hepatitis C in HIV patients. *Seminars in Liver Disease* 2012;32(2):138-46.
1783. Berzigotti, A. Transient elastography and prognosis of cirrhosis. *Hepatology* 2012;55(5):1629-31.
1784. Bray, A. P., I. Barnova, et al. Liver fibrosis screening for patients with psoriasis taking methotrexate: a cross-sectional study comparing transient elastography and liver biopsy. *British Journal of Dermatology* 2012;166(5):1125-7.
1785. Boursier, J. and P. Cales. Controlled attenuation parameter (CAP): a new device for fast evaluation of liver fat? *Liver International* 2012;32(6):875-7.
1786. Boursier, J., V. de Ledinghen, et al. Comparison of eight diagnostic algorithms for liver fibrosis in hepatitis C: new algorithms are more precise and entirely noninvasive. *Hepatology* 2012;55(1):58-67.
1787. Boursier, J. and P. Calès. Reply. *Hepatology* 2012;55(2):653-653.
1788. Barsic, N., I. Lerotic, et al. Overview and developments in noninvasive diagnosis of nonalcoholic fatty liver disease. *World Journal of Gastroenterology* 2012;18(30):3945-54.
1789. Cardoso, A. C., R. J. Carvalho-Filho, et al. Direct comparison of diagnostic performance of transient elastography in patients with chronic hepatitis B and chronic hepatitis C. *Liver International* 2012;32(4):612-21.
1790. Afdhal, N. H. Fibroscan (transient elastography) for the measurement of liver fibrosis. *Gastroenterology & Hepatology* 2012;8(9):605-7.
1791. Arienti, V., L. Aluigi, et al. Ultrasonography (US) and non-invasive diagnostic methods for non-alcoholic fatty liver disease (NAFLD) and early vascular damage. Possible application in a population study on the metabolic syndrome (MS). *Internal and Emergency Medicine* 2012;7 Suppl 3:S283-90.
1792. Abenavoli, L. and M. Beaugrand. Transient elastography in non-alcoholic fatty liver disease. *Annals of Hepatology* 2012;11(2):172-8.
1793. Adebajo, C. O., J. A. Talwalkar, et al. Ultrasound-based transient elastography for the detection of hepatic fibrosis in patients with recurrent hepatitis C virus after liver transplantation: a systematic review and meta-analysis. *Liver Transplantation* 2012;18(3):323-31.
1794. Aalaei-Andabili, S. H., L. Mehrnoush, et al. Liver Hemangioma Might Lead to overestimation of Liver Fibrosis by Fibroscan; A Missed Issue in Two Cases. *Hepatitis Monthly* 2012;12(6):408-10.
1795. Arora, A. and P. Sharma. Non-invasive Diagnosis of Fibrosis in Non-alcoholic Fatty Liver Disease. *Journal of Clinical and Experimental Hepatology* 2012;2(2):145-55.
1796. Andreozzi, P., G. Viscogliosi, et al. [Predictors of liver fibrosis in patients with non-alcoholic fatty liver disease. The role of metabolic syndrome, insulin-resistance and inflammation]. *Recenti Progressi in Medicina* 2012;103(12):570-4.
1797. Abraldes, J. G., I. K. Araujo, et al. Diagnosing and monitoring cirrhosis: Liver biopsy, hepatic venous pressure gradient and elastography. *Gastroenterología y hepatología* 2012;35(7):488-95.
1798. Al-Dahshan, M. Clinical application of transient elastography in prediction of portal hypertension related complication in patients with chronic liver diseases. *Journal of the Egyptian Society of Parasitology* 2012;42(1):79-88.
1799. Alvarez, D., M. Anders, et al. [Usefulness of transient elastography (Fibroscan(R)) in the assessment of fibrosis in patients with chronic liver disease]. *Medicina (B Aires)* 2012;72(1):10-4.
- 2011**
1800. Sporea, I., R. Sirli, et al. Is it better to use two elastographic methods for liver fibrosis assessment? *World Journal of Gastroenterology* 2011;17(33):3824-9.
1801. Zhu, X., L. C. Wang, et al. Prospective evaluation of FibroScan for the diagnosis of hepatic fibrosis compared with liver biopsy/AST platelet ratio index and FIB-4 in patients with chronic HBV infection. *Digestive Diseases & Sciences* 2011;56(9):2742-9.
1802. Wang, Q. X., L. Shen, et al. [Validation of transient elastography (Fibroscan) in assessment of hepatic fibrosis in autoimmune hepatitis]. *Zhonghua Gan Zang Bing Za Zhi* 2011;19(10):782-4.
1803. Wong, V. W. and J. J. Sung. Application of transient elastography in chronic hepatitis B. *Alimentary Pharmacology & Therapeutics* 2011;34(7):817-8; author reply 818-9.
1804. Trifan, A., C. Sfarti, et al. Increased liver stiffness in extrahepatic cholestasis caused by choledocholithiasis. *Hepatitis Monthly* 2011;11(5):372-5.
1805. Tarantino, G., A. Spano, et al. Is spleen circulation impaired in systemic sclerosis and what is the role of liver fibrosis? *World Journal of Gastroenterology* 2011;17(12):1606-13.
1806. Sporea, I., D. Nicolita, et al. Assessment of noninvasive liver stiffness in inactive HBsAg carriers by transient elastography: Fibroscan in inactive HBsAg carriers. *Hepatitis Monthly* 2011;11(3):182-5.
1807. Stefanescu, H., M. Grigorescu, et al. A new and simple algorithm for the noninvasive assessment of esophageal varices in cirrhotic patients using serum

- fibrosis markers and transient elastography. *Journal of Gastrointestinal & Liver Diseases* 2011;20(1):57-64.
1808. Scourfield, A., L. Waters, et al. Non-cirrhotic portal hypertension in HIV-infected individuals. *International Journal of STD & AIDS* 2011;22(6):324-8.
1809. Vergniol, J., J. Foucher, et al. Noninvasive tests for fibrosis and liver stiffness predict 5 year outcomes of patients with chronic hepatitis C. *Gastroenterology* 2011;140(7):1970-9, 1979 1-3.
1810. Yang, H. I., M. F. Yuen, et al. Risk estimation for hepatocellular carcinoma in chronic hepatitis B (REACH-B): development and validation of a predictive score. *The Lancet Oncology* 2011;12(6):568-74.
1811. Sporea, I., R. Badea, et al. How efficient is acoustic radiation force impulse elastography for the evaluation of liver stiffness? *Hepatitis Monthly* 2011;11(7):532-8.
1812. Stibbe, K. J., C. Verveer, et al. Comparison of non-invasive assessment to diagnose liver fibrosis in chronic hepatitis B and C patients. *Scandinavian Journal of Gastroenterology* 2011;46(7-8):962-72.
1813. Vignier, N., G. Esmat, et al. Reproducibility of liver stiffness measurements in hepatitis C virus (HCV)-infected patients in Egypt. *Journal of viral hepatitis* 2011;18(7):e358-65.
1814. Sporea, I., R. L. Sirli, et al. Acoustic radiation force impulse elastography as compared to transient elastography and liver biopsy in patients with chronic hepatopathies. *Ultraschall in der Medizin* 2011;32 Suppl 1:S46-52.
1815. Stefanescu, H., M. Grigorescu, et al. Spleen stiffness measurement using Fibroscan for the noninvasive assessment of esophageal varices in liver cirrhosis patients. *Journal of Gastroenterology & Hepatology* 2011;26(1):164-70.
1816. Stoean, R., C. Stoean, et al. Evolutionary-driven support vector machines for determining the degree of liver fibrosis in chronic hepatitis C. *Artificial Intelligence in Medicine* 2011;51(1):53-65.
1817. Vallet-Pichard, A., S. Pol, et al. [Assessment and reversibility of liver fibrosis in viral hepatitis]. *La Revue du Praticien* 2011;61(1):39-43.
1818. Yashima, Y., T. Tsujino, et al. Increased liver elasticity in patients with biliary obstruction. *Journal of Gastroenterology* 2011;46(1):86-91.
1819. Sporea, I., R. L. Sirli, et al. What did we learn from the first 3,459 cases of liver stiffness measurement by transient elastography (FibroScan(R))? *Ultraschall in der Medizin* 2011;32(1):40-5.
1820. Thabut, D., R. Moreau, et al. Noninvasive assessment of portal hypertension in patients with cirrhosis. *Hepatology* 2011;53(2):683-94.
1821. Wong, G. L., V. W. Wong, et al. Factors associated with unreliable liver stiffness measurement and its failure with transient elastography in the Chinese population. *Journal of Gastroenterology & Hepatology* 2011;26(2):300-5.
1822. Vigano, M., S. Paggi, et al. Dual cut-off transient elastography to assess liver fibrosis in chronic hepatitis B: a cohort study with internal validation. *Alimentary Pharmacology & Therapeutics* 2011;34(3):353-62.
1823. Wong, G. L., V. W. Wong, et al. Is transient elastography inaccurate in chronic hepatitis B and non-alcoholic fatty liver disease? *Journal of Hepatology* 2011;55(2):497; author reply 498.
1824. Stoean, C., R. Stoean, et al. Feature selection for a cooperative coevolutionary classifier in liver fibrosis diagnosis. *Computers in Biology and Medicine* 2011;41(4):238-46.
1825. Tsochatzis, E. A., K. S. Gurusamy, et al. Elastography for the diagnosis of severity of fibrosis in chronic liver disease: a meta-analysis of diagnostic accuracy. *Journal of Hepatology* 2011;54(4):650-9.
1826. Stabinski, L., S. J. Reynolds, et al. High prevalence of liver fibrosis associated with HIV infection: a study in rural Rakai, Uganda. *Antiviral Therapy* 2011;16(3):405-11.
1827. Topilskaja, N. V., S. V. Morozov, et al. [Elastography efficiency in the liver fibrosis determination in patients with nonalcoholic fatty liver disease]. *Eksperimental Klinical Gastroenterology* 2011;-(6):26-31.
1828. Wong, G. L., V. W. Wong, et al. On-treatment monitoring of liver fibrosis with transient elastography in chronic hepatitis B patients. *Antiviral Therapy* 2011;16(2):165-72.
1829. Wong GL1, W. V., Chim AM, Yiu KK, Chu SH, Li MK, Chan HL. Factors associated with unreliable liver stiffness measurement and its failure with transient elastography in the chinese population. *Journal of Gastroenterology & Hepatology* 2011;26(2):300-5.
1830. Fransen van de Putte, D. E., K. Fischer, et al. Liver stiffness measurements to assess progression of fibrosis in HCV-infected patients with inherited bleeding disorders. *Haemophilia* 2011;17(5):e975-80.
1831. Hezode, C., L. Castera, et al. Liver stiffness diminishes with antiviral response in chronic hepatitis C. *Alimentary Pharmacology & Therapeutics* 2011;34(6):656-63.
1832. Malbrunot-Wagner, A. C., L. Bridoux, et al. Transient elastography and portal hypertension in pediatric patients with cystic fibrosis Transient elastography and cystic fibrosis. *Journal of Cystic Fibrosis* 2011;10(5):338-42.
1833. Mori, M., H. Fujii, et al. Close correlation of liver stiffness with collagen deposition and presence of myofibroblasts in non-alcoholic fatty liver disease. *Hepatology Research* 2011;41(9):897-903.

1834. Fransen van de Putte, D., R. Blom, et al. Impact of Fibroscan on management of chronic viral hepatitis in clinical practice. *Annals of Hepatology* 2011;10(4):469-76.
1835. Ebinuma, H., H. Saito, et al. Evaluation of liver fibrosis by transient elastography using acoustic radiation force impulse: comparison with Fibroscan(R). *Journal of Gastroenterology* 2011;46(10):1238-48.
1836. Fung, J., C. L. Lai, et al. Prognostic significance of liver stiffness for hepatocellular carcinoma and mortality in HBeAg-negative chronic hepatitis B. *Journal of viral hepatitis* 2011;18(10):738-44.
1837. Guha, I. N., R. P. Myers, et al. Biomarkers of liver fibrosis: what lies beneath the receiver operating characteristic curve? *Hepatology* 2011;54(4):1454-62.
1838. Lesmana, C. R., S. Salim, et al. Diagnostic accuracy of transient elastography (FibroScan) versus the aspartate transaminase to platelet ratio index in assessing liver fibrosis in chronic hepatitis B: the role in primary care setting. *Journal of clinical pathology* 2011;64(10):916-20.
1839. Nobili, V., L. Monti, et al. Transient elastography for assessment of fibrosis in paediatric liver disease. *Pediatric Radiology* 2011;41(10):1232-8.
1840. Sanchez-Conde, M., P. Miralles, et al. Use of transient elastography (FibroScan(R)) for the noninvasive assessment of portal hypertension in HIV/HCV-coinfected patients. *Journal of viral hepatitis* 2011;18(10):685-91.
1841. Koch, A., A. Horn, et al. Increased liver stiffness denotes hepatic dysfunction and mortality risk in critically ill non-cirrhotic patients at a medical ICU. *Critical Care* 2011;15(6):R266.
1842. Patel, K., M. Friedrich-Rust, et al. FibroSURE and FibroScan in relation to treatment response in chronic hepatitis C virus. *World Journal of Gastroenterology* 2011;17(41):4581-9.
1843. Floreani, A., N. Cazzagon, et al. Performance and utility of transient elastography and noninvasive markers of liver fibrosis in primary biliary cirrhosis. *Digestive & Liver Disease* 2011;43(11):887-92.
1844. Floreani, A., N. Cazzagon, et al. Female patients in fertile age with chronic hepatitis C, easy genotype, and persistently normal transaminases have a 100% chance to reach a sustained virological response. *European Journal of Gastroenterology & Hepatology* 2011;23(11):997-1003.
1845. Mehta, S. H. and G. C. Buckle. Assessment of liver disease (noninvasive methods). *Current Opinion in HIV and AIDS* 2011;6(6):465-71.
1846. Moessner, B. K., E. S. Andersen, et al. Previously unrecognized advanced liver disease unveiled by transient elastography in patients with Haemophilia and chronic hepatitis C. *Haemophilia* 2011;17(6):938-43.
1847. Osakabe, K., N. Ichino, et al. Reduction of liver stiffness by antiviral therapy in chronic hepatitis B. *Journal of Gastroenterology* 2011;46(11):1324-34.
1848. Poupon, T., V. de Ledinghen, et al. FibroTest and FibroScan performances revisited in patients with chronic hepatitis C. Impact of the spectrum effect and the applicability rate. *Clinics and Research in Hepatology and Gastroenterology* 2011;35(11):720-30.
1849. Robic, M. A., B. Procopet, et al. Liver stiffness accurately predicts portal hypertension related complications in patients with chronic liver disease: a prospective study. *Journal of Hepatology* 2011;55(5):1017-24.
1850. Deng, K. S., M. L. Cheng, et al. [Eighty three cases of post transfusion HCV infection: a 10-year follow up]. *Zhonghua Gan Zang Bing Za Zhi* 2011;19(5):349-51.
1851. Harata, M., S. Hashimoto, et al. Liver stiffness in extrahepatic cholestasis correlates positively with bilirubin and negatively with alanine aminotransferase. *Hepatology Research* 2011;41(5):423-9.
1852. Kummer, S., A. Sagir, et al. Liver fibrosis in recessive multicystic kidney diseases: transient elastography for early detection. *Pediatric Nephrology* 2011;26(5):725-31.
1853. Liu, C. H., C. C. Liang, et al. Transient elastography to assess hepatic fibrosis in hemodialysis chronic hepatitis C patients. *Clinical Journal of the American Society of Nephrology* 2011;6(5):1057-65.
1854. Moessner, B. K., T. R. Jorgensen, et al. Outreach screening of drug users for cirrhosis with transient elastography. *Addiction* 2011;106(5):970-6.
1855. Pichon, N., V. Loustaud-Ratti, et al. Value of liver stiffness measured by transient elastography in the liver transplant pre-operative evaluation of the potential deceased liver donors: preliminary study. *Clinical Transplantation* 2011;25(2):E205-10.
1856. Frulio, N., C. Balabaud, et al. Can elastometry be used for a better identification of cirrhosis? *Clinics and Research in Hepatology and Gastroenterology* 2011;35(3):166-8.
1857. Fung, J., C. L. Lai, et al. Mild-to-moderate elevation of alanine aminotransferase increases liver stiffness measurement by transient elastography in patients with chronic hepatitis B. *American Journal of Gastroenterology* 2011;106(3):492-6.
1858. Gelsi, E., R. Dainese, et al. Effect of detoxification on liver stiffness assessed by FibroScan(R) in alcoholic patients. *Alcoholism, clinical and experimental research* 2011;35(3):566-70.

1859. Jung, K. S., S. U. Kim, et al. Risk assessment of hepatitis B virus-related hepatocellular carcinoma development using liver stiffness measurement (FibroScan). *Hepatology* 2011;53(3):885-894.
1860. Kitson, M., S. Roberts, et al. The prevalence of significant liver fibrosis and cirrhosis in haemophilia patients infected with hepatitis C using FibroScan. *Haemophilia* 2011;17(2):316-7.
1861. Loustaud-Ratti, V. R., A. Cypierre, et al. Non-invasive detection of hepatic amyloidosis: FibroScan, a new tool. *Amyloid* 2011;18(1):19-24.
1862. Malekzadeh, R. and H. Poustchi. Fibroscan for assessing liver fibrosis: An acceptable alternative for liver biopsy: Fibroscan: an acceptable alternative for liver biopsy. *Hepatitis Monthly* 2011;11(3):157-8.
1863. Morikawa, H., K. Fukuda, et al. Real-time tissue elastography as a tool for the noninvasive assessment of liver stiffness in patients with chronic hepatitis C. *Journal of Gastroenterology* 2011;46(3):350-8.
1864. Rosen, H. R. Clinical practice. Chronic hepatitis C infection. *The New England Journal of Medicine* 2011;364(25):2429-38.
1865. Lee, H. J., Y. S. Seo, et al. Application of the HALF index obviates the need for liver biopsy in half of all patients with chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2011;26(6):987-95.
1866. Petta, S., V. Di Marco, et al. Reliability of liver stiffness measurement in non-alcoholic fatty liver disease: the effects of body mass index. *Alimentary Pharmacology & Therapeutics* 2011;33(12):1350-60.
1867. Pownard, T., Y. Ngo, et al. Noninvasive Markers of Hepatic Fibrosis in Chronic Hepatitis B. *Current Hepatitis Reports* 2011;10(2):87-97.
1868. Rifai, K., J. Cornberg, et al. Clinical feasibility of liver elastography by acoustic radiation force impulse imaging (ARFI). *Digestive & Liver Disease* 2011;43(6):491-7.
1869. Di Marco, V., V. Calvaruso, et al. Liver stiffness measurement by transient elastography predicts early recovery from acute hepatitis. *Gut* 2011;60(7):1023.
1870. Honsawek, S., K. Praianantathavorn, et al. High serum matrix metalloproteinase-3 and liver stiffness in postoperative biliary atresia. *Pediatric Surgery International* 2011;27(7):681-7.
1871. Reiberger, T., K. Rutter, et al. Portal Pressure Predicts Outcome and Safety of Antiviral Therapy in Cirrhotic Patients with HCV infection. *Clinical Gastroenterology & Hepatology* 2011;9(7):602-608 e 1.
1872. Roulot, D., J. L. Costes, et al. Transient elastography as a screening tool for liver fibrosis and cirrhosis in a community-based population aged over 45 years. *Gut* 2011;60(7):977-984.
1873. Sanchez-Conde, M., M. L. Montes Ramirez, et al. Impact of liver steatosis on the correlation between liver stiffness and fibrosis measured by transient elastography in patients coinfected with human immunodeficiency virus and hepatitis C virus. *Journal of viral hepatitis* 2011;18(7):278-83.
1874. Martin-Carbonero, L., T. Teixeira, et al. Clinical and virological outcomes in HIV-infected patients with chronic hepatitis B on long-term nucleos(t)ide analogues. *Aids* 2011;25(1):73-9.
1875. Fraquelli, M. and F. Branchi. The role of transient elastography in patients with hepatitis B viral disease. *Digestive & Liver Disease* 2011;43 Suppl 1:S25-31.
1876. Gaia, S., S. Carenzi, et al. Reliability of transient elastography for the detection of fibrosis in non-alcoholic fatty liver disease and chronic viral hepatitis. *Journal of Hepatology* 2011;54(1):64-71.
1877. Kahraman, A., O. Witzke, et al. Impact of immunosuppressive therapy on hepatitis C infection after renal transplantation. *Clinical Nephrology* 2011;75(1):16-25.
1878. Martinez, S. M., G. Crespo, et al. Noninvasive assessment of liver fibrosis. *Hepatology* 2011;53(1):325-35.
1879. Mialhes, P., P. Pradat, et al. Proficiency of transient elastography compared to liver biopsy for the assessment of fibrosis in HIV/HBV-coinfected patients. *Journal of viral hepatitis* 2011;18(1):61-69.
1880. Rallon, N. I., P. Barreiro, et al. Elevated TGF-beta1 levels might protect HCV/ HIV-coinfected patients from liver fibrosis. *European Journal of Clinical Investigation* 2011;41(1):70-6.
1881. Hodge, A. and W. Kemp. Transient elastography: the bigger we are, the harder to scan. *Journal of Gastroenterology & Hepatology* 2011;26(2):217-9.
1882. Honsawek, S., M. Chayanupatkul, et al. Serum adiponectin and transient elastography as non-invasive markers for postoperative biliary atresia. *BMC Gastroenterology* 2011;11:16.
1883. Messina, M. F., G. Squadrito, et al. Fibroscan: a new noninvasive method for evaluation of liver dysfunction in Turner syndrome. *European Journal of Clinical Investigation* 2011;41(2):183-8.
1884. Fung, J., C. L. Lai, et al. The use of transient elastography in the management of chronic hepatitis B. *Hepatology International* 2011;5(4):868-875.
1885. Lai, M. and N. H. Afdhal. Editorial: staging liver fibrosis in hepatitis C: a challenge for this decade. *American Journal of Gastroenterology* 2011;106(12):2121-2.
1886. Liang, X. E., Y. P. Chen, et al. Dynamic evaluation of liver stiffness measurement to improve diagnostic accuracy of liver cirrhosis in patients with chronic hepatitis B acute exacerbation. *Journal of viral hepatitis* 2011;18(12):884-91.

1887. Musso, G., R. Gambino, et al. Meta-analysis: natural history of non-alcoholic fatty liver disease (NAFLD) and diagnostic accuracy of non-invasive tests for liver disease severity. *Annals of Medicine* 2011;43(8):617-49.
1888. Ogawa, E., N. Furusyo, et al. Longitudinal assessment of liver stiffness by transient elastography for chronic hepatitis B patients treated with nucleoside analog. *Hepatology Research* 2011;41(12):1178-88.
1889. Rizzo, L., V. Calvaruso, et al. Comparison of transient elastography and acoustic radiation force impulse for non-invasive staging of liver fibrosis in patients with chronic hepatitis C. *American Journal of Gastroenterology* 2011;106(12):2112-20.
1890. Honsawek, S., V. Chongsrisawat, et al. Elevation of serum galectin-3 and liver stiffness measured by transient elastography in biliary atresia. *European Journal of Pediatric Surgery* 2011;21(4):250-4.
1891. Krawczyk, M., F. Grunhage, et al. Variant adiponutrin (PNPLA3) represents a common fibrosis risk gene: non-invasive elastography-based study in chronic liver disease. *Journal of Hepatology* 2011;55(2):299-306.
1892. Kumar, M., P. Sharma, et al. Transient elastographic evaluation in adult subjects without overt liver disease: influence of alanine aminotransferase levels. *Journal of Gastroenterology & Hepatology* 2011;26(8):1318-25.
1893. Martin-Carbonero, L., P. Tuma, et al. Treatment of chronic hepatitis C in HIV-infected patients with compensated liver cirrhosis. *Journal of viral hepatitis* 2011;18(8):542-8.
1894. Ong, A., V. W. Wong, et al. The effect of caffeine and alcohol consumption on liver fibrosis - a study of 1045 Asian hepatitis B patients using transient elastography. *Liver International* 2011;31(7):1047-53.
1895. Fraquelli, M., C. Rigamonti, et al. Etiology-related determinants of liver stiffness values in chronic viral hepatitis B or C. *Journal of Hepatology* 2011;54(4):621-8.
1896. Piscaglia, F., V. Salvatore, et al. Accuracy of VirtualTouch Acoustic Radiation Force Impulse (ARFI) imaging for the diagnosis of cirrhosis during liver ultrasonography. *Ultraschall in der Medizin* 2011;32(2):167-75.
1897. Pritchett, S., A. Cardenas, et al. The optimal cut-off for predicting large oesophageal varices using transient elastography is disease specific. *Journal of viral hepatitis* 2011;18(4):e75-80.
1898. Gaia, S. Reply to: "Is transient elastography inaccurate in chronic hepatitis B and non-alcoholic fatty liver disease?". *Journal of Hepatology* 2011;55(2):498.
1899. listed, N. a. [Elastography efficiency in the liver fibrosis determination in patients with nonalcoholic fatty liver disease]. *Eksperimental Klinical Gastroenterology* 2011;(6):26-31.
1900. Mbaye, P. S., A. Sarr, et al. Liver stiffness measurement and biochemical markers in Senegalese chronic hepatitis B patients with normal ALT and high viral load. *PLoS ONE* 2011;6(7):e22291.
1901. Papageorgiou, M. V., G. V. Papatheodoridis, et al. Elastography for hepatic fibrosis severity in chronic hepatitis B or C. *Case Reports in Gastroenterology* 2011;5(1):63-72.
1902. Pascu, O., M. Voiculescu, et al. Early virological response in 1220 patients with HCV (genotype 1b) chronic hepatitis and cirrhosis treated with PegInterferon plus Ribavirin. *Romanian Journal of Internal Medicine* 2011;49(2):105-12.
1903. Rockstroh, J. K., L. Peters, et al. Is there a need for liver disease monitoring in HIV patients in Africa? *Antiviral Therapy* 2011;16(3):287-9.
1904. Chen, Y. P., X. E. Liang, et al. [Transient elastography accurately predicts the severity of disease in patients with chronic hepatitis B]. *Zhonghua Nei Ke Za Zhi* 2011;50(9):758-62.
1905. Cohen-Naftaly, M. and S. L. Friedman. Current status of novel antifibrotic therapies in patients with chronic liver disease. *Therapeutic Advances in Gastroenterology* 2011;4(6):391-417.
1906. Clark, P. J. and K. Patel. Noninvasive tools to assess liver disease. *Current Opinion in Gastroenterology* 2011;27(3):210-6.
1907. Colombo, S., L. Belloli, et al. Normal liver stiffness and its determinants in healthy blood donors. *Digestive & Liver Disease* 2011;43(3):231-6.
1908. Castera, L. Non-invasive assessment of liver fibrosis in chronic hepatitis C. *Hepatology International* 2011;5(2):625-34.
1909. Castera, L. Screening the general population for cirrhosis using transient elastography: finding a needle in a haystack? *Gut* 2011;60(7):883-4.
1910. Collazos, J., J. A. Carton, et al. Gender differences in liver fibrosis and hepatitis C virus-related parameters in patients coinfected with human immunodeficiency virus. *Current HIV Research* 2011;9(5):339-45.
1911. Castera, L. and P. Bedossa. How to assess liver fibrosis in chronic hepatitis C: serum markers or transient elastography vs. liver biopsy? *Liver International* 2011;31 Suppl 1:13-7.
1912. Chan, H. L. and J. Jia. Chronic hepatitis B in Asia-new insights from the past decade. *Journal of Gastroenterology & Hepatology* 2011;26 Suppl 1:131-7.
1913. Chitturi, S., V. W. Wong, et al. Nonalcoholic fatty liver in Asia: Firmly entrenched and rapidly gaining ground. *Journal of Gastroenterology & Hepatology* 2011;26 Suppl 1:163-72.
1914. Castera, L., P. H. Bernard, et al. Transient elastography and biomarkers for liver fibrosis

- assessment and follow-up of inactive hepatitis B carriers. *Alimentary Pharmacology & Therapeutics* 2011;33(4):455-65.
1915. Chayanupatkul, M., S. Honsawek, et al. Serum retinol binding protein 4 and clinical outcome in postoperative biliary atresia. *Hepatology International* 2011;5(4):906-12.
1916. Carrión, J. A., M. Navasa, et al. [Hepatic elastography. Position paper of the Catalan Society of Gastroenterology]. *Gastroenterología y Hepatología* 2011;34(7):504-10.
1917. Chongsrisawat, V., P. Vejapipat, et al. Transient elastography for predicting esophageal/gastric varices in children with biliary atresia. *BMC Gastroenterology* 2011;11(1):41.
1918. Castera, L. Invasive and non-invasive methods for the assessment of fibrosis and disease progression in chronic liver disease. *Best Practice & Research Clinical Gastroenterology* 2011;25(2):291-303.
1919. Chen, X. B., X. Zhu, et al. [Accuracy of FibroScan for the diagnosis of liver fibrosis influenced by serum alanine aminotransferase levels in patients with chronic hepatitis B]. *Zhonghua Gan Zang Bing Za Zhi* 2011;19(4):286-90.
1920. Collazos, J., J. A. Carton, et al. Immunological status does not influence hepatitis c virus or liver fibrosis in HIV-hepatitis C virus-coinfected patients. *AIDS Research and Human Retroviruses* 2011;27(4):383-9.
1921. Carton, J. A., J. Collazos, et al. Factors associated with liver fibrosis in intravenous drug users coinfected with HIV and HCV. *Antiviral Therapy* 2011;16(1):27-35.
1922. Boursier, J., S. Bertrais, et al. Comparison of accuracy of fibrosis degree classifications by liver biopsy and non-invasive tests in chronic hepatitis C. *BMC Gastroenterology* 2011;11(1):132.
1923. Cardoso, A. C., R. J. Carvalho-Filho, et al. Transient elastography in chronic viral hepatitis: a critical appraisal. *Gut* 2011;60(6):759-64.
1924. Boursier, J., V. de Ledinghen, et al. A new combination of blood test and fibroscan for accurate non-invasive diagnosis of liver fibrosis stages in chronic hepatitis C. *American Journal of Gastroenterology* 2011;106(7):1255-63.
1925. Blanco, F., P. Barreiro, et al. Risk factors for advanced liver fibrosis in HIV-infected individuals: role of antiretroviral drugs and insulin resistance. *Journal of viral hepatitis* 2011;18(1):11-6.
1926. Bavu, E., J. L. Gennisson, et al. Noninvasive in vivo liver fibrosis evaluation using supersonic shear imaging: a clinical study on 113 hepatitis C virus patients. *Ultrasound in Medicine and Biology* 2011;37(9):1361-73.
1927. Barbero-Villares, A., J. Mendoza, et al. Evaluation of liver fibrosis by transient elastography in methotrexate treated patients. *Medicina clínica* 2011;137(14):637-9.
1928. Baba, M., K. Furuya, et al. Discrimination of individuals in a general population at high-risk for alcoholic and non-alcoholic fatty liver disease based on liver stiffness: a cross section study. *BMC Gastroenterology* 2011;11(1):70.
1929. Barreiro, P., J. A. Pineda, et al. Influence of interleukin-28B single-nucleotide polymorphisms on progression to liver cirrhosis in human immunodeficiency virus-hepatitis C virus-coinfected patients receiving antiretroviral therapy. *The Journal of Infectious Diseases* 2011;203(11):1629-36.
1930. Bastard, C., M. R. Bosisio, et al. Transient micro-elastography: A novel non-invasive approach to measure liver stiffness in mice. *World Journal of Gastroenterology* 2011;17(8):968-75.
1931. Baranova, A., P. Lal, et al. Non-invasive markers for hepatic fibrosis. *BMC Gastroenterology* 2011;11(1):91.
1932. Akima, T., M. Tamano, et al. Liver stiffness measured by transient elastography is a predictor of hepatocellular carcinoma development in viral hepatitis. *Hepatology Research* 2011;41(10):965-70.
1933. Andersen, E. S., M. Ruhwald, et al. Twelve potential fibrosis markers to differentiate mild liver fibrosis from cirrhosis in patients infected with chronic hepatitis C genotype 1. *European Journal of Clinical Microbiology & Infectious Diseases* 2011;30(6):761-6.
1934. Andersen, E. S., B. K. Moessner, et al. Lower liver stiffness in patients with sustained virological response 4 years after treatment for chronic hepatitis C. *European Journal of Gastroenterology & Hepatology* 2011;23(1):41-4.
1935. Abenavoli, L. (2011). Transient Elastography for Assessment of Non-Alcoholic Fatty Liver Disease. *Liver Biopsy*. H. Takahashi. **18:** 284-292.
1936. Ahmad, W., B. Ijaz, et al. A brief review on molecular, genetic and imaging techniques for HCV fibrosis evaluation. *Virology Journal* 2011;8(1):53.
- ## 2010
1937. Sporea, I., R. Sirli, et al. Liver stiffness measurements in patients with HBV vs HCV chronic hepatitis: a comparative study. *World Journal of Gastroenterology* 2010;16(38):4832-7.
1938. Wong, V. W. and H. L. Chan. Transient elastography. *Journal of Gastroenterology & Hepatology* 2010;25(11):1726-31.
1939. Wang, J.-H., C.-S. Changchien, et al. Liver stiffness decrease after effective antiviral therapy in patients with chronic hepatitis C: Longitudinal study using FibroScan. *Journal of Gastroenterology & Hepatology* 2010;25(5):964-969.
1940. Wong, G. L., V. W. Wong, et al. Development of a non-invasive algorithm with transient

- elastography (Fibroscan) and serum test formula for advanced liver fibrosis in chronic hepatitis B. *Alimentary Pharmacology & Therapeutics* 2010;31(10):1095-103.
1941. Tuma, P., I. Jarrin, et al. Survival of HIV-infected patients with compensated liver cirrhosis. *Aids* 2010;24(5):745-53.
1942. Sporea, I., R. Sirli, et al. Acoustic Radiation Force Impulse (ARFI)--a new modality for the evaluation of liver fibrosis. *Medical Ultrasonography Journal* 2010;12(1):26-31.
1943. Stebbing, J., L. Farouk, et al. A meta-analysis of transient elastography for the detection of hepatic fibrosis. *Journal of Clinical Gastroenterology* 2010;44(3):214-9.
1944. Seitz, M., S. Reichenbach, et al. Hepatoprotective effect of tumour necrosis factor alpha blockade in psoriatic arthritis: a cross-sectional study. *Annals of Rheumatic Diseases* 2010;69(6):1148-50.
1945. Tyagi, S. and S. Kumar. Clinical applications of elastography: an overview. *International Journal of Pharma & Bio Sciences* 2010;1(3).
1946. Tatsumi, C., M. Kudo, et al. Non-invasive evaluation of hepatic fibrosis for type C chronic hepatitis. *Intervirology* 2010;53(1):76-81.
1947. Wasmuth, H. E. and C. Trautwein. [Liver fibrosis : clinics, diagnostics and management]. *Der Internist* 2010;51(1):14-20.
1948. Vigano, M., S. Massironi, et al. Transient elastography assessment of the liver stiffness dynamics during acute hepatitis B. *European Journal of Gastroenterology & Hepatology* 2010;22(2):180-4.
1949. Wong, V. W., J. Vergniol, et al. Diagnosis of fibrosis and cirrhosis using liver stiffness measurement in nonalcoholic fatty liver disease. *Hepatology* 2010;51(2):454-462.
1950. Yoneda, M., K. Suzuki, et al. Nonalcoholic Fatty Liver Disease: US-based Acoustic Radiation Force Impulse Elastography. *Radiology* 2010;256(2):640-647.
1951. Zhang, Y. G., B. E. Wang, et al. Assessment of hepatic fibrosis by transient elastography in patients with chronic hepatitis B. *Pathology International* 2010;60(4):284-90.
1952. Scott, D. R. and M. T. Levy. Liver transient elastography (Fibroscan): a place in the management algorithms of chronic viral hepatitis. *Antiviral Therapy* 2010;15(1):1-11.
1953. Trufanova Iu, M., N. V. Topil'skaia, et al. [The efficacy of transient liver elastography overweight patients with chronic liver diseases]. *Eksperimental' Klinicheskaya Gastroenterologiya* 2010;(5):19-26.
1954. Tuma, P., J. Medrano, et al. Incidence of liver cirrhosis in HIV-infected patients with chronic hepatitis B or C in the era of highly active antiretroviral therapy. *Antiviral Therapy* 2010;15(6):881-6.
1955. Malik, R., M. Lai, et al. Comparison of transient elastography, serum markers and clinical signs for the diagnosis of compensated cirrhosis. *Journal of Gastroenterology & Hepatology* 2010;25(9):1562-1568.
1956. Pineda, J. A., J. Gonzalez, et al. Prevalence and factors associated with significant liver fibrosis assessed by transient elastometry in HIV/hepatitis C virus-coinfected patients. *Journal of viral hepatitis* 2010;17(10):714-9.
1957. Myers, R. P., M. Elkashab, et al. Transient elastography for the noninvasive assessment of liver fibrosis: a multicentre Canadian study. *Canadian Journal of Gastroenterology* 2010;24(11):661-670.
1958. Par, A. and G. Par. [Non-invasive fibrosis assessment in chronic hepatitis C: aspartate-aminotransferase to platelet ratio index (APRI) and transient elastography (FibroScan)]. *Orvosi Hetilap* 2010;151(47):1951-5.
1959. Li Vecchi, V., M. Soresi, et al. Transient elastography: a non-invasive tool for assessing liver fibrosis in HIV/HCV patients. *World Journal of Gastroenterology* 2010;16(41):5225-32.
1960. Myers, R. P., P. Crotty, et al. Prevalence, risk factors and causes of discordance in fibrosis staging by transient elastography and liver biopsy. *Liver International* 2010;30(10):1471-80.
1961. Sasso, M., M. Beaugrand, et al. Controlled attenuation parameter (CAP): a novel VCTE guided ultrasonic attenuation measurement for the evaluation of hepatic steatosis: preliminary study and validation in a cohort of patients with chronic liver disease from various causes. *Ultrasound in Medicine and Biology* 2010;36(11):1825-1835.
1962. Mueller, S. and L. Sandrin. Liver stiffness: a novel parameter for the diagnosis of liver disease. *Hepatic Medicine: Evidence and Research* 2010;2:49-67.
1963. Lupsor, M., R. Badea, et al. Performance of unidimensional transient elastography in staging non-alcoholic steatohepatitis. *Journal of Gastrointestinal & Liver Diseases* 2010;19(1):53-60.
1964. Pambrun, E., V. Bouteloup, et al. On the validity of meta-analyses: exhaustivity must be warranted, exclusion of duplicate patients too. *Journal of clinical epidemiology* 2010;63(3):342-3.
1965. Lanzi, A., A. Gianstefani, et al. Liver AL amyloidosis as a possible cause of high liver stiffness values. *European Journal of Gastroenterology & Hepatology* 2010;22(7):895-7.
1966. Mendoza, J., M. Trapero-Marugan, et al. Hepatic fibrosis in patients with chronic hepatitis C assessed by transient elastography: implications for determining the efficacy of antiviral therapy. *Revista*

- Española de Enfermedades Digestivas 2010;102(7):426-34.
1967. Menten, R., A. Leonard, et al. Transient elastography in patients with cystic fibrosis. *Pediatric Radiology* 2010;40(7):1231-5.
1968. Neukam, K., E. Recio, et al. Interobserver concordance in the assessment of liver fibrosis in HIV/HCV-coinfected patients using transient elastometry. *European Journal of Gastroenterology & Hepatology* 2010;22(7):801-7.
1969. Nguyen-Khac, E., P. Saint-Leger, et al. Noninvasive diagnosis of large esophageal varices by Fibroscan: strong influence of the cirrhosis etiology. *Alcoholism, clinical and experimental research* 2010;34(7):1146-53.
1970. Maor, Y., P. Halfon, et al. Fibrotest or Fibroscan for evaluation of liver fibrosis in haemophilia patients infected with hepatitis C. *Haemophilia* 2010;16(1):148-54.
1971. Mueller, S., G. Millonig, et al. Increased liver stiffness in alcoholic liver disease: differentiating fibrosis from steatohepatitis. *World Journal of Gastroenterology* 2010;16(8):966-72.
1972. listed., N. a. Diagnosis of fibrosis and cirrhosis. Liver biopsy is not always necessary. *Prescrire International* 2010;19(105):38-42.
1973. Millonig, G., S. Friedrich, et al. Liver stiffness is directly influenced by central venous pressure. *Journal of Hepatology* 2010;52(2):206-10.
1974. Laharie, D., J. Seneschal, et al. Assessment of liver fibrosis with transient elastography and FibroTest in patients treated with methotrexate for chronic inflammatory diseases: a case-control study. *Journal of Hepatology* 2010;53(6):1035-40.
1975. Lee, M. H., J. Y. Cheong, et al. Comparison of surrogate serum markers and transient elastography (Fibroscan) for assessing cirrhosis in patients with chronic viral hepatitis. *Digestive Diseases & Sciences* 2010;55(12):3552-60.
1976. Maida, I., L. Martin-Carbonero, et al. Differences in liver fibrosis and response to pegylated interferon plus ribavirin in HIV/HCV-coinfected patients with normal vs elevated liver enzymes. *Journal of viral hepatitis* 2010;17(12):866-71.
1977. Park, S.-H., J.-Y. Choe, et al. Assessment of liver fibrosis by transient elastography in rheumatoid arthritis patients treated with methotrexate. *Joint Bone Spine* 2010;77(6):588-592.
1978. Rigamonti, C., M. F. Donato, et al. Transient elastography in the early prediction of progressive recurrent hepatitis C following liver transplantation. *Hepatology* 2010;52(2):800-1.
1979. Riggio, S., F. Mamone, et al. Assessment of liver stiffness in subjects affected by familial combined hyperlipidaemia with hepatic steatosis. *European Journal of Clinical Investigation* 2010;40(8):722-8.
1980. Laharie, D., J. Vergniol, et al. Usefulness of noninvasive tests in nodular regenerative hyperplasia of the liver. *European Journal of Gastroenterology & Hepatology* 2010;22(4):487-93.
1981. Puoti, C., L. Bellis, et al. HCV carriers with normal alanine aminotransferase levels: healthy persons or severely ill patients? Dealing with an everyday clinical problem. *European Journal of Internal Medicine* 2010;21(2):57-61.
1982. Sanchez-Conde, M., M. L. Montes-Ramirez, et al. Comparison of transient elastography and liver biopsy for the assessment of liver fibrosis in HIV/hepatitis C virus-coinfected patients and correlation with noninvasive serum markers. *Journal of viral hepatitis* 2010;17(4):280-6.
1983. Lazebnik, L. B., E. V. Vinnitskaia, et al. [Liver fibroelastometry in the elderly patients]. *Eksperimental Klinikal Gastroenterology* 2010;2010(12):24-6.
1984. Lazebnik, L. B., E. V. Vinnitskaia, et al. [The diagnostic significance of ultrasound elastometry in evaluation of fibrosis in chronic diffuse liver diseases]. *Eksperimental Klinikal Gastroenterology* 2010;2010(5):10-3.
1985. Nunnari, G., C. Vancheri, et al. Circulating fibrocytes as a marker of liver fibrosis in chronic hepatitis C. *Frontiers in Bioscience* 2010;2:1241-5.
1986. Poynard, T., Y. Ngo, et al. Biomarkers of liver injury for hepatitis clinical trials: a meta-analysis of longitudinal studies. *Antiviral Therapy* 2010;15(4):617-31.
1987. Kamphues, C., K. Lotz, et al. Chances and limitations of non-invasive tests in the assessment of liver fibrosis in liver transplant patients. *Clinical Transplantation* 2010;24(5):652-9.
1988. Friedrich-Rust, M., W. Rosenberg, et al. Comparison of ELF, FibroTest and FibroScan for the non-invasive assessment of liver fibrosis. *BMC Gastroenterology* 2010;10(1):103.
1989. Dechene, A., J. P. Sowa, et al. Acute liver failure is associated with elevated liver stiffness and hepatic stellate cell activation. *Hepatology* 2010;52(3):1008-16.
1990. Enomoto, M., M. Mori, et al. Usefulness of transient elastography for assessment of liver fibrosis in chronic hepatitis B: Regression of liver stiffness during entecavir therapy. *Hepatology Research* 2010;40(9):853-61.
1991. Janssens, F., N. de Suray, et al. Can transient elastography replace liver histology for determination of advanced fibrosis in alcoholic patients: a real-life study. *Journal of Clinical Gastroenterology* 2010;44(8):575-82.
1992. Kim, S. U., K. H. Han, et al. Non-invasive assessment of liver fibrosis: time to move from cross-sectional studies to longitudinal ones. *Journal of Gastroenterology & Hepatology* 2010;25(9):1472-3.

1993. Ichino, N., K. Osakabe, et al. A new index for non-invasive assessment of liver fibrosis. *World Journal of Gastroenterology* 2010;16(38):4809-16.
1994. Friedrich-Rust, M., H. Hadji-Hosseini, et al. Transient elastography with a new probe for obese patients for non-invasive staging of non-alcoholic steatohepatitis. *European Radiology* 2010;20(10):2390-6.
1995. Hagymasi, K. and G. Lengyel. [Non-alcoholic steatosis/steatohepatitis -- 2010]. *Orvosi Hetilap* 2010;151(47):1940-5.
1996. Honsawek, S., M. Chayanupatkul, et al. Increased osteopontin and liver stiffness measurement by transient elastography in biliary atresia. *World Journal of Gastroenterology* 2010;16(43):5467-73.
1997. de Ledinghen, V. and J. Vergniol. Transient elastography for the diagnosis of liver fibrosis. *Expert Review of Medical Devices* 2010;7(6):811-23.
1998. Forestier, J., J. Dumortier, et al. Noninvasive diagnosis and prognosis of liver cirrhosis: a comparison of biological scores, elastometry, and metabolic liver function tests. *European Journal of Gastroenterology & Hepatology* 2010;22(5):532-40.
1999. Fung, J., C. L. Lai, et al. Correlation of liver stiffness and histological features in healthy persons and in patients with occult hepatitis B, chronic active hepatitis B, or hepatitis B cirrhosis. *American Journal of Gastroenterology* 2010;105(5):1116-22.
2000. Friedrich-Rust, M. and E. Herrmann. Response to letter: on the validity of meta-analysis. *Journal of clinical epidemiology* 2010;63(3):343.
2001. Gonzalez Guilabert, M. I., C. Hinojosa Menabernal, et al. [Retrospective study of FibroScan, APRI, FIB-4 and FORNS indexes compared with liver biopsy in the evaluation of liver fibrosis in patients with chronic hepatitis C monoinfection and HIV coinfection]. *Gastroenterologia y hepatología* 2010;33(6):425-32.
2002. Haafiz, A. B. Liver fibrosis in biliary atresia. *Expert Review of Gastroenterology & Hepatology* 2010;4(3):335-43.
2003. Kim, B. K., K. H. Han, et al. A liver stiffness measurement-based, noninvasive prediction model for high-risk esophageal varices in B-viral liver cirrhosis. *American Journal of Gastroenterology* 2010;105(6):1382-90.
2004. Ibanez, J. G., M. Perez, et al. [Grade of consistency existing in the grade of hepatic fibrosis calculated with the APRI and FORNS biochemistry indexes and transient elastography (Fibroscan in patients coinfected with HIV-HCV)]. *Revista clinica española* 2010;210(7):317-22.
2005. Kim, S. U., Y. Kim do, et al. The impact of steatosis on liver stiffness measurement in patients with chronic hepatitis B. *Hepatogastroenterology* 2010;57(101):832-8.
2006. Janssens, F., L. Spahr, et al. Hepatic amyloidosis increases liver stiffness measured by transient elastography. *Acta Gastroenterologica Belgica* 2010;73(1):52-4.
2007. Friedrich-Rust, M., C. Muller, et al. Assessment of liver fibrosis and steatosis in PBC with FibroScan, MRI, MR-spectroscopy, and serum markers. *Journal of Clinical Gastroenterology* 2010;44(1):58-65.
2008. Kim, S. U., Y. Kim do, et al. How can we enhance the performance of liver stiffness measurement using FibroScan in diagnosing liver cirrhosis in patients with chronic hepatitis B? *Journal of Clinical Gastroenterology* 2010;44(1):66-71.
2009. Di Marco, V., F. Bronte, et al. Noninvasive assessment of liver fibrosis in thalassaemia major patients by transient elastography (TE) - lack of interference by iron deposition. *British Journal of Haematology* 2010;148(3):476-9.
2010. Kim, S. U., G. H. Choi, et al. What are 'true normal' liver stiffness values using FibroScan?: a prospective study in healthy living liver and kidney donors in South Korea. *Liver International* 2010;30(2):268-74.
2011. Degos, F., P. Perez, et al. Diagnostic accuracy of FibroScan and comparison to liver fibrosis biomarkers in chronic viral hepatitis: a multicenter prospective study (the FIBROSTIC study). *Journal of Hepatology* 2010;53(6):1013-21.
2012. de Ledinghen, V., J. Vergniol, et al. Feasibility of liver transient elastography with FibroScan using a new probe for obese patients. *Liver International* 2010;30(7):1043-8.
2013. Fraquelli, M., E. Cassinero, et al. Transient elastography in the assessment of liver fibrosis in adult thalassemia patients. *American Journal of Hematology* 2010;85(8):564-8.
2014. Kim, S. U., Y. S. Seo, et al. Factors that affect the diagnostic accuracy of liver fibrosis measurement by FibroScan in patients with chronic hepatitis B. *Alimentary Pharmacology & Therapeutics* 2010;32(3):498-505.
2015. Kim, B. K., K.-H. Han, et al. A novel liver stiffness measurement-based prediction model for cirrhosis in hepatitis B patients. *Liver International* 2010;30(7):1073-1081.
2016. Donato, M. F., C. Rigamonti, et al. Can protocol liver biopsy be avoided to evaluate post-transplant hepatitis C recurrence? Transient elastography makes it possible. *Digestive & Liver Disease* 2010;42(4):307.
2017. Fung, J., C. L. Lai, et al. Reduction of liver stiffness following resolution of acute flares of chronic hepatitis B. *Hepatology International* 2010;4(4):716-22.
2018. Golovanova, E. V., E. V. Vinnitskaia, et al. [Effectiveness of the new plant hepatoprotector--Ropren in patients with non-alcoholic

- steatohepatitis]. *Eksperimental Klinical Gastroenterology* 2010;2010(7):97-102.
2019. Horster, S., P. Mandel, et al. Comparing acoustic radiation force impulse imaging to transient elastography to assess liver stiffness in healthy volunteers with and without valsalva manoeuvre. *Clinical Hemorheology and Microcirculation* 2010;46(2-3):159-68.
2020. Inglot, M., A. Szymczak, et al. [Pathogenesis and methods evaluation of liver fibrosis in HIV/HCV co-infection]. *Przeglad Epidemiologiczny* 2010;64(4):465-71.
2021. Kim, S. U., J. Y. Park, et al. Non-invasive assessment of changes in liver fibrosis via liver stiffness measurement in patients with chronic hepatitis B: impact of antiviral treatment on fibrosis regression. *Hepatology International* 2010;4(4):673-80.
2022. Chan, H. L. Letters to the Editors: Potential pitfalls on the validation of liver stiffness measurement by Fibroscan with histological staging in chronic hepatitis B. *Alimentary Pharmacology & Therapeutics* 2010;32(7):937-8.
2023. Cohen, E. B. and N. H. Afdhal. Ultrasound-based hepatic elastography: origins, limitations, and applications. *Journal of Clinical Gastroenterology* 2010;44(9):637-45.
2024. Cesari, M., M. Schiavini, et al. Noncirrhotic portal hypertension in HIV-infected patients: a case control evaluation and review of the literature. *AIDS Patient Care STDS* 2010;24(11):697-703.
2025. Castera, L., J. Foucher, et al. Pitfalls of liver stiffness measurement: a 5-year prospective study of 13,369 examinations. *Hepatology* 2010;51(3):828-35.
2026. Castera, L. Diagnosing cirrhosis non-invasively: sense the stiffness but don't forget the nodules! *Journal of Hepatology* 2010;52(6):786-7.
2027. Castera, L. and M. Pinzani. Biopsy and non-invasive methods for the diagnosis of liver fibrosis: does it take two to tango? *Gut* 2010;59(7):861-6.
2028. Carrion, J. A., F. Torres, et al. Liver stiffness identifies two different patterns of fibrosis progression in patients with hepatitis C virus recurrence after liver transplantation. *Hepatology* 2010;51(1):23-34.
2029. Castera, L., G. Sebastiani, et al. Prospective comparison of two algorithms combining non-invasive methods for staging liver fibrosis in chronic hepatitis C. *Journal of Hepatology* 2010;52(2):191-8.
2030. Colli, A., P. Pozzoni, et al. Decompensated chronic heart failure: increased liver stiffness measured by means of transient elastography. *Radiology* 2010;257(3):872-8.
2031. Cobbold, J. F., M. M. Crossey, et al. Optimal combinations of ultrasound-based and serum markers of disease severity in patients with chronic hepatitis C. *Journal of viral hepatitis* 2010;17(8):537-45.
2032. Cross, T. J., V. Calvaruso, et al. Prospective comparison of Fibroscan, King's score and liver biopsy for the assessment of cirrhosis in chronic hepatitis C infection. *Journal of viral hepatitis* 2010;17(8):546-54.
2033. Boursier, J., G. Isselin, et al. Acoustic radiation force impulse: a new ultrasonographic technology for the widespread noninvasive diagnosis of liver fibrosis. *European Journal of Gastroenterology & Hepatology* 2010;22(9):1074-84.
2034. Cales, P., J. Boursier, et al. Optimization and robustness of blood tests for liver fibrosis and cirrhosis. *Clinical Biochemistry* 2010;43(16-17):1315-22.
2035. Calvaruso, V., C. Camma, et al. Fibrosis staging in chronic hepatitis C: analysis of discordance between transient elastography and liver biopsy. *Journal of viral hepatitis* 2010;17(7):469-74.
2036. Boozari, B., A. Potthoff, et al. Evaluation of sound speed for detection of liver fibrosis: prospective comparison with transient dynamic elastography and histology. *Journal of ultrasound in medicine* 2010;29(11):1581-8.
2037. Boursier, J. and P. Cales. Clinical interpretation of Fibroscan(R) results: a real challenge. *Liver International* 2010;30(10):1400-2.
2038. Bonnard, P., R. Sombie, et al. Comparison of elastography, serum marker scores, and histology for the assessment of liver fibrosis in hepatitis B virus (HBV)-infected patients in Burkina Faso. *American Journal of Tropical Medicine and Hygiene* 2010;82(3):454-8.
2039. Berzigotti, A., J. G. Abraldes, et al. Ultrasonographic evaluation of liver surface and transient elastography in clinically doubtful cirrhosis. *Journal of Hepatology* 2010;52(6):846-53.
2040. Bonino, F., U. Arena, et al. Liver stiffness, a non-invasive marker of liver disease: a core study group report. *Antiviral Therapy* 2010;15 Suppl 3:69-78.
2041. Arndt, R., S. Schmidt, et al. Noninvasive evaluation of renal allograft fibrosis by transient elastography--a pilot study. *Transplant International* 2010;23(9):871-7.
2042. Bae, R. C., H. J. Cho, et al. [Clinical factors influencing liver stiffness as measured by transient elastography (Fibroscan) in patients with chronic liver disease]. *Korean Journal of Hepatology* 2010;16(2):123-30.
2043. Abdelaal, U. M., E. Morita, et al. Evaluation of portal hypertensive enteropathy by scoring with capsule endoscopy: is transient elastography of clinical impact? *Journal of Clinical Biochemistry and Nutrition* 2010;47(1):37-44.
2044. Audiere, S., M. Charbit, et al. Measurement of the skin-liver capsule distance on ultrasound RF data for 1D transient elastography. *Medical image*

- computing and computer assisted intervention 2010;13(Pt 2):34-41.
2045. Adams, L. Transient elastography in nonalcoholic fatty liver disease: making sense of echoes. *Hepatology* 2010;51(2):370-2.
2046. Anastasiou, J., A. Alisa, et al. Noninvasive markers of fibrosis and inflammation in clinical practice: prospective comparison with liver biopsy. *European Journal of Gastroenterology & Hepatology* 2010;22(4):474-80.
2047. Arima, Y., N. Kawabe, et al. Reduction of liver stiffness by interferon treatment in the patients with chronic hepatitis C. *Hepatology Research* 2010;40(4):383-92.

2009

2048. Petta, S., C. Camma, et al. Insulin resistance is a major determinant of liver stiffness in nondiabetic patients with HCV genotype 1 chronic hepatitis. *Alimentary Pharmacology & Therapeutics* 2009;30(6):603-13.
2049. Smith, J. O. and R. K. Sterling. Systematic review: non-invasive methods of fibrosis analysis in chronic hepatitis C. *Alimentary Pharmacology & Therapeutics* 2009;30(6):557-76.
2050. Cales, P., J. Boursier, et al. [FibroMeters: a family of blood tests for liver fibrosis with high diagnostic performance and applicability in clinical practice]. *Pathologie Biologie* 2009;57(6):459-62.
2051. Chang, H. K., Y. J. Park, et al. Hepatic fibrosis scan for liver stiffness score measurement: a useful preendoscopic screening test for the detection of varices in postoperative patients with biliary atresia. *Journal of Pediatric Gastroenterology and Nutrition* 2009;49(3):323-8.
2052. Kim, J. H., H. J. Yim, et al. [Changes in liver stiffness after acute or chronic liver injury due to viral hepatitis--does fibrosis exist after recovery from acute viral hepatitis?]. *Korean Journal of Gastroenterology* 2009;54(3):155-61.
2053. Lupsor, M., R. Badea, et al. Performance of a new elastographic method (ARFI technology) compared to unidimensional transient elastography in the noninvasive assessment of chronic hepatitis C. Preliminary results. *Journal of Gastrointestinal & Liver Diseases* 2009;18(3):303-10.
2054. Nie, J. J., J. Li, et al. [Value of FibroScan in clinical diagnosis]. *Zhonghua Gan Zang Bing Za Zhi* 2009;17(9):715-7.
2055. Pineda, J. A., M. Aguilar-Guisado, et al. Natural history of compensated hepatitis C virus-related cirrhosis in HIV-infected patients. *Clinical Infectious Diseases* 2009;49(8):1274-82.
2056. Friedrich-Rust, M., A. Schwarz, et al. Real-time tissue elastography versus FibroScan for noninvasive assessment of liver fibrosis in chronic

- liver disease. *Ultraschall in der Medizin* 2009;30(5):478-84.
2057. Segawa, M. and I. Sakaida. Diagnosis and treatment of portal hypertension. *Hepatology Research* 2009;39(10):1039-43.
2058. Vergniol, J. and V. de Ledinghen. Transient elastography (FibroScan): a new tool in hepatology. *La Presse Médicale* 2009;38(10):1516-25.
2059. Boursier, J., J. Vergniol, et al. The combination of a blood test and Fibroscan improves the non-invasive diagnosis of liver fibrosis. *Liver International* 2009;29(10):1507-15.
2060. Maimone, S., V. Calvaruso, et al. An evaluation of transient elastography in the discrimination of HBeAg-negative disease from inactive hepatitis B carriers. *Journal of viral hepatitis* 2009;16(11):769-74.
2061. Marin-Gabriel, J. C. and J. A. Solis-Herruzo. Noninvasive assessment of liver fibrosis. Serum markers and transient elastography (FibroScan). *Revista Española de Enfermedades Digestivas* 2009;101(11):787-99.
2062. Martin-Carbonero, L., V. de Ledinghen, et al. Liver fibrosis in patients with chronic hepatitis C and persistently normal liver enzymes: influence of HIV infection. *Journal of viral hepatitis* 2009;16(11):790-5.
2063. Mederacke, I., K. Wursthorn, et al. Food intake increases liver stiffness in patients with chronic or resolved hepatitis C virus infection. *Liver International* 2009;29(10):1500-6.
2064. Rubio, A., F. Monpoux, et al. Noninvasive procedures to evaluate liver involvement in HIV-1 vertically infected children. *Journal of Pediatric Gastroenterology and Nutrition* 2009;49(5):599-606.
2065. Zioli, M., A. Kettaneh, et al. Relationships between fibrosis amounts assessed by morphometry and liver stiffness measurements in chronic hepatitis or steatohepatitis. *European Journal of Gastroenterology & Hepatology* 2009;21(11):1261-8.
2066. Sebastiani, G. Non-invasive assessment of liver fibrosis in chronic liver diseases: implementation in clinical practice and decisional algorithms. *World Journal of Gastroenterology* 2009;15(18):2190-203.
2067. Cardenas, A. and P. Gines. Portal hypertension. *Current Opinion in Gastroenterology* 2009;25(3):195-201.
2068. Carlson, J. J., K. V. Kowdley, et al. An evaluation of the potential cost-effectiveness of non-invasive testing strategies in the diagnosis of significant liver fibrosis. *Journal of Gastroenterology & Hepatology* 2009;24(5):786-91.
2069. Castera, L. Transient elastography and other noninvasive tests to assess hepatic fibrosis in patients with viral hepatitis. *Journal of viral hepatitis* 2009;16(5):300-14.

2070. Friedrich-Rust, M. and S. Zeuzem. Reproducibility and limitations of transient elastography. *Liver International* 2009;29(5):619-20.
2071. Kim, S. U., J. K. Kim, et al. Variability in liver stiffness values from different intercostal spaces. *Liver International* 2009;29(5):760-6.
2072. Rotman, Y., T. A. Brown, et al. Evaluation of the patient with hepatitis B. *Hepatology* 2009;49(5 Suppl):S22-7.
2073. Takemoto, R., M. Nakamura, et al. Validity of FibroScan values for predicting hepatic fibrosis stage in patients with chronic HCV infection. *Journal of Digestive Diseases* 2009;10(2):145-8.
2074. Rockstroh, J. K. Hot topics in HIV and hepatitis coinfection: noninvasive diagnosis of liver disease, liver transplantation, and new drugs for treatment of hepatitis coinfection. *HIV Clinical Trials* 2009;10(2):110-5.
2075. Del Poggio, P. and S. Colombo. Is transient elastography a useful tool for screening liver disease? *World Journal of Gastroenterology* 2009;15(12):1409-14.
2076. Wang, J. H., C. S. Changchien, et al. FibroScan and ultrasonography in the prediction of hepatic fibrosis in patients with chronic viral hepatitis. *Journal of Gastroenterology* 2009;44(5):439-46.
2077. Corradi, F., F. Piscaglia, et al. Assessment of liver fibrosis in transplant recipients with recurrent HCV infection: usefulness of transient elastography. *Digestive & Liver Disease* 2009;41(3):217-25.
2078. Gatti, F., P. Nasta, et al. Unboosted fosamprenavir is associated with low drug exposure in HIV-infected patients with mild-moderate liver impairment resulting from HCV-related cirrhosis. *The Journal of Antimicrobial Chemotherapy* 2009;63(3):575-8.
2079. Kim, S. U., S. H. Ahn, et al. Liver stiffness measurement in combination with noninvasive markers for the improved diagnosis of B-viral liver cirrhosis. *Journal of Clinical Gastroenterology* 2009;43(3):267-71.
2080. Kim, S. G., Y. S. Kim, et al. [The usefulness of transient elastography to diagnose cirrhosis in patients with alcoholic liver disease]. *Korean Journal of Hepatology* 2009;15(1):42-51.
2081. Montaldo, G., M. Tanter, et al. Coherent plane-wave compounding for very high frame rate ultrasonography and transient elastography. *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control* 2009;56(3):489-506.
2082. Sirli, R., I. Sporea, et al. Transient elastographic evaluation of subjects without known hepatic pathology: does age change the liver stiffness? *Journal of Gastrointestinal & Liver Disease* 2009;18(1):57-60.
2083. Song, Z. Z. Extrahepatic cholestasis and liver stiffness by transient elastography. *Hepatology* 2009;49(3):1053; author reply 1053.
2084. Lemoine, M. and L. Serfaty. [How to diagnose the nonalcoholic fatty liver disease?]. *La Lettre de l'Hépato-Gastroentérologue* 2009;12(3):80-85.
2085. Carrion, J. A. [Utility of Fibroscan in the evaluation of liver fibrosis]. *Gastroenterología y Hepatología* 2009;32(6):415-23.
2086. Masuzaki, R., N. Yamashiki, et al. Assessment of liver stiffness in patients after living donor liver transplantation by transient elastography. *Scandinavian Journal of Gastroenterology* 2009;44(9):1115-20.
2087. Lee, A. U. When liver stiffness is not so straight forward and Fibroscan not so simple. *Journal of Gastroenterology & Hepatology* 2009;24(6):934-6.
2088. Masuzaki, R., R. Tateishi, et al. Prospective risk assessment for hepatocellular carcinoma development in patients with chronic hepatitis C by transient elastography. *Hepatology* 2009;49(6):1954-1961.
2089. Sebastiani, G., P. Halfon, et al. SAFE biopsy: a validated method for large-scale staging of liver fibrosis in chronic hepatitis C. *Hepatology* 2009;49(6):1821-7.
2090. Wong, G. L., V. W. Wong, et al. Increased liver stiffness measurement by transient elastography in severe acute exacerbation of chronic hepatitis B. *Journal of Gastroenterology & Hepatology* 2009;24(6):1002-7.
2091. Munoz, R., E. Ramirez, et al. Correlation between fibroscan, liver biopsy, and clinical liver function in patients with hepatitis C virus infection after renal transplantation. *Transplantation Proceedings* 2009;41(6):2425-6.
2092. Neukam, K., J. Macias, et al. Impact of observer experience on the reproducibility of transient elastometry in HIV/HCV co-infected patients. *HIV Clinical Trials* 2009;10(4):276-81.
2093. Ingiliz, P., K. P. Chhay, et al. Applicability and variability of liver stiffness measurements according to probe position. *World Journal of Gastroenterology* 2009;15(27):3398-404.
2094. Iser, D. M. and S. R. Lewin. Future directions in the treatment of HIV-HBV coinfection. *HIV Therapy* 2009;3(4):405-415.
2095. Breton, E., L. Bridoux-Henno, et al. [Value of transient elastography in noninvasive assessment in children's hepatic fibrosis]. *Archives de Pédiatrie* 2009;16(7):1005-10.
2096. Cales, P., J. Boursier, et al. Ultrasonographic elastometry vs. blood tests for the diagnosis of cirrhosis: a Pyrrhic victory? *Journal of Hepatology* 2009;51(1):228-9; author reply 229.
2097. Kim, S. U., K. H. Han, et al. Liver stiffness measurement using FibroScan is influenced by serum total bilirubin in acute hepatitis. *Liver International* 2009;29(6):810-5.
2098. Nitta, Y., N. Kawabe, et al. Liver stiffness measured by transient elastography correlates with

- fibrosis area in liver biopsy in patients with chronic hepatitis C. *Hepatology Research* 2009;39(7):675-84.
2099. Camma, C., S. Petta, et al. Insulin resistance is a risk factor for esophageal varices in hepatitis C virus cirrhosis. *Hepatology* 2009;49(1):195-203.
2100. Castéra, L., B. Le Bail, et al. Early detection in routine clinical practice of cirrhosis and oesophageal varices in chronic hepatitis C: comparison of transient elastography (FibroScan) with standard laboratory tests and non-invasive scores. *Journal of Hepatology* 2009;50(1):59-68.
2101. Chan, H. L., G. L. Wong, et al. Alanine aminotransferase-based algorithms of liver stiffness measurement by transient elastography (FibroScan) for liver fibrosis in chronic hepatitis B. *Journal of viral hepatitis* 2009;16(1):36-44.
2102. Ismail, M. H. and M. Pinzani. Reversal of liver fibrosis. *Saudi Journal of Gastroenterology* 2009;15(1):72-9.
2103. Wong, G. L., V. W. Wong, et al. Metabolic syndrome increases the risk of liver cirrhosis in chronic hepatitis B. *Gut* 2009;58(1):111-7.
2104. Foucher, J., B. Reiller, et al. FibroScan used in street-based outreach for drug users is useful for hepatitis C virus screening and management: a prospective study. *Journal of viral hepatitis* 2009;16(2):121-31.
2105. Marcellin, P., M. Ziol, et al. Non-invasive assessment of liver fibrosis by stiffness measurement in patients with chronic hepatitis B. *Liver International* 2009;29 (2):242-247.
2106. Nahon, P., A. Kettaneh, et al. Liver stiffness measurement in patients with cirrhosis and hepatocellular carcinoma: a case-control study. *European Journal of Gastroenterology & Hepatology* 2009;21(2):214-9.
2107. Salles, N., P. Dussarat, et al. Non-invasive evaluation of liver fibrosis by transient elastography and biochemical markers in elderly inpatients. *Gastroentérologie Clinique & Biologique* 2009;33(2):126-32.
2108. Vergniol, J., J. Foucher, et al. Changes of non-invasive markers and FibroScan values during HCV treatment. *Journal of viral hepatitis* 2009;16(2):132-40.
2109. Wong, G. L., V. W. Wong, et al. Clinical factors associated with liver stiffness in hepatitis B e antigen-positive chronic hepatitis B patients. *Clinical Gastroenterology & Hepatology* 2009;7(2):227-33.
2110. Zhou, K. and L. G. Lu. Assessment of fibrosis in chronic liver diseases. *Journal of Digestive Diseases* 2009;10(1):7-14.
2111. Castéra, L., X. Forns, et al. Interpreting liver stiffness in the cirrhotic range: What are we measuring? *Journal of Hepatology* 2009;50(2):424-425.
2112. Panos, G., L. Farouk, et al. Cryptogenic pseudocirrhosis: a new clinical syndrome of noncirrhotic portal hypertension (unassociated with advanced fibrosis) that can be detected by transient elastography in patients with HIV. *Journal of Acquired Immune Deficiency Syndromes* 2009;52(4):525-7.
2113. Chan, H. L., G. L. Wong, et al. Hepatitis B virus genotype C is associated with more severe liver fibrosis than genotype B. *Clinical Gastroenterology & Hepatology* 2009;7(12):1361-6.
2114. Seo, Y. S. [Transient elastography, true or false?]. *Korean Journal of Hepatology* 2009;15(4):431-7.
2115. Stasi, C., U. Arena, et al. Transient elastography for the assessment of liver fibrosis in patients with chronic viral hepatitis: the missing tool? *Digestive & Liver Disease* 2009;41(12):863-6.
2116. Witters, P., K. De Boeck, et al. Non-invasive liver elastography (FibroScan) for detection of cystic fibrosis-associated liver disease. *Journal of Cystic Fibrosis* 2009;8(6):392-9.
2117. Lebrec, D. and R. Moreau. [Progress in portal hypertension]. *Gastroentérologie Clinique & Biologique* 2009;33(8-9):799-810.
2118. Pineda, J. A., E. Recio, et al. Liver stiffness as a predictor of esophageal varices requiring therapy in HIV/hepatitis C virus-coinfected patients with cirrhosis. *Journal of Acquired Immune Deficiency Syndromes* 2009;51(4):445-9.
2119. Friedrich-Rust, M., K. Wunder, et al. Liver fibrosis in viral hepatitis: noninvasive assessment with acoustic radiation force impulse imaging versus transient elastography. *Radiology* 2009;252(2):595-604.
2120. Kim do, Y., S. U. Kim, et al. Usefulness of FibroScan for detection of early compensated liver cirrhosis in chronic hepatitis B. *Digestive Diseases & Sciences* 2009;54(8):1758-63.
2121. Ogawa, E., N. Furusyo, et al. The longitudinal quantitative assessment by transient elastography of chronic hepatitis C patients treated with pegylated interferon alpha-2b and ribavirin. *Antiviral Research* 2009;83(2):127-34.
2122. Sundaram, S. S., P. Zeitler, et al. The metabolic syndrome and nonalcoholic fatty liver disease in children. *Current Opinion in Pediatrics* 2009;21(4):529-35.
2123. Uribe-Ramos, M. and N. Mendez-Sanchez. Why is transient elastography essential in hepatology? *Annals of Hepatology* 2009;8(2):148-9.
2124. Kirk, G. D., J. Astemborski, et al. Assessment of liver fibrosis by transient elastography in persons with hepatitis C virus infection or HIV-hepatitis C virus coinfection. *Clinical Infectious Diseases* 2009;48(7):963-72.

2125. Della Valle, S., F. Branchi, et al. [Transient elastography]. *Recenti Progressi in Medicina* 2009;100(4):195-201.
2126. Lucidarme, D., J. Foucher, et al. Factors of accuracy of transient elastography (fibroscan) for the diagnosis of liver fibrosis in chronic hepatitis C. *Hepatology* 2009;49(4):1083-9.
2127. Moreno, S., J. Garcia-Samaniego, et al. Noninvasive diagnosis of liver fibrosis in patients with HIV infection and HCV/HBV co-infection. *Journal of viral hepatitis* 2009;16(4):249-58.
2128. Sanchez Antolin, G., F. Garcia Pajares, et al. FibroScan evaluation of liver fibrosis in liver transplantation. *Transplantation Proceedings* 2009;41(3):1044-6.
2129. Trinchet, J., V. Bourcier, et al. Compensated viral cirrhosis: descriptive analysis of 774 first patients included in a prospective French cohort. *BEH thématique* 2009;20-21(4):204-7.
2130. Chan, H. L., G. L. Wong, et al. A review of the natural history of chronic hepatitis B in the era of transient elastography. *Antiviral Therapy* 2009;14(4):489-99.
2131. De Ledighen, V. Évaluation non invasive de la fibrose hépatique. *La Revue du Praticien* 2009;23.
2132. de Ledighen, V. Diagnostic de la Fibrose Hépatique en 2009. *Axcan News* 2009;(16).
2133. Denzer, U. W. and S. Luth. Non-invasive diagnosis and monitoring of liver fibrosis and cirrhosis. *Best Practice & Research Clinical Gastroenterology* 2009;23(3):453-60.
2134. Dubertret, L. Retinoids, methotrexate and cyclosporine. *Current Problems in Dermatology* 2009;38:79-94.
2135. Myers, R. P. Noninvasive diagnosis of nonalcoholic fatty liver disease. *Annals of Hepatology* 2009;8 Suppl 1:S25-33.
2136. Pais, R., M. Lupsor, et al. Liver biopsy versus noninvasive methods--fibroscan and fibrotest in the diagnosis of non-alcoholic fatty liver disease: a review of the literature. *Romanian Journal of Internal Medicine* 2009;47(4):331-40.
2137. Vispo, E., P. Barreiro, et al. Overestimation of liver fibrosis staging using transient elastography in patients with chronic hepatitis C and significant liver inflammation. *Antiviral Therapy* 2009;14(2):187-93.
2138. Bedossa, P. and F. Carrat. Liver biopsy: the best, not the gold standard. *Journal of Hepatology* 2009;50(1):1-3.
2139. Bedossa, P. Assessment of hepatitis C: non-invasive fibrosis markers and/or liver biopsy. *Liver International* 2009;29 Suppl 1:19-22.
2140. Bonekamp, S., I. Kamel, et al. Can imaging modalities diagnose and stage hepatic fibrosis and cirrhosis accurately? *Journal of Hepatology* 2009;50(1):17-35.
2141. Bosch, J. Towards the non-invasive diagnosis of cirrhosis: the nuts-cirrhosis connection. *Journal of Hepatology* 2009;50(1):4-6.
2142. Bioulac-Sage, P., T. Couffinhal, et al. Interpreting liver stiffness in the cirrhotic range. *Journal of Hepatology* 2009;50(2):423-4; author reply 424-5.
2143. Angulo, P. Noninvasive assessment of fibrosis and steatosis in NASH and ASH. *Gastroentérologie Clinique & Biologique* 2009;33(10-11):940-8.
2144. Alric, L., N. Kamar, et al. Comparison of liver stiffness, fibrotest and liver biopsy for assessment of liver fibrosis in kidney-transplant patients with chronic viral hepatitis. *Transplant International* 2009;22(5):568-73.
2145. Andersen, E. S., P. B. Christensen, et al. Transient elastography for liver fibrosis diagnosis. *European Journal of Internal Medicine* 2009;20(4):339-42.
2146. Alisi, A., M. Pinzani, et al. Diagnostic power of fibroscan in predicting liver fibrosis in nonalcoholic fatty liver disease. *Hepatology* 2009;50(6):2048-9; author reply 2049-50.

2008

2147. Arena, U., F. Vizzutti, et al. Reliability of transient elastography for the diagnosis of advanced fibrosis in chronic hepatitis C. *Gut* 2008;57(9):1288-93.
2148. de Ledighen, V. and J. Vergniol. Transient elastography (FibroScan). *Gastroentérologie Clinique & Biologique* 2008;32(6 Suppl 1):58-67.
2149. Huwart, L. and B. E. van Beers. MR elastography. *Gastroentérologie Clinique & Biologique* 2008;32(6 Suppl 1):68-72.
2150. Jung, H. S., Y. S. Kim, et al. [Usefulness of liver stiffness measurement for predicting the presence of esophageal varices in patients with liver cirrhosis]. *Korean Journal of Hepatology* 2008;14(3):342-50.
2151. Masuzaki, R., R. Tateishi, et al. Comparison of liver biopsy and transient elastography based on clinical relevance. *Canadian Journal of Gastroenterology* 2008;22(9):753-7.
2152. Romanque, P., F. Stickel, et al. Disproportionally high results of transient elastography in patients with autoimmune hepatitis. *Liver International* 2008;28(8):1177-8.
2153. Wong, G. L., V. W. Wong, et al. Assessment of fibrosis by transient elastography compared with liver biopsy and morphometry in chronic liver diseases. *Clinical Gastroenterology & Hepatology* 2008;6(9):1027-35.
2154. Oliveri, F., B. Coco, et al. Liver stiffness in the hepatitis B virus carrier: a non-invasive marker of liver disease influenced by the pattern of

- transaminases. *World Journal of Gastroenterology* 2008;14(40):6154-62.
2155. de Franchis, R. Non-invasive (and minimally invasive) diagnosis of oesophageal varices. *Journal of Hepatology* 2008;49(4):520-7.
2156. Poinard, T., R. Morra, et al. Assessment of liver fibrosis: noninvasive means. *Saudi Journal of Gastroenterology* 2008;14(4):163-73.
2157. Song, Z. Z. Re: Transient elastography is unreliable for detection of cirrhosis in patients with acute liver damage. *Hepatology* 2008;48(4):1351-2.
2158. Yuen, M. F., D. K. Wong, et al. HBsAg Seroclearance in chronic hepatitis B in Asian patients: replicative level and risk of hepatocellular carcinoma. *Gastroenterology* 2008;135(4):1192-9.
2159. Nguyen-Khac, E., D. Chatelain, et al. Assessment of asymptomatic liver fibrosis in alcoholic patients using fibroscan: prospective comparison with seven non-invasive laboratory tests. *Alimentary Pharmacology & Therapeutics* 2008;28(10):1188-1198.
2160. Zois, C. D., G. H. Baltayiannis, et al. Systematic review: hepatic fibrosis - regression with therapy. *Alimentary Pharmacology & Therapeutics* 2008;28(10):1175-87.
2161. Sporea, I., R. Sirli, et al. Comparison of the liver stiffness measurement by transient elastography with the liver biopsy. *World Journal of Gastroenterology* 2008;14(42):6513-7.
2162. Lemoine, M., S. Katsahian, et al. Liver stiffness measurement as a predictive tool of clinically significant portal hypertension in patients with compensated hepatitis C virus or alcohol-related cirrhosis. *Alimentary Pharmacology & Therapeutics* 2008;28(9):1102-10.
2163. Bensamoun, S. F., L. Wang, et al. Measurement of liver stiffness with two imaging techniques: magnetic resonance elastography and ultrasound elastometry. *Journal of Magnetic Resonance Imaging* 2008;28(5):1287-92.
2164. Boursier, J., A. Konate, et al. Reproducibility of liver stiffness measurement by ultrasonographic elastometry. *Clinical Gastroenterology & Hepatology* 2008;6(11):1263-9.
2165. Millonig, G., F. M. Reimann, et al. Extrahepatic cholestasis increases liver stiffness (FibroScan) irrespective of fibrosis. *Hepatology* 2008;48(5):1718-23.
2166. Montes, M. L., J. M. Castro, et al. [Relationship between hepatic fibrosis measured by transient elastography and peripheral blood CD4+ lymphocyte values]. *Enfermedades Infecciosas y Microbiología Clínica* 2008;26(9):561-3.
2167. Nguyen-Khac, E., D. Capron, et al. The non-invasive diagnosis of cirrhosis using the FibroScan must be performed with cause-specific stiffness cut-offs. *Gut* 2008;57(11):1630; author reply 1630-1.
2168. Yoshioka, K., N. Kawabe, et al. Transient elastography: Applications and limitations. *Hepatology Research* 2008;38(11):1063-8.
2169. Castera, L., X. Forns, et al. Non-invasive evaluation of liver fibrosis using transient elastography. *Journal of Hepatology* 2008;48(5):835-47.
2170. de Franchis, R., A. Dell'Era, et al. Diagnosis and monitoring of portal hypertension. *Digestive & Liver Disease* 2008;40(5):312-7.
2171. Manning, D. S. and N. H. Afdhal. Diagnosis and quantitation of fibrosis. *Gastroenterology* 2008;134(6):1670-81.
2172. Van Leeuwen, D. J., C. Balabaud, et al. A clinical and histopathologic perspective on evolving noninvasive and invasive alternatives for liver biopsy. *Clinical Gastroenterology & Hepatology* 2008;6(5):491-6.
2173. Yoneda, M., M. Yoneda, et al. Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with nonalcoholic fatty liver disease (NAFLD). *Digestive & Liver Disease* 2008;40(5):371-8.
2174. Gomez-Dominguez, E., J. Mendoza, et al. Transient elastography to assess hepatic fibrosis in primary biliary cirrhosis. *Alimentary Pharmacology & Therapeutics* 2008;27(5):441-7.
2175. Castellares, C., P. Barreiro, et al. Liver cirrhosis in HIV-infected patients: prevalence, aetiology and clinical outcome. *Journal of viral hepatitis* 2008;15(3):165-72.
2176. de Ledinghen, V., T. Poinard, et al. [Non-invasive evaluation of liver fibrosis in hepatitis C]. *Gastroentérologie Clinique & Biologique* 2008;32(3 Pt 2):S90-5.
2177. Friedrich-Rust, M., C. Koch, et al. Noninvasive assessment of liver fibrosis in patients with Fontan circulation using transient elastography and biochemical fibrosis markers. *Journal of thoracic and cardiovascular surgery* 2008;135(3):560-7.
2178. Han, K. H. and K. T. Yoon. New diagnostic method for liver fibrosis and cirrhosis. *Intervirology* 2008;51 Suppl 1:11-6.
2179. Tatsumi, C., M. Kudo, et al. Noninvasive evaluation of hepatic fibrosis using serum fibrotic markers, transient elastography (FibroScan) and real-time tissue elastography. *Intervirology* 2008;51 Suppl 1:27-33.
2180. Rosa, H. and E. R. Parise. Is there a place for serum laminin determination in patients with liver disease and cancer? *World Journal of Gastroenterology* 2008;14(23):3628-32.
2181. Sporea, I., A. Popescu, et al. Why, who and how should perform liver biopsy in chronic liver diseases. *World Journal of Gastroenterology* 2008;14(21):3396-402.
2182. Bureau, C., S. Metivier, et al. Transient elastography accurately predicts presence of

- significant portal hypertension in patients with chronic liver disease. *Alimentary Pharmacology & Therapeutics* 2008;27(12):1261-8.
2183. de Ledinghen, V., P. Barreiro, et al. Liver fibrosis on account of chronic hepatitis C is more severe in HIV-positive than HIV-negative patients despite antiretroviral therapy. *Journal of viral hepatitis* 2008;15(6):427-33.
2184. Fung, J., C. L. Lai, et al. Prevalence of fibrosis and cirrhosis in chronic hepatitis B: implications for treatment and management. *American Journal of Gastroenterology* 2008;103(6):1421-6.
2185. Lupsor, M., R. Badea, et al. Analysis of histopathological changes that influence liver stiffness in chronic hepatitis C. Results from a cohort of 324 patients. *Journal of Gastrointestinal & Liver Diseases* 2008;17(2):155-63.
2186. Panos, G., P. Holmes, et al. Transient elastography, liver stiffness values, and acute hepatopathy. *Hepatology* 2008;47(6):2140; author reply 2141.
2187. Rigamonti, C., M. F. Donato, et al. Transient elastography predicts fibrosis progression in patients with recurrent hepatitis C after liver transplantation. *Gut* 2008;57(6):821-7.
2188. Abenavoli, L., G. Addolorato, et al. Elastography assessment in patients with chronic HCV infection. *Internal Journal of Clinical Practice* 2008;62(7):1108-12.
2189. Boursier, J., A. Konate, et al. Learning curve and interobserver reproducibility evaluation of liver stiffness measurement by transient elastography. *European Journal of Gastroenterology & Hepatology* 2008;20(7):693-701.
2190. Chang, P. E., H. F. Lui, et al. Prospective evaluation of transient elastography for the diagnosis of hepatic fibrosis in Asians: comparison with liver biopsy and aspartate transaminase platelet ratio index. *Alimentary Pharmacology & Therapeutics* 2008;28(1):51-61.
2191. Huwart, L., C. Sempoux, et al. Magnetic resonance elastography for the noninvasive staging of liver fibrosis. *Gastroenterology* 2008;135(1):32-40.
2192. Jacqueminet, S., P. Lebray, et al. Screening for liver fibrosis by using a noninvasive biomarker in patients with diabetes. *Clinical Gastroenterology & Hepatology* 2008;6(7):828-31.
2193. Song, Z. Z. Acute viral hepatitis increases liver stiffness values measured by transient elastography. *Hepatology* 2008;48(1):349-50; author reply 350.
2194. Talwalkar, J. A. Elastography for detecting hepatic fibrosis: options and considerations. *Gastroenterology* 2008;135(1):299-302.
2195. Fartoux, L. and L. Serfaty. [Comparison of liver biopsy, FibroScan® and FibroTest® for the assessment of liver fibrosis]. *La Lettre de l'Infectiologue* 2008;23(1):15-21.
2196. Harada, N., Y. Soejima, et al. Assessment of graft fibrosis by transient elastography in patients with recurrent hepatitis C after living donor liver transplantation. *Transplantation* 2008;85(1):69-74.
2197. Rockey, D. C. Noninvasive assessment of liver fibrosis and portal hypertension with transient elastography. *Gastroenterology* 2008;134(1):8-14.
2198. Adhoute, X., J. Foucher, et al. Diagnosis of liver fibrosis using FibroScan and other noninvasive methods in patients with hemochromatosis: a prospective study. *Gastroentérologie Clinique & Biologique* 2008;32(2):180-7.
2199. Arena, U., F. Vizzutti, et al. Acute viral hepatitis increases liver stiffness values measured by transient elastography. *Hepatology* 2008;47(2):380-4.
2200. Baranova, A. and Z. M. Younossi. The future is around the corner: Noninvasive diagnosis of progressive nonalcoholic steatohepatitis. *Hepatology* 2008;47(2):373-5.
2201. Bellentani, S., R. Dalle Grave, et al. Behavior therapy for nonalcoholic fatty liver disease: The need for a multidisciplinary approach. *Hepatology* 2008;47(2):746-54.
2202. Cobbold, J. F. and S. D. Taylor-Robinson. Transient elastography in acute hepatitis: All that's stiff is not fibrosis. *Hepatology* 2008;47(2):370-2.
2203. Laharie, D., E. Terrebonne, et al. [The liver and methotrexate]. *Gastroentérologie Clinique & Biologique* 2008;32(2):134-42.
2204. Pinzani, M., F. Vizzutti, et al. Technology insight: noninvasive assessment of liver fibrosis by biochemical scores and elastography. *Nature clinical practice. Gastroenterology & hepatology* 2008;5(2):95-106.
2205. Sagir, A., A. Erhardt, et al. Transient elastography is unreliable for detection of cirrhosis in patients with acute liver damage. *Hepatology* 2008;47(2):592-5.
2206. Castera, L. Non-invasive diagnosis of steatosis and fibrosis. *Diabetes & Metabolism* 2008;34(6 Pt 2):674-9.
2207. Dolmazashvili, E., M. Zhamutashvili, et al. Fibroscan and FibroTest/FibroMax to assess liver fibrosis/cirrhosis in patients with chronic HBV and HCV infection in Georgia. *Georgian Medical News* 2008;(165):83-7.
2208. Fung, J., C. L. Lai, et al. Correlation of liver biochemistry with liver stiffness in chronic hepatitis B and development of a predictive model for liver fibrosis. *Liver International* 2008;28(10):1408-16.
2209. Gheorghe, L., S. Iacob, et al. Real-time sonelastography - a new application in the field of liver disease. *Journal of Gastrointestinal & Liver Diseases* 2008;17(4):469-74.
2210. Kim, S. U., S. H. Ahn, et al. Prediction of postoperative hepatic insufficiency by liver stiffness measurement (FibroScan(R)) before curative

- resection of hepatocellular carcinoma: a pilot study. *Hepatology International* 2008;2(4):471-7.
2211. Lebray, P., S. Varnous, et al. Liver stiffness is an unreliable marker of liver fibrosis in patients with cardiac insufficiency. *Hepatology* 2008;48(6):2089.
2212. Macias, J., E. Recio, et al. Application of transient elastometry to differentiate mild from moderate to severe liver fibrosis in HIV/HCV co-infected patients. *Journal of Hepatology* 2008;49(6):916-22.
2213. Nahon, P., A. Kettaneh, et al. Assessment of liver fibrosis using transient elastography in patients with alcoholic liver disease. *Journal of Hepatology* 2008;49(6):1062-8.
2214. Seo, Y. S., S. H. Um, et al. Changes in liver stiffness during the course of acute hepatitis A. *Korean Journal of Hepatology* 2008;14(4):465-73.
2215. Sporea, I., R. Sirli, et al. Liver stiffness measurement by transient elastography in clinical practice. *Journal of Gastrointestinal & Liver Diseases* 2008;17(4):395-9.
2216. Tservadze, T., E. Dolmazashvili, et al. Assessment of liver fibrosis and cirrhosis by transient elastography among patients with chronic HBV and HCV infection in Georgia. *Georgian Medical News* 2008;(165):38-42.
2217. Vergniol, J., J. Foucher, et al. Non-invasive evaluation of liver fibrosis using FibroScan in long-term sustained-virological responder patients after HCV treatment. *Gastroentérologie Clinique & Biologique* 2008;32(12):1061-3.
2218. Wong, G. L., V. W. Wong, et al. Evaluation of alanine transaminase and hepatitis B virus DNA to predict liver cirrhosis in hepatitis B e antigen-negative chronic hepatitis B using transient elastography. *American Journal of Gastroenterology* 2008;103(12):3071-81.
2219. Mossner, B. K., C. Pedersen, et al. [Transient elastography for diagnosing liver fibrosis]. *Ugeskrift for Laeger* 2008;170(33):2453-6.
2220. Castera, L. Assessing liver fibrosis. *Expert Review of Gastroenterology & Hepatology* 2008;2(4):541-52.
2221. Kotlyar, D. S., W. Blonski, et al. Noninvasive monitoring of hepatitis C fibrosis progression. *Clinics in Liver Disease* 2008;12(3):557-71, viii.
2222. Masuzaki, R., R. Tateishi, et al. Risk assessment of hepatocellular carcinoma in chronic hepatitis C patients by transient elastography. *Journal of Clinical Gastroenterology* 2008;42(7):839-43.
2223. Nobili, V., F. Vizzutti, et al. Accuracy and reproducibility of transient elastography for the diagnosis of fibrosis in pediatric nonalcoholic steatohepatitis. *Hepatology* 2008;48(2):442-8.
2224. Oki, E., Y. Kakeji, et al. Transient elastography for the prediction of oxaliplatin-associated liver injury in colon cancer patients: a preliminary analysis. *Journal of Gastrointestinal Cancer* 2008;39(1-4):82-5.
2225. Cobbold, J. F. and S. D. Taylor-Robinson. Liver stiffness values in healthy subjects: implications for clinical practice. *Journal of Hepatology* 2008;48(4):529-31.
2226. Friedrich-Rust, M., M. F. Ong, et al. Performance of transient elastography for the staging of liver fibrosis: a meta-analysis. *Gastroenterology* 2008;134(4):960-74.
2227. Jung, M. K., H. J. Cho, et al. [Comparison of transient elastography and hepatic fibrosis assessed by histology in chronic liver disease]. *Korean Journal of Gastroenterology* 2008;51(4):241-7.
2228. Li, L. F., L. Dai, et al. [Factors influencing the success rate and stability of transient elastography for liver fibrosis evaluation]. *Nan Fang Yi Ke Da Xue Xue Bao* 2008;28(4):595-7.
2229. Mirault, T., D. Lucidarme, et al. Non-invasive assessment of liver fibrosis by transient elastography in post transfusional iron overload. *European Journal of Haematology* 2008;80(4):337-40.
2230. Roulot, D., S. Czernichow, et al. Liver stiffness values in apparently healthy subjects: influence of gender and metabolic syndrome. *Journal of Hepatology* 2008;48(4):606-13.
2231. Aita, J. S. and S. A. Harrison. Liver biopsy versus noninvasive testing in chronic hepatitis C: Where do we stand in 2008? *Current Hepatitis Reports* 2008;7(2):51-59.
2232. Leroy, V. and J.-P. Zarski. Clinical applicability of non-invasive markers for liver fibrosis in chronic hepatitis C. *Journal of Hepatology* 2008;48(5):882-884.
2233. Cainelli, F. Hepatitis C virus infection in the elderly: epidemiology, natural history and management. *Drugs & Aging* 2008;25(1):9-18.
2234. Maida, I., P. Garcia-Gasco, et al. Antiretroviral-associated portal hypertension: a new clinical condition? Prevalence, predictors and outcome. *Antiviral Therapy* 2008;13(1):103-7.
2235. Obara, N., Y. Ueno, et al. Transient elastography for measurement of liver stiffness measurement can detect early significant hepatic fibrosis in Japanese patients with viral and nonviral liver diseases. *Journal of Gastroenterology* 2008;43(9):720-8.
2236. Poynard, T., P. Ingiliz, et al. Concordance in a world without a gold standard: a new non-invasive methodology for improving accuracy of fibrosis markers. *PLoS ONE* 2008;3(12):e3857.
2237. Sheldon, J., B. Ramos, et al. Does treatment of hepatitis B virus (HBV) infection reduce hepatitis delta virus (HDV) replication in HIV-HBV-HDV-coinfected patients? *Antiviral Therapy* 2008;13(1):97-102.

2238. Vinnitskaia, E. V., N. A. Shaposhnikova, et al. [Liver cirrhosis of alcoholic etiology complicated by the spontaneous bacterial peritonitis]. *Eksperimental Klinical Gastroenterology* 2008;2008(7):129-32.
2239. Yeshua, H. and R. Oren. Non invasive assessment of liver fibrosis. *Annals of Transplantation* 2008;13(2):5-11.

2007

2240. Castera, L. [How to assess disease severity in chronic hepatitis B using non invasive fibrosis markers?]. *Hépato-Gastro & Oncologie Digestive* 2007;14(numéro spécial):23-28.
2241. Gressner, O. A., R. Weiskirchen, et al. Biomarkers of hepatic fibrosis, fibrogenesis and genetic pre-disposition pending between fiction and reality. *Journal of Cellular and Molecular Medicine* 2007;11(5):1031-51.
2242. Cobbold, J. F., S. Morin, et al. Transient elastography for the assessment of chronic liver disease: ready for the clinic? *World Journal of Gastroenterology* 2007;13(36):4791-7.
2243. Labarga, P., V. Soriano, et al. Hepatotoxicity of antiretroviral drugs is reduced after successful treatment of chronic hepatitis C in HIV-infected patients. *The Journal of Infectious Diseases* 2007;196(5):670-6.
2244. Yoneda, M., M. Yoneda, et al. Transient elastography in patients with non-alcoholic fatty liver disease (NAFLD). *Gut* 2007;56(9):1330-1.
2245. Vergara, S., J. Macias, et al. The use of transient elastometry for assessing liver fibrosis in patients with HIV and hepatitis C virus coinfection. *Clinical Infectious Diseases* 2007;45(8):969-74.
2246. de Ledinghen, V., B. Le Bail, et al. Liver stiffness measurement in children using FibroScan: feasibility study and comparison with Fibrotest, aspartate transaminase to platelets ratio index, and liver biopsy. *Journal of Pediatric Gastroenterology and Nutrition* 2007;45(4):443-50.
2247. Malik, R. and N. Afshar. Stiffness and impedance: the new liver biomarkers. *Clinical Gastroenterology & Hepatology* 2007;5(10):1144-6.
2248. Talwalkar, J. A., D. M. Kurtz, et al. Ultrasound-based transient elastography for the detection of hepatic fibrosis: systematic review and meta-analysis. *Clinical Gastroenterology & Hepatology* 2007;5(10):1214-20.
2249. Shaheen, A. A., A. F. Wan, et al. FibroTest and FibroScan for the prediction of hepatitis C-related fibrosis: a systematic review of diagnostic test accuracy. *American Journal of Gastroenterology* 2007;102(11):2589-600.
2250. Boursier, J., N. Dib, et al. Caractéristiques et interprétation des tests sanguins de fibrose hépatique. *Gastroentérologie Clinique & Biologique* 2007;31(5):511-523.
2251. Castera, L. [Use of elastometry (FibroScan) for the non-invasive staging of liver fibrosis]. *Gastroentérologie Clinique & Biologique* 2007;31(5):524-30; quiz 500, 531-2.
2252. Coco, B., F. Oliveri, et al. Transient elastography: a new surrogate marker of liver fibrosis influenced by major changes of transaminases. *Journal of viral hepatitis* 2007;14(5):360-9.
2253. Friedrich-Rust, M. and S. Zeuzem. [Transient elastography (FibroScan) for the non-invasive assessment of liver fibrosis: current status and perspectives]. *Zeitschrift für Gastroenterologie* 2007;45(5):387-94.
2254. Kim, K. M., W. B. Choi, et al. Diagnosis of hepatic steatosis and fibrosis by transient elastography in asymptomatic healthy individuals: a prospective study of living related potential liver donors. *Journal of Gastroenterology* 2007;42(5):382-8.
2255. Kim, H. S., K. K. Ja, et al. Non-invasive assessment of liver fibrosis by measuring the liver stiffness and biochemical markers in chronic hepatitis B patients. *Korean Journal of Medicine* 2007;72(5):459-469.
2256. Lim, J. K. and R. J. Groszmann. Transient elastography for diagnosis of portal hypertension in liver cirrhosis: is there still a role for hepatic venous pressure gradient measurement? *Hepatology* 2007;45(5):1087-90.
2257. Rendina, M., A. Schena, et al. The treatment of chronic hepatitis C with peginterferon alfa-2a (40 kDa) plus ribavirin in haemodialysed patients awaiting renal transplant. *Journal of Hepatology* 2007;46(5):768-74.
2258. Vizzutti, F., U. Arena, et al. Liver stiffness measurement predicts severe portal hypertension in patients with HCV-related cirrhosis. *Hepatology* 2007;45(5):1290-7.
2259. Castera, L. [FibroScan: a new tool for non-invasive assessment of fibrosis in chronic liver disease]. *Hépato-Gastro & Oncologie Digestive* 2007;14(2).
2260. Sebastiani, G. and A. Alberti. Non-invasive markers of liver fibrosis: How to use them in clinical practice? *Journal of Hepatology* 2007;46(3):529-530.
2261. Berrutti, M., A. Ciancio, et al. Assessment of liver fibrosis in the clinical setting: something is changing? *Minerva Gastroenterologica e Dietologica* 2007;53(1):111-4.
2262. Castera, L., J. Denis, et al. Evolving practices of non-invasive markers of liver fibrosis in patients with chronic hepatitis C in France: time for new guidelines? *Journal of Hepatology* 2007;46(3):528-9; author reply 529-30.

2263. Friedrich-Rust, M., M. F. Ong, et al. Real-time elastography for noninvasive assessment of liver fibrosis in chronic viral hepatitis. *American journal of roentgenology* 2007;188(3):758-64.
2264. Halfon, P., M. Bourliere, et al. [Noninvasive methods for predicting liver fibrosis in patients with chronic hepatitis C: alternatives to liver biopsy]. *La Presse Médicale* 2007;36(3 Pt 2):457-66.
2265. Berends, M. A., J. Snoek, et al. Biochemical and biophysical assessment of MTX-induced liver fibrosis in psoriasis patients: Fibrotest predicts the presence and Fibroscan predicts the absence of significant liver fibrosis. *Liver International* 2007;27(5):639-45.
2266. Cho, S. W. and J. Y. Cheong. [Clinical application of non-invasive diagnosis for hepatic fibrosis]. *Korean Journal of Hepatology* 2007;13(2):129-37.
2267. Maida, I., V. Soriano, et al. Liver fibrosis stage and HCV genotype distribution in HIV-HCV coinfected patients with persistently normal transaminases. *AIDS Research and Human Retroviruses* 2007;23(6):801-4.
2268. Rigamonti, C. and M. Fraquelli. Do not trivialize the Fibroscan examination, value its accuracy. *Journal of Hepatology* 2007;46(6):1149.
2269. Al-Ghamdi, A. S. Fibroscan: a noninvasive test of liver fibrosis assessment. *Saudi Journal of Gastroenterology* 2007;13(3):147-149.
2270. Arends, J. E., R. J. de Knegt, et al. [Diagnosis of hepatic fibrosis and cirrhosis]. *Nederlands Tijdschrift Voor Geneeskunde* 2007;151(27):1502-6.
2271. Fraquelli, M., C. Rigamonti, et al. Reproducibility of transient elastography in the evaluation of liver fibrosis in patients with chronic liver disease. *Gut* 2007;56(7):968-73.
2272. Fraquelli, M. and C. Rigamonti. Diagnosis of cirrhosis by transient elastography: what is hidden behind misleading results. *Hepatology* 2007;46(1):282; author reply 282-3.
2273. Lörke, J., A. Erhardt, et al. Non-invasive investigation of liver cirrhosis. *Deutsches Ärzteblatt international* 2007;24(104):1752-7.
2274. Castera, L. Reproductibilité de l'élastométrie pour la mesure de l'élasticité hépatique. *Hépato-Gastro & Oncologie Digestive* 2007;15(1):77.
2275. Pauwels, A. Le Fibroscan[®] : un nouvel outil pour l'évaluation de l'hypertension portale ? *Hépato-Gastro & Oncologie Digestive* 2007;15(1):75-76.
2276. Carrión, J. A., J. Colmenero, et al. Diagnóstico no invasivo de la fibrosis hepática. *Gastroenterología y hepatología* 2007;30(Suppl. 1):106-112.
2277. de Franchis, R. and A. Dell'Era. Non-invasive diagnosis of cirrhosis and the natural history of its complications. *Best Practice & Research Clinical Gastroenterology* 2007;21(1):3-18.
2278. de Ledinghen, V., P. Trimoulet, et al. Outbreak of hepatitis C virus infection during sclerotherapy of varicose veins: long-term follow-up of 196 patients (4535 patient-years). *Journal of Hepatology* 2007;46(1):19-25.
2279. Posthouwer, D., E. P. Mauser-Bunschoten, et al. Significant liver damage in patients with bleeding disorders and chronic hepatitis C: non-invasive assessment of liver fibrosis using transient elastography. *Journal of Thrombosis and Haemostasis* 2007;5(1):25-30.
2280. Nguyen-Khac, E. Résultats et place du FibroScan[®] dans le diagnostic non invasif de la fibrose hépatique [*Results and place of FibroScan[®] in the noninvasive diagnosis of liver fibrosis*]. *Revue de médecine interne* 2007;28(2):94-102.
2281. Nguyen-Khac, E. [Results and place of Fibroscan in the non-invasive diagnosis of hepatic fibrosis]. *La Revue de médecine interne* 2007;28(2):94-102.
2282. Abenavoli, L., C. Corpechot, et al. Elastography in hepatology. *Canadian Journal of Gastroenterology* 2007;21(12):839-42.
2283. Kang, J. K., J. Y. Cheong, et al. [Liver stiffness measurement for the diagnosis of hepatic fibrosis in patients with chronic viral hepatitis]. *Korean Journal of Hepatology* 2007;13(4):521-9.
2284. Ogawa, E., N. Furusyo, et al. Transient elastography for patients with chronic hepatitis B and C virus infection: Non-invasive, quantitative assessment of liver fibrosis. *Hepatology Research* 2007;37(12):1002-10.
2285. Stauber, R. E. and C. Lackner. Noninvasive diagnosis of hepatic fibrosis in chronic hepatitis C. *World Journal of Gastroenterology* 2007;13(32):4287-94.
2286. Talwalkar, J. A. Methodologic issues with transfer of ultrasound-based transient elastography into clinical practice. *Journal of Hepatology* 2007;47(2):301-2.
2287. Wieckowska, A., A. J. McCullough, et al. Noninvasive diagnosis and monitoring of nonalcoholic steatohepatitis: present and future. *Hepatology* 2007;46(2):582-9.
2288. Barreiro, P., S. Rodriguez-Novoa, et al. Influence of liver fibrosis stage on plasma levels of antiretroviral drugs in HIV-infected patients with chronic hepatitis C. *The Journal of Infectious Diseases* 2007;195(7):973-9.
2289. Afdhal, N. H. and M. Curry. Technology evaluation: a critical step in the clinical utilization of novel diagnostic tests for liver fibrosis. *Journal of Hepatology* 2007;46(4):543-5.
2290. Kettaneh, A., P. Marcellin, et al. Features associated with success rate and performance of FibroScan measurements for the diagnosis of

- cirrhosis in HCV patients: a prospective study of 935 patients. *Journal of Hepatology* 2007;46(4):628-34.
2291. Dib, N., A. Sawadogo, et al. [Noninvasive evaluation of liver fibrosis in patients coinfected with HIV and HCV and/or HBV]. *Journal Africain d'Hépato-Gastroentérologie* 2007;1(1):54-58.
2292. Delasalle, P. Le FibroScan Mode d'emploi et Coût. *Acta Endoscopica* 2007;37(S1):400-401.
2293. Chastagner, M., S. Hourcades, et al. [Assessment of liver fibrosis during Hepatitis C: liver needle-biopsy and non-invasive methods]. *Pratiques et organisation des soins* 2007;38(4):269-281.
2294. Fujirawa, S., Y. Hongou, et al. Relationship between liver fibrosis noninvasively measured by FibroScan and blood test. *Bulletin of the Osaka Medical College* 2007;2(53):93-105.
2295. Lamprocye, A., J. Belaiche, et al. [The FibroScan: a new non invasive method of liver fibrosis evaluation]. *Revue Médicale de Liège* 2007;62 Spec No:68-72.

2006

2296. Maida, I., V. Soriano, et al. Liver fibrosis in HIV-infected patients with chronic hepatitis B extensively exposed to antiretroviral therapy with anti-HBV activity. *HIV Clinical Trials* 2006;7(5):246-50.
2297. Abenavoli, L., L. Leggio, et al. [Evaluation of liver fibrosis in clinical practice]. *Recenti Progressi in Medicina* 2006;97(9):481-5.
2298. Guechot, J. and H. Groupe de Travail "Marqueurs Biologiques de Fibrose. [Noninvasive assessment of liver fibrosis in patients with chronic hepatitis virus C]. *La Presse Médicale* 2006;35(9 Pt 2):1317-26.
2299. Murtagh, J. and V. Foerster. Transient elastography (FibroScan) for non-invasive assessment of liver fibrosis. Issues in Emerging Health Technologies 2006;(90):1-4.
2300. Pol, S. and V. O. Mallet. Improving anti-hepatitis C virus therapy. *Expert Opinion on Biological Therapy* 2006;6(9):923-33.
2301. Stanciu, C., A. Trifan, et al. Transient elastography--an alternative to liver biopsy in patients with chronic hepatitis C? *Revista medico-chirurgicala a Societății de Medici și Naturaliști din Iasi* 2006;110(4):765-70.
2302. McHutchison, J., T. Poynard, et al. Fibrosis as an end point for clinical trials in liver disease: a report of the international fibrosis group. *Clinical Gastroenterology & Hepatology* 2006;4(10):1214-1220.
2303. Cherry, K. Accurate detection of cirrhosis by transient elastography. *Nature clinical practice. Gastroenterology & hepatology* 2006;3(5):243-244.

2304. Corpechot, C., A. El Naggar, et al. Assessment of biliary fibrosis by transient elastography in patients with PBC and PSC. *Hepatology* 2006;43(5):1118-24.
2305. Luo, J. W., J. H. Shao, et al. [Using non-invasive transient elastography for the assessment of hepatic fibrosis]. *Zhonghua Gan Zang Bing Za Zhi* 2006;14(5):395-7.
2306. Verveer, C. and R. J. de Knegt. Non-invasive measurement of liver fibrosis: application of the FibroScan in hepatology. *Scandinavian Journal of Gastroenterology* 2006;41(Suppl 243):85-88.
2307. Beaugrand, M. How to assess liver fibrosis and for what purpose? *Journal of Hepatology* 2006;44(3):444-445.
2308. Foucher, J., E. Chanteloup, et al. Diagnosis of cirrhosis by transient elastography (FibroScan): a prospective study. *Gut* 2006;55(3):403-8.
2309. Melin, P. Quel traitement après un échec? SOS Hépatites 2006;(34):10-44.
2310. Riou-Milliot, S. Un "stéthoscope" pour le foie. *Sciences et avenir* 2006;709:60-61.
2311. Soriano, V., M. Nunez, et al. Chronic hepatitis C in HIV-infected patients: those who more need therapy are those who respond less. *Infectious Disorders - Drug Targets* 2006;6(1):57-88.
2312. Lemoine, M. [Place and perspectives of diagnosis with FibroScan]. *Gastroenterologie Pratique* 2006.
2313. Sebastiani, G. and A. Alberti. Non invasive fibrosis biomarkers reduce but not substitute the need for liver biopsy. *World Journal of Gastroenterology* 2006;12(23):3682-94.
2314. Laharie, D., F. Zerbib, et al. Diagnosis of liver fibrosis by transient elastography (FibroScan) and non-invasive methods in Crohn's disease patients treated with methotrexate. *Alimentary Pharmacology & Therapeutics* 2006;23(11):1621-8.
2315. Maida, I., M. Nunez, et al. Severe liver disease associated with prolonged exposure to antiretroviral drugs. *Journal of Acquired Immune Deficiency Syndromes* 2006;42(2):177-82.
2316. Masaki, N., M. Imamura, et al. Usefulness of elastometry in evaluating the extents of liver fibrosis in hemophiliacs coinfected with hepatitis C virus and human immunodeficiency virus. *Hepatology Research* 2006;35(2):135-9.
2317. Sandrin, L., C. Fournier, et al. Fibroscan®: a noninvasive device that measures liver stiffness by transient elastography. *ITBM-RBM* 2006;27(1):25-31.
2318. Kawamoto, M., T. Mizuguchi, et al. Assessment of liver fibrosis by a noninvasive method of transient elastography and biochemical markers. *World Journal of Gastroenterology* 2006;12(27):4325-30.
2319. Moreno-Otero, R., M. Trapero-Marugan, et al. Liver fibrosis assessment by transient elastography

- in hepatitis C patients with normal alanine aminotransferase. Gut 2006;55(7):1055-6.
2320. Barreiro, P., P. Labarga, et al. Sustained virological response following HCV therapy is associated with non-progression of liver fibrosis in HCV/HIV-coinfected patients. Antiviral Therapy 2006;11(7):869-77.
2321. Mendoza, J., E. Gomez-Dominguez, et al. [Transient elastography (Fibroscan), a new non-invasive method to evaluate hepatic fibrosis]. Medicina clinica 2006;126(6):220-2.
2322. de Ledinghen, V., C. Douvin, et al. Diagnosis of hepatic fibrosis and cirrhosis by transient elastography in HIV/hepatitis C virus-coinfected patients. Journal of Acquired Immune Deficiency Syndromes 2006;41(2):175-179.
2323. Castera, L., J. Foucher, et al. FibroScan and FibroTest to assess liver fibrosis in HCV with normal aminotransferases. Hepatology 2006;43(2):373-374.
2324. Takeda, T., T. Yasuda, et al. Usefulness of noninvasive transient elastography for assessment of liver fibrosis stage in chronic hepatitis C. World Journal of Gastroenterology 2006;12(48):7768-73.
2325. Erhardt, A., J. Lorke, et al. [Transient elastography for diagnosing liver cirrhosis]. Deutsche Medizinische Wochenschrift 2006;113(49):2765-9.
2326. Carrion, J. A., M. Navasa, et al. Transient elastography for diagnosis of advanced fibrosis and portal hypertension in patients with hepatitis C recurrence after liver transplantation. Liver Transplantation 2006;12(12):1791-8.
2327. Ganne-Carrie, N., M. Ziol, et al. Accuracy of liver stiffness measurement for the diagnosis of cirrhosis in patients with chronic liver diseases. Hepatology 2006;44(6):1511-7.
2328. Nahon, P., G. Thabut, et al. Liver stiffness measurement versus clinicians' prediction or both for the assessment of liver fibrosis in patients with chronic hepatitis C. American Journal of Gastroenterology 2006;101(12):2744-2751.
2329. Nguyen-Khac, E. and D. Capron. Noninvasive diagnosis of liver fibrosis by ultrasonic transient elastography (Fibroscan). European Journal of Gastroenterology & Hepatology 2006;18(12):1321-5.
2330. Gomez-Dominguez, E., J. Mendoza, et al. Transient elastography: a valid alternative to biopsy in patients with chronic liver disease. Alimentary Pharmacology & Therapeutics 2006;24(3):513-8.
2331. Bosch, J. Predictions from a hard liver. Journal of Hepatology 2006;45(2):174-7.
2332. Corpechot, C., A. El Naggar, et al. Gender and liver: is the liver stiffness weaker in weaker sex? Hepatology 2006;44(2):513-514.
2333. Kazemi, F., A. Kettaneh, et al. Liver stiffness measurement selects patients with cirrhosis at risk of bearing large oesophageal varices. Journal of Hepatology 2006;45(2):230-5.
2334. Barreiro, P., L. Martin-Carbonero, et al. Predictors of liver fibrosis in HIV-infected patients with chronic hepatitis C virus (HCV) infection: assessment using transient elastometry and the role of HCV genotype 3. Clinical Infectious Diseases 2006;42(7):1032-9.
2335. Beaugrand, M. [Fibroscan: instructions for use]. Gastroentérologie Clinique & Biologique 2006;30(4):513-4.
2336. Foucher, J., L. Castera, et al. Prevalence and factors associated with failure of liver stiffness measurement using FibroScan in a prospective study of 2114 examinations. European Journal of Gastroenterology & Hepatology 2006;18(4):411-2.
2337. Kelleher, T. B. and N. Afdhal. Assessment of liver fibrosis in co-infected patients. Journal of Hepatology 2006;44(1 Suppl):S126-31.

2002-2005

2338. Colletta, C., C. Smirne, et al. Value of two noninvasive methods to detect progression of fibrosis among HCV carriers with normal aminotransferases. Hepatology 2005;42(4):838-45.
2339. Ghany, M. G. and E. Doo. Assessment of liver fibrosis: palpate, poke or pulse? Hepatology 2005;42(4):759-61.
2340. Sogni, P., D. Salmon-Ceron, et al. [Management of cirrhosis complications in HIV patients coinfected with hepatitis B or C virus]. La Presse Médicale 2005;34(20 Pt 2):1579-83.
2341. Kelleher, T. B. and N. Afdhal. Noninvasive assessment of liver fibrosis. Clinics in liver disease 2005;9(4):667-83, vii.
2342. Blanc, J. F., P. Bioulac-Sage, et al. Investigation of liver fibrosis in clinical practice. Hepatology Research 2005;32(1):1-8.
2343. Melin, P., A. Dacon, et al. Dépistage non invasif de la fibrose : Intérêt du FibroScan® en consultation d'alcoologie [Noninvasive screening of fibrosis: interest of FibroScan® in alcohol addiction consultation]. Alcoologie et Addictologie 2005;27(3):191-196.
2344. Ziol, M., A. Handra-Luca, et al. Noninvasive assessment of liver fibrosis by measurement of stiffness in patients with chronic hepatitis C. Hepatology 2005;41(1):48-54.
2345. Castera, L., J. Vergniol, et al. Prospective comparison of transient elastography, Fibrotest, APRI, and liver biopsy for the assessment of fibrosis in chronic hepatitis C. Gastroenterology 2005;128(2):343-50.
2346. Oberti, F. [Fibrosis diagnosis: still needs to prove itself]. Transcriptases 2005;(121).

2347. Saito, H., S. Tada, et al. Efficacy of non-invasive elastometry on staging of hepatic fibrosis. *Hepatology Research* 2004;29(2):97-103.
2348. Sandrin, L., D. Cassereau, et al. The role of the coupling term in transient elastography. *Journal of the Acoustical Society of America* 2004;115(1):73-83.
2349. Sandrin, L., B. Fourquet, et al. Transient elastography: a new noninvasive method for assessment of hepatic fibrosis. *Ultrasound in Medicine and Biology* 2003;29(12):1705-1713.
2350. Sandrin, L., M. Tanter, et al. Shear elasticity probe for soft tissues with 1-D transient elastography. *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control* 2002;49(4):436-446.
2351. Sandrin, L., M. Tanter, et al. Shear modulus imaging with 2-D transient elastography. *IEEE Transactions on Ultrasonics, Ferroelectrics, and Frequency Control* 2002;49(4):426-435.
2352. Beaugrand, M. L'élastométrie impulsionnelle : une simple "pichenette" pour mesurer la fibrose hépatique. *Revues Réseaux*.